



Corrigendum: *In vitro* Characterization of Insulin-Producing β -Cell Spheroids

Yonela Ntamo^{1,2}, Ebrahim Samodien¹, Joleen Burger^{1,3}, Nolan Muller⁴,
Christo J. F. Muller^{1,2,3} and Nireshni Chellan^{1,3*}

¹ Biomedical Research and Innovation Platform, South African Medical Research Council, Cape Town, South Africa,

² Department of Biochemistry and Microbiology, Faculty of Science and Agriculture, University of Zululand, KwaDlangezwa, South Africa, ³ Division of Medical Physiology, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa, ⁴ National Health Laboratory Service, Anatomical Pathology, Tygerberg Hospital, Cape Town, South Africa

Keywords: 3D culture spheroids, β -cell, transmission electron microscopy, insulin secretion, viability

OPEN ACCESS

Edited and reviewed by:

Allen Liu,
University of Michigan, United States

*Correspondence:

Nireshni Chellan
Nireshni.Chellan@mrc.ac.za;
nchellan@mrc.ac.za

Specialty section:

This article was submitted to
Cellular Biochemistry,
a section of the journal
Frontiers in Cell and Developmental
Biology

Received: 01 February 2021

Accepted: 15 February 2021

Published: 03 March 2021

Citation:

Ntamo Y, Samodien E, Burger J,
Muller N, Muller CJF and Chellan N
(2021) Corrigendum: *In vitro*
Characterization of Insulin-Producing
 β -Cell Spheroids.
Front. Cell Dev. Biol. 9:662574.
doi: 10.3389/fcell.2021.662574

A Corrigendum on

In vitro Characterization of Insulin-Producing β -Cell Spheroids

by Ntamo, Y., Samodien, E., Burger, J., Muller, N., Muller, C. J. F., and Chellan, N. (2021). *Front. Cell Dev. Biol.* 8:623889. doi: 10.3389/fcell.2020.623889

In the original article, there was an error. The name of the suppliers of the ProtoTissue™ bioreactors and BioArray Matrix (BAM) system were incorrectly provided as the suppliers have pointed out the error to the authors.

A correction has been made to *Materials and method section; subsection 2.1.1. Two-dimensional monolayers and three-dimensional spheroids.*

Compact, intact cell aggregates were hand-picked and transferred into ProtoTissue™ bioreactors (CelVivo ApS, DK-5491 Blommenslyst, Denmark; Cat. no 010) to facilitate the formation of spheroids. The spheroids were incubated in a rotating BioArray Matrix (BAM) system (CelVivo ApS, Denmark) at standard culture conditions for 30 days, with media exchanged every 2–3 days.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Ntamo, Samodien, Burger, Muller, Muller and Chellan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.