



OPEN ACCESS

EDITED AND REVIEWED BY
Philipp Kaldis,
Lund University, Sweden

*CORRESPONDENCE
Asish Dasgupta,
✉ asish.dasgupta@queensu.ca

RECEIVED 28 September 2023
ACCEPTED 11 October 2023
PUBLISHED 19 October 2023

CITATION

Wu D, Tian L, Hoskin V and Dasgupta A (2023), Editorial: The effects of mitochondrial dysfunction on the cell cycle.
Front. Cell Dev. Biol. 11:1303834.
doi: 10.3389/fcell.2023.1303834

COPYRIGHT

© 2023 Wu, Tian, Hoskin and Dasgupta. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Editorial: The effects of mitochondrial dysfunction on the cell cycle

Danchen Wu¹, Lian Tian², Victoria Hoskin³ and Asish Dasgupta^{1*}

¹Department of Medicine, Queen's University, Kingston, ON, Canada, ²Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, United Kingdom, ³Division of Cancer Biology and Genetics, Department of Pathology and Molecular Medicine, Queen's Cancer Research Institute, Queen's University, Kingston, ON, Canada

KEYWORDS

pulmonary arterial hypertension, cancer, mitochondrial fission, dynamin-related protein 1 (Drp1), cell cycle, diabetes, circular RNA (circRNA), cell proliferation

Editorial on the Research Topic

[The effects of mitochondrial dysfunction on the cell cycle](#)

Hyperproliferative disorders, such as pulmonary arterial hypertension (PAH) and cancers, are characterized by excessive cell proliferation and resistance to apoptosis (Dasgupta et al., 2021). This “neoplastic phenotype” is due, at least in part, to acquired changes in mitochondrial metabolism. While perturbation of mitochondrial metabolism, notably a shift to aerobic glycolysis (*Warburg phenomenon*) contributes to the proliferation/apoptosis imbalance in cells from hyperproliferative disease origin, a newly recognized abnormality, namely, dysregulation of mitochondrial dynamics has been identified (Rehman et al., 2012). Mitochondria continuously join together (fusion) and divide (fission) thereby maintaining network quality control (Mao and Klionsky, 2013), mediating cell death (Tian et al., 2017) and regulating metabolism and the cell cycle (Chen et al., 2018). The major mediator of mitochondrial fission is dynamin-related protein 1 (Drp1); while fusion is mediated by mitofusin-1 and mitofusin-2 (Archer, 2013). Upon activation, Drp1 is recruited from the cytosol to the mitochondrial outer membrane (OMM) via interaction with its receptor proteins in a poorly understood multimerization reaction. In mammals, there are four proteins on the mitochondrial outer membrane that act as Drp1 receptors: mitochondrial fission 1 (Fis1), mitochondrial fission factor (Mff), mitochondrial dynamics proteins of 49 and 51 kDa (MiD49 and MiD51, respectively) (Atkins et al., 2016). Mitotic fission coordinates mitochondrial and nuclear division, ensuring equitable distribution of mitochondria between daughter cells. Mitotic fission occurs via a Drp1-dependent process. Several studies have shown that inhibition of mitotic fission triggers a cell cycle checkpoint and results in cell cycle arrest and apoptosis (Chen et al., 2018) both in cancers and in non-malignant, hyperproliferative diseases such as PAH (Marsboom et al., 2012). Thus, mitotic fission is an appealing therapeutic target. It has also been shown that Drp1 expression is upregulated in hyperproliferative diseases and Drp1 is postrationally activated (Marsboom et al., 2012; Rehman et al., 2012; Tian et al., 2018; Abu-Hanna et al., 2023). Inhibition of Drp1 regresses cancer and PAH in animal models (Marsboom et al., 2012; Rehman et al., 2012). In this special edition of Frontiers in Cell and Developmental Biology, four journal articles on *The effects of mitochondrial dysfunction on the cell cycle* were

published: Two original research articles, one review, and one mini-review. Three articles are relevant to pulmonary hypertension (PH), and the other is related to diabetes.

The perturbation of mitochondrial dynamics leading to disordered metabolism in PAH is thoroughly discussed in a comprehensive review by Breault et al. In this review article, the authors summarized the current knowledge on the causes and consequences of disordered mitochondrial function in PAH, focusing on aberrant mitochondrial metabolism, disruption of oxygen sensing, and abnormal mitochondrial dynamics in different cell and tissues in PAH.

A mini-review by Xiao et al. provided a comprehensive summary highlighting the mechanism of mitochondrial dynamics emphasizing the role of Drp1 in the pathogenesis of PAH. This mini-review article further summarized pharmacological inhibitors that are used to target Drp1. Drp1 is a large GTPase (Archer, 2013). Following activation, Drp1 is recruited to OMM where, at a site demarcated by the endoplasmic reticulum (Friedman et al., 2011), it assembles by multimerization with its receptor proteins by hydrolyzing GTP to form a macromolecular ring-like structure. This ring constricts and divides the mitochondria. Several approaches have been taken to inhibit Drp1's ability to execute mitochondrial fission such as developing pharmacological inhibitors to inhibit Drp1's GTPase activity (Mdivi1 and Drpitor1) (Cassidy-Stone et al., 2008; Wu et al., 2020). In addition to blocking Drp1's GTPase activity, a competitive small peptide (P110) which blocks binding of Drp1 to its receptor protein Fis1 has been developed, thereby inhibiting Drp1's recruitment to OMM (Qi et al., 2013). Inhibition of Drp1's activity by mdivi-1/Drpitor1 or the Drp1-Fis1 interaction by P110 attenuated mitochondrial fission, cell proliferation, regressed cancer (Rehman et al., 2012; Wu et al., 2020) and improved right ventricular function during ischemia-reperfusion injury in experimental animal models (Tian et al., 2017). In addition, Fis1-specific inhibitor SS-31 is reported to regress PAH in a murine model (Lu et al., 2016).

In the research article entitled *Alterations in inflammatory cytokines and redox homeostasis in LPS-induced pancreatic beta-cell toxicity and mitochondrial stress: protection by azadirachtin*, the authors investigated the protective role of an anti-inflammatory, anticancer, and antioxidant phytochemical, azadirachtin (AZD) on the lipopolysaccharide (LPS)-induced nitro/oxidative stress in a rat insulin-secreting pancreatic beta cell line, Rin-5F (John and Raza). Using *in vitro* experiments, they discovered that the bacterial endotoxin LPS not only leads to nitro/oxidative stress but also triggers subsequent pro-inflammatory changes in cells, which is marked by translocation of nuclear factor kappa B (NF- κ B) and release of inflammatory cytokines, such as tumor necrosis factor alpha (TNF- α) and interleukin 6 (IL-6). Furthermore, LPS also impaired mitochondrial function in Rin-5F cells. It reduced mitochondrial membrane potential, decreased electron transport chain (ETC) complex activities, decreased ATP production, and induced mitochondria-mediated apoptosis. AZD can successfully reverse these detrimental changes in pancreatic beta cells caused by LPS.

The pancreatic beta cell produces, stores, and releases insulin, a hormone that lowers blood glucose levels (Marchetti et al., 2017). Insulin plays a crucial role in glucose homeostasis. Beta cell dysfunction is a hallmark of metabolic syndrome (MetS) and the degree of β cell dysfunction correlates with the severity of MetS (Hudish et al., 2019). In the early stage of MetS, to compensate for

the increased metabolic demand caused by changes in lifestyle, there is increased secretion of insulin in pancreatic β cells (Hudish et al., 2019). However, if the imbalance continues, β cell function will decompensate, eventually leading to a vicious loop and exacerbating MetS (Hudish et al., 2019).

Decompensated β cells in MetS and diabetes demonstrate many cellular pathological changes, including oxidative stress, inflammation, and mitochondrial dysfunction. These changes eventually lead to cell apoptosis. Increased plasma level of glucose, as seen in MetS and diabetes, induces glucose transportation across β cell membrane. As a result of increased glucose oxidative metabolism, the level of its byproduct, reactive oxygen species (ROS), is elevated. Mitochondria is the main organelle that produces ROS in β cells (Turrens, 2003). Excessive ROS exhausts the cytosolic antioxidants. In insulin-secreting cells, acute ROS attacks result in long-lasting mitochondrial inactivation, including partial loss of electron transport chain components (Li et al., 2009).

Altered mitochondrial morphology, especially fragmented and swollen mitochondrial networks are observed in diabetic animal models and β cells from type 2 diabetes (T2D) patients (Higa et al., 1999; Bindokas et al., 2003; Mizukami et al., 2008; Dlaskova et al., 2010). However, manipulating mitochondrial dynamics proteins does not always normalize β cell function (Supale et al., 2012). Since mitochondrial fission and fusion are closely related to mitophagy and calcium regulation, the involvement of mitochondrial dynamics in β cell function requires further investigation.

AZD is a chemical compound belonging to the limonoid group and an extract of fruit from the Neem tree. It acts as an antifeedant and growth disruptor to insects. The pharmaceutical use of AZD in medicine is less explored. However, preliminary studies with insects have shown that it has protective roles against oxidative stress (Zhang et al., 2018). Therefore, it may be a promising pharmacological agent for treating β cell oxidative dysfunction.

In the research article entitled *CircGSAP regulates the cell cycle of pulmonary microvascular endothelial cells via the miR-942-5p sponge in pulmonary hypertension*, Sun et al. explored the molecular mechanism for cell proliferation, apoptosis and cell cycle in pulmonary microvascular endothelial cells (PMECs). Non-coding RNAs are important molecular regulators of RNA activity and protein function and play important roles in cardiovascular, pulmonary, and muscle diseases (Bonnet et al., 2020). Circular RNAs (circRNA), a new type of non-coding RNAs, form covalently closed continuous loops and can regulate cellular functions by acting as microRNA or protein inhibitors ("sponges") (Kristensen et al., 2019; Wang et al., 2022). Many circRNAs have been found to be dysregulated in plasma or lung tissues in PH and the circRNAs are predicted to regulate the function of pulmonary arterial cells such as endothelial cell and smooth muscle cells (SMCs), resulting in pulmonary vascular remodeling in PH (Wang et al., 2022). Circular RNA gamma-secretase activating protein (circGSAP) has been found to decrease in lung tissues from idiopathic PAH patients (Yuan et al., 2021). This study demonstrates that circGSAP is downregulated in lung tissues from chronic obstructive pulmonary disease (COPD)-PH patients and in hypoxic PMECs; upregulated miR-942-5p and downregulated SMAD4 are also observed in hypoxic PMECs. The authors also showed that decreased circGSAP promotes cell proliferation, apoptosis resistance and G1/S transition and

circGSAP plays such role through competitively binding miR-942-5p to module SMAD4. These results demonstrate the important role of circGSAP/miR-942-5p/SMAD4 axis in regulating PMECs under hypoxic condition. However, no study is presented to demonstrate the role of this axis in other cell types in the lungs such as pulmonary artery SMCs and *in vivo* preclinical animal models of PH. Although this study did not investigate the mitochondrial function, loss of SMAD4 has been demonstrated to decrease mitochondrial respiration and increase mitochondrial fission in pancreatic cancer cells (Ezrova et al., 2021). In addition, numerous studies have demonstrated mitochondria are critical component in regulating cell cycle, cell proliferation, apoptosis and many cell functions in both health and diseases (Ryan et al., 2015; Dasgupta et al., 2020; Wu et al., 2021). Future work is therefore needed to identify the role of mitochondria in the pathway in both *in vitro* and *in vivo* studies.

Author contributions

DW: Writing–original draft, Writing–review and editing. LT: Writing–original draft, Writing–review and editing. VH:

Writing–review and editing. AD: Writing–original draft, Writing–review and editing.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The authors declared that they were an editorial board member of *Frontiers*, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Abu-Hanna, J., Anastasakis, E., Patel, J. A., Eddama, M. M. R., Denton, C. P., Taanman, J. W., et al. (2023). Prostacyclin mimetics inhibit DRP1-mediated pro-proliferative mitochondrial fragmentation in pulmonary arterial hypertension. *Vasc. Pharmacol.* 151, 107194. doi:10.1016/j.vph.2023.107194
- Archer, S. L. (2013). Mitochondrial dynamics--mitochondrial fission and fusion in human diseases. *N. Engl. J. Med.* 369, 2236–2251. doi:10.1056/NEJMr1215233
- Atkins, K., Dasgupta, A., Chen, K. H., Mewburn, J., and Archer, S. L. (2016). The role of Drp1 adaptor proteins MiD49 and MiD51 in mitochondrial fission: implications for human disease. *Clin. Sci. (Lond)* 130, 1861–1874. doi:10.1042/CS20160030
- Bindokas, V. P., Kuznetsov, A., Sreenan, S., Polonsky, K. S., Roe, M. W., and Philipson, L. H. (2003). Visualizing superoxide production in normal and diabetic rat islets of Langerhans. *J. Biol. Chem.* 278, 9796–9801. doi:10.1074/jbc.M206913200
- Bonnet, S., Boucherat, O., Paulin, R., Wu, D., Hindmarch, C. C. T., Archer, S. L., et al. (2020). Clinical value of non-coding RNAs in cardiovascular, pulmonary, and muscle diseases. *Am. J. Physiol. Cell. Physiol.* 318, C1–C28. doi:10.1152/ajpcell.00078.2019
- Cassidy-Stone, A., Chipuk, J. E., Ingeman, E., Song, C., Yoo, C., Kuwana, T., et al. (2008). Chemical inhibition of the mitochondrial division dynamin reveals its role in Bax/Bak-dependent mitochondrial outer membrane permeabilization. *Dev. Cell.* 14, 193–204. doi:10.1016/j.devcel.2007.11.019
- Chen, K. H., Dasgupta, A., Lin, J., Potus, F., Bonnet, S., Iremonger, J., et al. (2018). Epigenetic dysregulation of the dynamin binding partners MiD49 and MiD51 increases mitotic mitochondrial fission and promotes pulmonary arterial hypertension: mechanistic and therapeutic implications. *Circulation* 138, 287–304. doi:10.1161/CIRCULATIONAHA.117.031258
- Dasgupta, A., Chen, K. H., Lima, P. D. A., Mewburn, J., Wu, D., Al-Qazazi, R., et al. (2021). PINK1-induced phosphorylation of mitofusin 2 at serine 442 causes its proteasomal degradation and promotes cell proliferation in lung cancer and pulmonary arterial hypertension. *FASEB J. official Publ. Fed. Am. Soc. Exp. Biol.* 35, e21771. doi:10.1096/fj.202100361R
- Dasgupta, A., Wu, D., Tian, L., Xiong, P. Y., Dunham-Snary, K. J., Chen, K. H., et al. (2020). Mitochondria in the pulmonary vasculature in health and disease: oxygen-sensing, metabolism, and dynamics. *Compr. Physiol.* 10, 713–765. doi:10.1002/cphy.c190027
- Blaskova, A., Spacek, T., Santorova, J., Plecita-Hlavata, L., Berkova, Z., Saudek, F., et al. (2010). 4Pi microscopy reveals an impaired three-dimensional mitochondrial network of pancreatic islet beta-cells, an experimental model of type-2 diabetes. *Biochim. Biophys. Acta* 1797, 1327–1341. doi:10.1016/j.bbabi.2010.02.003
- Ezrova, Z., Nahacka, Z., Stursa, J., Werner, L., Vlcak, E., Kralova Viziova, P., et al. (2021). SMAD4 loss limits the vulnerability of pancreatic cancer cells to complex I inhibition via promotion of mitophagy. *Oncogene* 40, 2539–2552. doi:10.1038/s41388-021-01726-4
- Friedman, J. R., Lackner, L. L., West, M., DiBenedetto, J. R., Nunnari, J., and Voeltz, G. K. (2011). ER tubules mark sites of mitochondrial division. *Science* 334, 358–362. doi:10.1126/science.1207385
- Higa, M., Zhou, Y. T., Ravazzola, M., Baetens, D., Orci, L., and Unger, R. H. (1999). Troglitazone prevents mitochondrial alterations, beta cell destruction, and diabetes in obese prediabetic rats. *Proc. Natl. Acad. Sci. U. S. A.* 96, 11513–11518. doi:10.1073/pnas.96.20.11513
- Hudish, L. I., Reusch, J. E., and Sussel, L. (2019). β Cell dysfunction during progression of metabolic syndrome to type 2 diabetes. *J. Clin. Invest.* 129, 4001–4008. doi:10.1172/JCI129188
- Kristensen, L. S., Andersen, M. S., Stagsted, L. V. W., Ebbesen, K. K., Hansen, T. B., and Kjems, J. (2019). The biogenesis, biology and characterization of circular RNAs. *Nat. Rev. Genet.* 20, 675–691. doi:10.1038/s41576-019-0158-7
- Li, N., Brun, T., Cnop, M., Cunha, D. A., Eizirik, D. L., and Maechler, P. (2009). Transient oxidative stress damages mitochondrial machinery inducing persistent beta-cell dysfunction. *J. Biol. Chem.* 284, 23602–23612. doi:10.1074/jbc.M109.024323
- Lu, H. I., Huang, T. H., Sung, P. H., Chen, Y. L., Chua, S., Chai, H. Y., et al. (2016). Administration of antioxidant peptide SS-31 attenuates transverse aortic constriction-induced pulmonary arterial hypertension in mice. *Acta Pharmacol. Sin.* 37, 589–603. doi:10.1038/aps.2015.162
- Mao, K., and Klionsky, D. J. (2013). Participation of mitochondrial fission during mitophagy. *Cell. cycle* 12, 3131–3132. doi:10.4161/cc.26352
- Marchetti, P., Bugliani, M., De Tata, V., Suleiman, M., and Marselli, L. (2017). Pancreatic beta cell identity in humans and the role of type 2 diabetes. *Front. Cell. Dev. Biol.* 5, 55. doi:10.3389/fcell.2017.00055
- Marsboom, G., Toth, P. T., Ryan, J. J., Hong, Z., Wu, X., Fang, Y. H., et al. (2012). Dynamin-related protein 1-mediated mitochondrial mitotic fission permits hyperproliferation of vascular smooth muscle cells and offers a novel therapeutic target in pulmonary hypertension. *Circ. Res.* 110, 1484–1497. doi:10.1161/CIRCRESAHA.111.263848
- Mizukami, H., Wada, R., Koyama, M., Takeo, T., Suga, S., Wakui, M., et al. (2008). Augmented beta cell loss and mitochondrial abnormalities in sucrose-fed GK rats. *Virchows Arch.* 452, 383–392. doi:10.1007/s00428-007-0508-2
- Qi, X., Qvit, N., Su, Y. C., and Mochly-Rosen, D. (2013). A novel Drp1 inhibitor diminishes aberrant mitochondrial fission and neurotoxicity. *J. Cell. Sci.* 126, 789–802. doi:10.1242/jcs.114439
- Rehman, J., Zhang, H. J., Toth, P. T., Zhang, Y., Marsboom, G., Hong, Z., et al. (2012). Inhibition of mitochondrial fission prevents cell cycle progression in lung cancer. *FASEB J.* 26, 2175–2186. doi:10.1096/fj.11-196543
- Ryan, J., Dasgupta, A., Huston, J., Chen, K. H., and Archer, S. L. (2015). Mitochondrial dynamics in pulmonary arterial hypertension. *J. Mol. Med. Berl.* 93, 229–242. doi:10.1007/s00109-015-1263-5
- Supale, S., Li, N., Brun, T., and Maechler, P. (2012). Mitochondrial dysfunction in pancreatic beta cells. *Trends Endocrinol. Metab.* 23, 477–487. doi:10.1016/j.tem.2012.06.002

- Tian, L., Neuber-Hess, M., Mewburn, J., Dasgupta, A., Dunham-Snary, K., Wu, D., et al. (2017). Ischemia-induced Drp1 and Fis1-mediated mitochondrial fission and right ventricular dysfunction in pulmonary hypertension. *J. Mol. Med. Berl.* 95, 381–393. doi:10.1007/s00109-017-1522-8
- Tian, L., Potus, F., Wu, D., Dasgupta, A., Chen, K. H., Mewburn, J., et al. (2018). Increased drp1-mediated mitochondrial fission promotes proliferation and collagen production by right ventricular fibroblasts in experimental pulmonary arterial hypertension. *Front. Physiol.* 9, 828. doi:10.3389/fphys.2018.00828
- Turrens, J. F. (2003). Mitochondrial formation of reactive oxygen species. *J. Physiol.* 552, 335–344. doi:10.1113/jphysiol.2003.049478
- Wang, Q., Sun, Y., Zhao, Q., Wu, W., Wang, L., Miao, Y., et al. (2022). Circular RNAs in pulmonary hypertension: emerging biological concepts and potential mechanism. *Anim. Model. Exp. Med.* 5, 38–47. doi:10.1002/ame2.12208
- Wu, D., Dasgupta, A., Chen, K. H., Neuber-Hess, M., Patel, J., Hurst, T. E., et al. (2020). Identification of novel dynamin-related protein 1 (Drp1) GTPase inhibitors: therapeutic potential of Drpitor1 and Drpitor1a in cancer and cardiac ischemia-reperfusion injury. *FASEB J.* 34, 1447–1464. doi:10.1096/fj.201901467R
- Wu, D., Dasgupta, A., Read, A. D., Bentley, R. E. T., Motamed, M., Chen, K. H., et al. (2021). Oxygen sensing, mitochondrial biology and experimental therapeutics for pulmonary hypertension and cancer. *Free Radic. Biol. Med.* 170, 150–178. doi:10.1016/j.freeradbiomed.2020.12.452
- Yuan, P., Wu, W. H., Gong, S. G., Jiang, R., Zhao, Q. H., Pudasaini, B., et al. (2021). Impact of circGSAP in peripheral blood mononuclear cells on idiopathic pulmonary arterial hypertension. *Am. J. Respir. Crit. Care Med.* 203, 1579–1583. doi:10.1164/rccm.202005-2052LE
- Zhang, J., Sun, T., Sun, Z., Li, H., Qi, X., Zhong, G., et al. (2018). Azadirachtin acting as a hazardous compound to induce multiple detrimental effects in *Drosophila melanogaster*. *J. Hazard Mater* 359, 338–347. doi:10.1016/j.jhazmat.2018.07.057