



The *Escherichia coli* Serogroup O1 and O2 Lipopolysaccharides Are Encoded by Multiple O-antigen Gene Clusters

Sabine Delannoy^{1*}, Lothar Beutin^{2,3}, Patricia Mariani-Kurkdjian^{4,5,6}, Aubin Fleiss^{1†}, Stéphane Bonacorsi^{4,5,6} and Patrick Fach¹

¹ IdentyPath Platform, Food Safety Laboratory, Anses, Université Paris-Est, Maisons-Alfort, France, ² National Reference Laboratory for *Escherichia coli*, Federal Institute for Risk Assessment (BfR), Berlin, Germany, ³ Department of Biology, Chemistry, Pharmacy, Institute for Biology - Microbiology, Freie Universität Berlin, Berlin, Germany, ⁴ CNR Associé *Escherichia coli*, Service de Microbiologie, Hôpital Robert-Debré, Paris, France, ⁵ IAME, UMR 1137, INSERM, Paris, France, ⁶ IAME, UMR 1137, University Paris Diderot, Sorbonne Paris Cité, Paris, France

OPEN ACCESS

Edited by:

Nora Lía Padola,
National University of Central Buenos
Aires, Argentina

Reviewed by:

Charles Martin Dozois,
Institut National de la Recherche
Scientifique, Canada
Roxane Piazza,
Instituto Butantan, Brazil

*Correspondence:

Sabine Delannoy
sabine.delannoy@anses.fr

† Present Address:

Aubin Fleiss,
Laboratory of Computational and
Quantitative Biology, Centre National
de la Recherche Scientifique, Institut
de Biologie Paris-Seine, Sorbonne
Universités, UPMC, Paris, France

Received: 21 November 2016

Accepted: 23 January 2017

Published: 07 February 2017

Citation:

Delannoy S, Beutin L,
Mariani-Kurkdjian P, Fleiss A,
Bonacorsi S and Fach P (2017) The
Escherichia coli Serogroup O1 and O2
Lipopolysaccharides Are Encoded by
Multiple O-antigen Gene Clusters.
Front. Cell. Infect. Microbiol. 7:30.
doi: 10.3389/fcimb.2017.00030

Escherichia coli strains belonging to serogroups O1 and O2 are frequently associated with human infections, especially extra-intestinal infections such as bloodstream infections or urinary tract infections. These strains can be associated with a large array of flagellar antigens. Because of their frequency and clinical importance, a reliable detection of *E. coli* O1 and O2 strains and also the frequently associated K1 capsule is important for diagnosis and source attribution of *E. coli* infections in humans and animals. By sequencing the O-antigen clusters of various O1 and O2 strains we showed that the serogroups O1 and O2 are encoded by different sets of O-antigen encoding genes and identified potentially new O-groups. We developed qPCR-assays to detect the various O1 and O2 variants and the K1-encoding gene. These qPCR assays proved to be 100% sensitive and 100% specific and could be valuable tools for the investigations of zoonotic and food-borne infection of humans with O1 and O2 extra-intestinal (ExPEC) or Shiga toxin-producing *E. coli* (STEC) strains.

Keywords: *E. coli*, extraintestinal *E. coli*, ExPEC, Shiga toxin-producing *E. coli*, STEC, serotyping, O-antigen gene cluster, O-AGC

INTRODUCTION

Strains of the *E. coli* species are currently divided into 183 O-groups (lipopolysaccharide) and 53 H-types (flagellar antigen) (Joensen et al., 2015). For many decades serotyping of O- and H-antigens and in some cases capsular (K-antigens) was the only way to classify *E. coli* which shared similarities in their fermentation reactions, hosts, virulence markers, and their pathogenicity to humans or animals (Orskov and Orskov, 1984, 1992; Edwards and Ewing, 1986). Most of the genes responsible for the biosynthesis of the O-antigen are located in a single cluster, the O-antigen gene cluster or O-AGC (Reeves et al., 1996; Iguchi et al., 2015a; DebRoy et al., 2016). The size and gene content of the O-AGC vary between serogroups, and two genes, *wzx* (O-antigen flippase) and *wzy* (O-antigen polymerase) or *wzm* (O-antigen ABC transporter permease gene) and *wzt* (ABC transporter ATP-binding gene), appear highly specific for each serogroup. These can serve as targets for molecular determination of the serogroup by PCR, qPCR or nucleotide sequencing (Iguchi et al., 2015b; Joensen et al., 2015; Fratamico et al., 2016). The conventional serotype identification,

based on the serological agglutination of the O-antigens and flagellar H-antigens, is difficult, time consuming and expensive. In addition, many samples remain untypeable by agglutination. Indeed, cross-reactions, the existence of “O rough” strains that have lost their ability to express the O antigen, and non-motile strains that do not express a flagellar antigen, all hamper the efficiency of the method. Molecular serotyping, targeting genes involved in surface antigen synthesis, appears to be a rapid, specific and cheaper alternative to conventional serotyping.

Escherichia coli strains belonging to serogroups O1 and O2 were among the first 20 O-serogroups defined when the serotyping scheme was introduced by Kauffmann in 1944 (Orskov and Orskov, 1984). The prototype strains for the *E. coli* O1 and O2 serogroups were from human extra-intestinal infections (Orskov and Orskov, 1984). *Escherichia coli* O1 and O2 strains were since found to be associated with disease in humans and animals and also heterogeneous for their hosts, phenotypical traits and the disease they cause in humans and animals. Certain *E. coli* O1 and O2 strains such as O1:K1:H7 or O2:K1:H4, O2:K1:H5, O2:K1:H6, and O2:K1:H7 were identified as causative agents of urinary tract infection, septicaemia, and neonatal meningitis in humans and animals (Orskov, 1978; Achtman et al., 1986; Olesen et al., 1995). Both serogroups account for the majority of strains causing bacteraemia and sepsis in human patients (Orskov and Orskov, 1992; Olesen et al., 1995).

Escherichia coli O1:K1:H7, O2:K1:H4, O2:K1:H5, and O2:K1:H7 strains belong to a clonal group (sequence type 95) frequently associated with extra-intestinal diseases in humans and poultry (Dziva and Stevens, 2008; Riley, 2014). These extra-intestinal pathogenic *E. coli* (ExPEC) constitute an emerging public health problem (Johnson et al., 2012a), in particular if they carry plasmids encoding resistance to antimicrobial agents. Meanwhile, the avian pathogenic *Escherichia coli* (APEC) strains, associated with colibacillosis, a serious illness in poultry worldwide, are the cause of significant economic losses in poultry farms. Poultry strains share some ExPEC virulence traits with human O1:K1:H7 or O2:K1:H7 isolates and with other ExPEC strains frequently isolated from human patients (Johnson et al., 2006, 2007, 2012b; Dziva and Stevens, 2008; Houdouin et al., 2008; Mora et al., 2009; Peigne et al., 2009; Nandanwar et al., 2014). *Escherichia coli* O1:H6 strains form another group of ExPEC that were isolated from extra-intestinal infections of dogs, cats and horses (Ewers et al., 2014; Guo et al., 2015).

The human gut microbiota constitutes an important reservoir of *E. coli* O1:K1:H1, O1:K1:H7 and O2:K1:H4 and these strains can thus serve as source of extra-intestinal infections (Damborg et al., 2009; Moreno et al., 2009). A possible route of human contamination is through the food supply with an initial colonization of the intestinal tract before onset of an extra-intestinal infection (Manges and Johnson, 2015). Indeed, apart from poultry, many of these ExPEC strains sharing virulence markers with human isolates were isolated from a variety of sources, including animals, food, and the environment (Jakobsen et al., 2010; Singer, 2015). O2 strains such as O2:H6 or O2:H42, are frequently isolated from cattle (Diarra et al., 2009). *Escherichia coli* O2:H7 similar to isolates causing human infections were also isolated from food (Vincent et al., 2010).

Some *Escherichia coli* O1 and O2 strains producing Shiga toxins (STEC) such as O1:H20, O1:H7, O1:H2, and O1:H1 were sporadically isolated from feces of human patients with diarrhea and STEC O2:K1:H7 have caused sporadic cases of HUS in humans (Kobayashi et al., 1993; Beutin et al., 2004; Blanco et al., 2004; Chen et al., 2014). Cattle most likely serves as a reservoir for these strains as STEC O1:H20 and other STEC O1 strains were isolated from bovine meat products and cattle feces (Hussein and Bollinger, 2005; Mekata et al., 2014) while STEC O2:H27 and other STEC O2 have been recovered from feces of cattle, pigs and wild ruminants (Sánchez et al., 2009; Scott et al., 2009; Hutchinson et al., 2011). Interestingly, poultry, pigs and sheep can also harbor enteropathogenic *E. coli* (EPEC) O2 strains such as O2:H40, O2:H49, or O2:H8 (Fröhlicher et al., 2008; Alonso et al., 2016).

By analyzing different types of O1 strains, five different serological factors (O1a–O1e) were described (Moll et al., 1986) and three different polysaccharide structures (O1A, O1B, and O1C) were characterized which differ slightly in their composition (Gupta et al., 1992). The O1A type corresponded to *E. coli* O1:K1:H7 strains from human patients. The closely related O1B and O1C types were associated with other types of *E. coli* O1 strains (Achtman et al., 1983; Moll et al., 1986; Gupta et al., 1992). These differences in the LPS structures suggest that serogroup O1 strains show alterations in their O-antigen encoding genes. Moreover, serogroup O1 was reported to show multiple cross-reactions with other O-serogroups such as O2, O10, O14, O50, O53, O64, O70, O107, O115, O117, O148, O149, and O154 (Orskov and Orskov, 1984; Edwards and Ewing, 1986). The O2 antisera can also give cross-reactions with serogroups O1, O50, O53, O74, and O117 (Edwards and Ewing, 1986).

In this work, we investigated ExPEC, EPEC, and STEC strains of serogroup O1 and O2 for their O-antigen encoding genes to explore the nature of different antigens described for O1 and to explain the cross-reactivity between O1, O2, and other *E. coli* O-groups. As the nucleotide sequences of O-antigen encoding genes O1–O186 are now available (Iguchi et al., 2015a; DebRoy et al., 2016) we developed a qPCR-based detection procedure to investigate O1 and O2 strains of different H-types and from various sources. By sequencing the O-AGC of non-reacting strains, we showed that the serogroups O1 and O2 are encoded by different sets of O-antigen encoding genes and identified potentially new serotypes. The capsular antigen K1 contributes to the survival of ExPEC strains in the blood stream and is frequently associated with strains causing septicaemia and meningitis. We developed an additional qPCR assay for the specific detection of the capsular antigen K1 often associated with the clinically highly important O1 and O2 and other ExPEC strains.

MATERIALS AND METHODS

Bacteria

Escherichia coli strains used in this study were derived from the collections of the National Reference Laboratory for *E. coli* (NRL *E. coli*) at the Federal Institute for Risk Assessment (BfR) in Berlin, Germany, the associated French National Reference

Center for *E. coli* at the pediatric Hospital Robert-Debré in Paris (France), and from the French Agency for Food, Environmental and Occupational Health and Safety (Anses) in Maisons-Alfort, France.

Most of the strains studied here belong to previously described collections (Wullenweber et al., 1993; Beutin et al., 2004, 2007, 2016; Bidet et al., 2007; Bugarel et al., 2010; Martin and Beutin, 2011). *E. coli* strains used to test for the specificity of the qPCR assays included in particular the *E. coli* reference strains belonging to serogroups O1-O186 and H-types H1-H56. *E. coli* strains carrying the capsular antigen K1 (Orskov and Orskov, 1984; Wullenweber et al., 1993; Bettelheim et al., 2003) were used as reference strains for the *neuB* gene specific real-time PCR described in **Table 1**.

Two STEC O1 strains, CB13533 (O1:H33) and CB14293 (O1:H20), the *E. coli* O1:H12 strain CB11070, the O1 reference strain A47 (O1:HNT), the O50 reference strain U18-41 and two EPEC O2 strains CB11127 (O2:H49) and CB15123 (O2:H40) were investigated for the gene sequences of their O-antigen locus (**Table 2**).

All strains were grown overnight at 37°C in Luria broth, and DNA was extracted according to the manufacturer's instructions using InstaGene matrix (BioRad laboratories, Marnes-La-Coquette, France).

Serotyping of *E. coli* Strains and Detection of Capsular Antigen K1

Serotyping of O (lipopolysaccharide) and H (flagellar) antigens of the strains received at the BfR was performed with O-specific and H-specific rabbit antisera prepared according to standard methods (Orskov and Orskov, 1984). The O1 and O2 serogroups reference strains U5-41 (O1:K1:H7) and U9-41 (O2:K1:H4) (Orskov and Orskov, 1984) were used for preparation of *E. coli* O1 and O2 antisera following previously published protocols (Orskov and Orskov, 1984; Edwards and Ewing, 1986). Two-fold dilutions of O-antiserum in a range 1:50–1:12,800 were used for testing boiled cultures of bacteria for agglutination reactions. Agglutination reactions \pm one titer step as obtained with the positive control strains U5-41 and U9-41 (both end point titer 1:12,800) were interpreted as positive for serogroup O1 or O2 respectively following published serotyping methods (Orskov and Orskov, 1984; Edwards and Ewing, 1986). Strains carrying the K1 capsule were detected using K1 specific bacteriophages as previously described (Wullenweber et al., 1993).

Clinical strains isolated at the Robert Debré hospital were either serotyped by agglutination as described above or serogrouped using a multiplex PCR assay (Clermont et al., 2007). The K1 antigen was detected using the Wellcogen™ Bacterial Antigen Rapid Latex Agglutination Test according to the manufacturer's instructions.

PCR Detection and Mapping of *E. coli* O-antigen Genes

Nucleotide sequence data obtained from the previously published sequence of the lipopolysaccharide O1 and O2 antigens (Li et al., 2010) and capsular K1 antigen (Lu et al., 2011) encoding genes

TABLE 1 | Primers and probes for the real-time PCR assays.

Target gene	Forward primer, reverse primer, and probe sequences (5'–3')
<i>wzyO1A</i>	CCTTTTGTATTTCTTTCTGGCTAGTG CGAATATTTTATCCGATGGCTTTAG [6FAM]- AGTCCC GGTCATTGCTTATCGAACTGC-[BHQ1]
<i>wzxO1non-A</i>	AATCATTTGATGTCGGCATGTC GATTTTATACTTACATGGTGGATCGTATC [6FAM]-TCAGCTATACGCACTGGGCGTCCC-[BHQ1]
<i>wzxO2</i>	GCCAAGTGCAAAGTTTAAATCAACAAT CTTGCCAATTTTCCGAGTATAT [6FAM]- CCTCTGCACCTGTAAGCACTGGCCTT-[BHQ1]
<i>wzxO2-2 (CB11127)</i>	TCCGTTATATTTTATGATGGCATTGA CCCTGTTACATCCACCCCTCT [6FAM]-CAGCGCATTAGTTTTTCCACTTGCCTTG-[BHQ1]
<i>wzxO2-3 (CB15123)</i>	CCACCACGGTGCACATTTAC GGACAGGTACAAAGCCTAATGAATATT [6FAM]-TTTCCCTTTACCTCCATCCCAATTTTCTGC-[BHQ1]
<i>neuBK1</i>	TCAATAGAACCTGATGAAGTGAACAT TCTGATCATTCTAGCGGGTTTTTATG [6FAM]-TTATTCATAAAGGCACCGCCGCAA-[BHQ1]

were used for designing TaqMan® qPCR probes for specific detection of *E. coli* O1-, O2-, and K1-antigen encoding strains. As some of the investigated O1 and O2 strains did not react with the *wzyO1A* and *wzxO2* qPCR assays, the nucleotide sequence of the O-antigen encoding genes of four qPCR-negative O1 strains (A47, CB11070, CB13533, and CB14293) and two qPCR-negative O2 strains (CB11127 and CB15123) were determined (see below). A qPCR for the variant *wzx* gene present in O1 strains CB13533 and CB14293 was developed and used for further typing of the collection of *E. coli* O1 strains. In a similar way, two qPCRs for the variant *wzx* genes of CB11127 and CB15123 were developed and used for molecular typing of *E. coli* O2 strains (**Table 1**). Real-time qPCR probes and primers used in this work were designed with the software Primer Express V3.0 (Applied Biosystems) and are described in **Table 1**.

Real-time PCR assays were performed with an ABI7500 instrument (Applied Biosystems, Foster City, CA, USA) in 25- μ l reaction volumes, a Light Cycler 1536 (Roche Diagnostics, Meylan, France) in 1.5- μ l reaction volumes or a BioMark (Fluidigm, South San Francisco, CA, USA) in nanofluidic format according to the recommendations of the suppliers. Primers and TaqMan probes were used at 300 nM final concentrations. The following two-steps thermal profile was applied to all instruments: enzyme activation at 95°C for 1–10 min (as recommended depending on the enzyme used), followed by 40 cycles of denaturation at 95°C and annealing at 60°C.

TABLE 2 | Properties of the *E. coli* strains used for sequencing of the O-AGC.

Strain	Source	Year isolated or reference	Serotype (as defined by agglutination)*	stx1	stx2	eae	Accession number
CB11070	Pig feces	2007	O1:H12	–	–	–	KY115223
CB13533	Bovine feces	2011	O1:H33	+	–	–	KY115224
CB14293	veal	2012	O1:H20	+	–	–	KY115225
A47	reference strain, pig feces	Moll et al., 1986	O1:[H21]	–	–	–	KY115226
U18-41	Human urine	1944	O50:H4	–	–	–	KY115227
CB11127	pig	Fröhlicher et al., 2008	O2:H49	–	–	+	KY115228
CB15123	Chicken meat	2013	O2:H40	–	–	+	KY115229

*Serotypes between brackets were determined with a qPCR-based assay.

Nucleotide Sequencing

Genomic DNA of strains CB11070, CB13533, CB14293, A47, CB11127, CB15123, and of the O50 reference strain U18-41 was extracted from an overnight culture in tryptic soy broth (TSB) medium using the DNeasy blood and tissue kit (Qiagen). The O-antigen cluster of the strains was amplified by long range PCR between the *galF* and *gnd* genes, using the Expand Long Range PCR system (Roche) and either the 1523/1524 primers (A47, CB14293, CB13533, CB11127), 412/482 primers (CB15123), or 412/482b primers (CB11070) as described previously (Coimbra et al., 2000; Guo et al., 2004; Perelle et al., 2005). The amplicons were purified using the Charge Switch kit (Invitrogen) and used to prepare libraries with the Nextera XT kit (Illumina) according to the recommendations of the manufacturers. Paired-end short-read sequencing was performed using an Illumina MiSeq instrument (Illumina), according to the manufacturer's instruction. The raw reads were trimmed (minimum length, 35 bp; quality score, 0.03) and assembled in CLC Genomics Workbench 7.5.1 by *de novo* assembly (minimum contig length, 1000 bp), producing a single contig for each sample. The sequences were annotated using PROKKA (Seemann, 2014) on a Galaxy platform. Unannotated ORFs were further manually curated and analyzed by functional domain analysis using InterPro (Mitchell et al., 2015) and Blastp.

The annotated O-antigen cluster sequences of these strains were deposited in DDBJ/ENA/GenBank under the accession numbers listed in Table 2.

RESULTS

Analysis of the O-antigen Encoding Genes in *E. coli* O1 Strains

A qPCR was developed for the *E. coli* *wzy*O1 gene derived from the sequence of the *E. coli* O1 strain G1632 (U5-41) (Li et al., 2010; GenBank Accession GU299791; Table 1). This PCR was tested on an initial panel of 36 *E. coli* O1 strains and reacted only with 17 of these (Table 3, see below). The O1 antigen encoding genes of four non-reacting strains, A47 (O1:H-[H21]), CB11070 (O1:H12), CB13533 (O1:H33), and CB14293 (O1:H20), were analyzed by nucleotide sequencing.

The O-antigen gene clusters (O-AGC) of strains CB11070, CB13533, and CB14293 are approximately 10 kb in length; they

all contain 10 ORFs between the *galF* and *gnd* genes which all have the same transcriptional direction from *galF* to *gnd* (Figure 1, Supplementary Table S1). The overall structure of their O-AGC is similar to that of G1632 with the same genes in the same order (Figure 1). The nucleotide sequence of the *wzx* and *wzy* genes of the 3 strains are closely related (99.8–100% identity; Figure 2). All the other genes from the cluster are similarly related with an overall similarity between the three clusters of 82.5–94.1% (Supplementary Data S2). All sequences, including *wzx* and *wzy*, are however distantly related to those of the O1 reference strain G1632 (41–55% identity). Besides the *rmlBDAC* genes, which are fairly common genes among various serogroups (Iguchi et al., 2015a; DebRoy et al., 2016), the other ORFs, including the *wzx* and *wzy* gene sequences, do not have any significant similarity in the non-redundant nucleotide database of Genbank (Supplementary Table S3).

The O-AGC of strain A47 is different from the other strains (Figure 1). It contains 12 ORFs between *galF* and *gnd* in the same transcriptional order from *galF* to *gnd* (Supplementary Table S1). It is approximately 16 kb in length and has an overall structure different from G1632 reference sequence. The 5'-end of the cluster is similar to the other O1 strains, containing the *rmlBDAC* gene cluster followed by the *wzx* gene, the *mnaA* gene (encoding a UDP-N-acetylglucosamine 2-epimerase) and a glycosyl transferase. However, the second half of the cluster harbors a different set of glycosyl transferase genes. The nucleotide sequence of the *wzx* gene of strain A47 is somewhat related to that of strains CB11070, CB13533, and CB14293 as they share 89% identity (89.1–89.2%). As expected, it is distantly related to that of G1632 (54.4% identity) and no significant similarity was found in the nucleotide database of Genbank (Supplementary Table S3). Similarly, the *mnaA* gene and glycosyl transferase (PROKKA_00007) following *wzx* do not reveal any significant similarity in the nucleotide database of Genbank (Supplementary Table S1). The nucleotide sequence of the *wzy* gene however is very different from that of G1632 (44.7% identity) and to those of the 3 other strains (48% identity). A blastn search indicates that it is 89% identical to the *wzy* gene of O10:K5:H4 strain Bi8337-41. Interestingly, a blast analysis of the glycosyl transferase genes surrounding *wzy* (PROKKA_00006, PROKKA_00004, and PROKKA_00003) shows 90–94% identity to the O10 strain Bi8337-41 O-AGC (Supplementary Table S3). The last ORF of A47 O-AGC

TABLE 3 | Investigation of *E. coli* O1 strains by qPCR with O1 and K1 specific assays.

Serotype ^a	Source, origin	Reference/year isolated	Stx	Nos of strains	wzy _{O1A}	wzx _{O1non-A}	neuBK1
O1:K1:H7	See footnote b	See footnote b	–	11	+	–	+
O1:H31	Sheep feces, Norway	Döpfer et al., 2010	–	1	+	–	–
O1:[H34]	Cattle feces, Germany	This work, 2008	–	1	+	–	–
O1:H42	Human feces, France	This work, 2009	–	1	+	–	–
O1:H6	Human feces, France, Switzerland	Geser et al., 2012	–	2	+	–	–
O1:H10	Cattle feces, Germany	This work, 2011	Stx1	1	+	–	–
O1:H12	Pig feces, Germany	This work, 2007	–	1	–	+	–
O1:H20	See footnote c	See footnote c	Stx1	15	–	+	–
O1:H33	Cattle feces, Germany	This work, 2011	Stx1	1	–	+	–
O1:H19	Pig feces, Germany	Beutin et al., 1993	–	1	–	+	–
O1:NM	A47, O1non-A reference strain, pig feces	Moll et al., 1986; Gupta et al., 1992	–	1	–	+	–

^aSerotypes between brackets were determined by *fljC* gene sequencing.

^bFive O1:K1:H7 strains were from extraintestinal infections of humans (Achtman et al., 1983) including the serogroup reference strains U5-41 (Orskov and Orskov, 1984), A220 and A25 (Achtman et al., 1983; Moll et al., 1986). Four O1:K1:H7 strains were from diseased poultry and two from feces of healthy infants (Bettelheim et al., 2003).

^cThirteen of the 15 O1:H20 strains were positive for *stx*₁. One of these was also positive for *stx*₂. An additional strain was positive only for *stx*₂. Eight O1:H20 strains were from bovine meat products and isolated in Germany between 2010 and 2012. One strain was from lamb meat product and isolated in Germany in 2008. Two O1:H20 strains were feces of cattle (Germany, 2011) and three from stool of humans with diarrhea (Beutin et al., 2004). The only O1:H20 strain that was *Stx*-negative was isolated from the feces of a healthy dog (Beutin et al., 1993).

(PROKKA_00002) shows 91% identity to the *wekS* gene of O2 strain G1674 O-AGC (Supplementary Table S3).

Analysis of the O-antigen Encoding Genes in *E. coli* O2 Strains

We confirmed the published sequence of O50 reference strain U18-41 by *de novo* sequencing. The O-AGC for serogroups O2 and O50 are over 99% identical (Iguchi et al., 2015a; DebRoy et al., 2016). An alignment of O2 and O50 O-AGC sequences show that they differ by less than 10 SNPs with only 2 SNPs apparently differentiating O2 and O50 (not due to inter-individual variations or sequencing errors) which could thus be considered as a single serogroup (Supplementary Table S4).


A qPCR was developed for the *E. coli* O2/O50wzx gene derived from the sequence of the *E. coli* O2 strain G1674 (Li et al., 2010; GenBank Accession GU299792.1; Table 1). This PCR was tested on an initial panel of 23 *E. coli* O2 strains and reacted only with 13 of these (Table 4, see below). The O2 antigen encoding genes of two non-reacting strains, CB11127 (O2:H49) and CB15123 (O2:H40), were analyzed by nucleotide sequencing.

The O-AGC of both strains are different from each other and from the reference sequence G1674 (Figure 1). The O-AGC of strains CB11127 and CB15123 are approximately 15 kb in length and contain 14 and 15 ORFs between *galF* and *gnd* in the same transcriptional order from *galF* to *gnd*, respectively (Supplementary Table S1). The nucleotide sequence of the *wzx* and *wzy* genes of both strains have only 37.7–60.2% identity to each other and to the O2 and O50 reference strains (Figure 3). While O2 reference sequence G1674, O50 reference sequence U18-41 and CB15123 possess the *rmlBDAC* genes, CB11127 is missing the *rmlD* and *rmlC* genes. Both CB11127 and CB15123 also harbor the extra *manB* and *manC* genes of the GDP-mannose pathway and different sets of putative glycosyl

transferases. Transposases are inserted between the *manC* and *manB* genes of CB11127 without any apparent gene disruption. Blast analyses of CB11127 ORFs indicate that most of them (including *wzx* and *wzy*) are significantly similar to a single sequence in Genbank nucleotide collection corresponding to strain FHI98 (strain of unknown serotype isolated from human feces in 2011) (Supplementary Table S3). Besides the *rmlBDA* genes which are found in the O-AGC of multiple isolates of various serogroups, blastn analyses of CB15123 do not find any significantly similar sequences for most of its ORFs (including *wzx* and *wzy*) in Genbank nucleotide collection (Supplementary Table S3).


Development and Evaluation of qPCR Assays for Identification of *E. coli* O1, O2, and K1 Strains

Major differences between O1A and O1non-A strains were found in their corresponding *wzx* and *wzy* genes which are the principal targets used for molecular O-typing of *E. coli* (Iguchi et al., 2015a; Joensen et al., 2015). Certain sequence alterations in these two genes could be attributed to *E. coli* O1A and O1non-A strains, respectively, and qPCR assays for their specific detection were developed (Table 1). When tested with an initial panel of 36 well characterized O1 strains, the qPCR assay derived from the sequence of the ExPEC O1A strain U5-41 (*wzy*O1A) detected all O1 strains belonging to H-types H6, H7, H10, H31, H34, and H42. The qPCR assay derived from the STEC O1:H20 strain CB13533 (*wzx*O1non-A) detected all O1 strains belonging to H-types H12, H19, H20, and H33 (Table 3). We have tested the qPCRs developed for detection of O1A type and O1 non-A strains for reaction with the reference strains for serogroups O1-O186. None of the reference strains belonging to other O-groups than O1 reacted with the *wzy*O1A or with the *wzx*O1non-A qPCR assays. Additionally, they were



tested against a panel of 153 strains (Table 5), including 130 strains of serogroups other than O1, belonging to 43 different serogroups. None of the strains belonging to O-groups other than O1 reacted with the wzyO1A or with the wzxO1non-A qPCR assays. The wzyO1A qPCR assay reacted only with strains of serotype O1:K1:H7 ($n = 17$), while the wzxO1non-A qPCR assay reacted only with strains of serotype O1:H12 ($n = 1$), O1:H20 ($n = 3$), O1:H21 ($n = 1$), and O1:H33 ($n = 1$). Thus, the qPCR detection method was found to be far more specific for identification of *E. coli* O1 strains than conventional O-serotyping. Moreover, the qPCR assays used here discerned clearly between *E. coli* O1A and O1non-A type strains, respectively.

The divergence between the wzx sequences of strains U9-41 (O2:K1:H4), CB11127 (O2:H49), and CB15123 (O2:H40) allowed us to design three qPCR assays, targeting the O2/O50 and two O2-like serogroups respectively (Table 1). When tested with an initial panel of 23 well characterized O2 strains, the qPCR assay derived from the sequence of the ExPEC O2 strain U9-41 (wzxO2) detected all O2 strains belonging to H-types H1, H4, H6, H7, and H25. The qPCR assay derived from the O2:H49 strain CB11127 (wzxO2-2) detected the O2 strains belonging to H-type H49, H34, and H29. The qPCR assay derived from the O2:H40 strain CB15123 (wzxO2-3) detected all O2:H40 strains and the O2:H8 strain (Table 4). The wzxO2, wzxO2-2, and wzxO2-3 qPCRs did not react with any of the O1-O186 reference strains



except for *wzxO2* qPCR test which reacted with reference strains for serogroups O2 and O50. Additionally, when tested against the panel of 153 strains (Table 5), including 115 strains of serogroups other than O2, none of the strains belonging to O-groups other than O2 reacted with the *wzxO2*, the *wzxO2-2* or with the *wzxO2-3* qPCR assays. The *wzxO2* qPCR assay reacted only with ExPEC and STEC strains of serotype O2:H1 ($n = 2$), O2:K5:H1 ($n = 1$), O2:K1:H1 ($n = 2$), O2:K5:H4 ($n = 2$), O2:K1:H4 ($n = 3$), O2:K1:H5 ($n = 3$), O2:K1:H6 ($n = 4$), O2:K1:H7 ($n = 1$), O2:H25 ($n = 4$), O2:H27 ($n = 3$), O2:H32 ($n = 1$) and O2:K1:H48 ($n = 1$). The *wzxO2-2* qPCR assay reacted with EPEC and STEC strains of serotype O2:H34 ($n = 1$), O2:H49 ($n = 3$), and O2:H29 ($n = 2$). The *wzxO2-3* qPCR reacted only with EPEC strains of serotype O2:H8 ($n = 1$) and O2:H40 ($n = 4$).

The specificity of the *neuB* qPCR for indicating strains producing the K1 antigen was tested on the serotype reference strains O1-O186 which have been analyzed for their capsular

antigens (Orskov and Orskov, 1984; Orskov et al., 1984; Scheutz et al., 2004; Joensen et al., 2015). None of the strains except those carrying the K1 antigen (U5-41, U9-41, Bi7509-41, F11119-41, and H61) reacted with the *neuB* qPCR. The *neuB* qPCR was further evaluated on *E. coli* K1 strains belonging to different O-groups which were from clinical samples and from stool of healthy infants (Wullenweber et al., 1993). In the group of *E. coli* O1 strains, the K1 antigen gene was only detected in O1:K1:H7 strains isolated from humans and poultry (Table 3). In the group of *E. coli* O2 strains, the K1 antigen gene was detected in O2:K1:H1, O2:K1:H4, O2:K1:H5, O2:K1:H6, O2:K1:H7, O2:K1:H9, and O2:K1:H48, from humans and animals (Table 4). Among the other strains of serogroups other than O1 and O2, only four ExPEC strains isolated from humans (ONT:K1:H4, O134:K1:HNT, O166:K1:H7, and O13/O135/O129:K1:H4) reacted with the K1 qPCR assay

TABLE 4 | Investigation of *E. coli* O2 strains by qPCR with O2 and K1 specific assays.

Serotype	Source, origin	References	Stx	Nos of strains	wzxO2	wzxO2-2	wzxO2-3	neuBK1
O2:K1:H6	Infant feces, Germany	Wullenweber et al., 1993; Bettelheim et al., 2003	–	1	+	–	–	+
O2:K1:H7	Infant feces, Germany	Wullenweber et al., 1993; Bettelheim et al., 2003	–	1	+	–	–	+
O2: H1	Calf feces, Germany	This work, 2010	–	1	+	–	–	–
O2:K1:H4	Infant feces, Germany	Wullenweber et al., 1993; Bettelheim et al., 2003	–	1	+	–	–	+
O2:K5:H4	Infant feces, Germany	Wullenweber et al., 1993; Bettelheim et al., 2003	–	2	+	–	–	–
O2:H25	See footnote a	See footnote a	+	3	+	–	–	–
O2:H32	Wild boar, Germany	This work, 2011	stx2	1	+	–	–	–
O2:H27	Cattle feces, Germany	This work, 2011	stx2	2	+	–	–	–
O2:K2:H1	Infant feces, Germany	Wullenweber et al., 1993; Bettelheim et al., 2003	–	1	+	–	–	–
O2:H40	See footnote b	See footnote b	–	3	–	–	+	–
O2:H34	Cat feces, Brazil	Morato et al., 2009	–	1	–	+	–	–
O2:H49	Pig feces, Switzerland	Fröhlicher et al., 2008	–	3	–	+	–	–
O2:H29	Cattle feces	Martin and Beutin, 2011	stx2	2	–	+	–	–
O2:H8	Sheep feces, Germany	Krause et al., 2005	–	1	–	–	+	–

^aOne stx2-positive O2:H25 strain was isolated from calf feces (Germany, 2011). One stx2-positive O2:H25 strain is a stx2g reference strain (Scheutz et al., 2012). The only stx1-positive O2:H25 strain was isolated from cattle (Beutin et al., 1993).

^bO2:H40 strains were isolated from pig and sheep feces, (2007) (Fröhlicher et al., 2008) and chicken meat (this work, 2013).

(Table 5). These four strains were previously found K1-positive by latex agglutination.

DISCUSSION


Escherichia coli strains of serogroups O1 and O2 belong to the most frequently isolated *E. coli* types from humans (Orskov and Orskov, 1992; Olesen et al., 1995; Bettelheim et al., 2003; Ciesielczuk et al., 2016) and also have a broad reservoir in pets and domestic animals (Moulin-Schouleur et al., 2007; Dziva and Stevens, 2008; Ewers et al., 2014). Because of their incidence, clinical importance and type diversity, a reliable identification of *E. coli* O1 and O2 serogroups together with the frequently associated K1 capsule is important for diagnosis and source attribution of *E. coli* infections in humans and animals. Detection of *E. coli* O1 and O2 strains by conventional serotyping with O-antigen specific rabbit antisera is hampered by cross reactions that were found between *E. coli* O1 and O2 and also with many other *E. coli* O-serogroups (Orskov and Orskov, 1984; Edwards and Ewing, 1986). Detection of the K1 capsule is made either by serotyping or by lysotyping with K1-specific bacteriophages (Gross et al., 1977; Orskov et al., 1977). Both methods are costly and rather difficult to introduce in routine clinical diagnosis and would be advantageously replaced by molecular methods.

The LPS of *E. coli* O1 strains exhibit some structural variations which cross-react in conventional serotyping assays. The common epitope between the O1 variants contains an *N*-acetylmannosamine residue branched on an *L*-rhamnose residue (Gupta et al., 1992). Moreover, *E. coli* O1 antisera were reported to show multiple cross-reactions with other O-serogroups. With the O1 antiserum used here for serotyping, serological cross-reactions with reference strains for O-serogroups O2, O10, O50, O53, O107, O115, O117, O148, O149, and O154 were observed,

corresponding to previously published findings (Orskov and Orskov, 1984; Edwards and Ewing, 1986). All of these O-groups possess the *rmlBDAC* genes (Iguchi et al., 2015a; DebRoy et al., 2016) involved in the synthesis of *L*-rhamnose residues which must be part of their antigenic epitope and may be the cause of the serological cross-reactions. Similarly, O2 antisera can show cross-reactions with serogroups O1, O2/O50, O53, O74, and O117 (Orskov et al., 1977). The O2 antiserum used here agglutinated strains belonging to these O-groups as well as O184 which was not in the serotyping scheme published by Orskov in 1977 (Orskov et al., 1977). The molecular basis of these serological cross-reactions is not clear as the O-AGC of these serogroups exhibit few common features.

In an effort to build a complete molecular serotyping scheme for *E. coli* we sought to develop qPCR assays for the specific detection of O1, O2 LPS and the K1 capsule. Based on the nucleotide sequence of the *wzy* gene of the O1 reference strain G1632 (U5-41) we developed a qPCR assay for specific detection of O1 strains. While this assay appeared highly specific, as it did not cross react with any other serogroups, several strains identified as O1 with the seroagglutination failed to react. In the same way, a highly specific qPCR assay developed based on the *wzx* gene of the O2 reference strain G1674 failed to react with several strains identified as O2 with the seroagglutination. The qPCR assay developed for specific detection of the *neuB* gene of K1 strains proved 100% sensitive and 100% specific.

Analysis of the O-AGC of non-reacting O1 and O2 strains revealed significant nucleotide sequence divergences. The O-AGC in the non-reacting O1 strains CB11070, CB13533, and CB14293 appears to be a variant of the one in O1A reference strain G1632. The O1A polysaccharide contains five sugar residues: 3 *L*-rhamnose residues, an *N*-acetylglucosamine residue and an *N*-Acetyl-*D*-mannosamine residue. The



O1B/O1C variants contain two *L*-rhamnose residues, an *N*-acetylglucosamine residue, an *N*-Acetyl-*D*-mannosamine residue and a *D*-Galactose residue (Moll et al., 1986; Gupta et al., 1992). Strains CB11070, CB13533, and CB14293 all apparently conserved the ability to synthesize *L*-rhamnose residues, *N*-acetylglucosamine residues and *N*-Acetyl-*D*-mannosamine residues. The presence of a *D*-Galactose residue in these strains cannot be inferred from the O-AGC sequence. Presence of other residues whose biosynthesis genes are located outside of the O-AGC cannot be ruled out. Indeed, genes involved in the biosynthesis of the nucleotide precursors of common sugars are usually located outside O-antigen gene clusters (Stenutz et al., 2006). Additional LPS-modifying enzymes can also be found elsewhere in the chromosome (Wang et al., 2002), especially in bacteriophages (Allison and Verma, 2000; Perry et al., 2009). The differences in genetic sequence of the whole cluster suggest a clonal divergence and

parallel evolution of O1A and O1nonA strains. The genetic structure of the O-AGC of strain A47 appears to be a mosaic structure containing three modules. The first module (in the transcriptional orientation), containing seven genes including *wzx*, derives from the same O1 variant. The second module, containing four genes including *wzy*, is related to the O10 O-AGC. The third module contains a single gene derived from the O2-OAGC. This mosaic structure suggests a recombination between the O-AGC of strains of serogroups O1, O10 and O2. Generation of new serogroup variants through recombination of parts of the O-AGC has already been reported for serogroup O9 (Sugiyama et al., 1998) and could be used to generate new epitopes and evade detection by the immune system of the host.

The O-AGC found in O2 strains CB11127 and CB15123 constitute two new O2-like serogroups. The genetic structure of the O-AGCs in strains CB11127 and CB15123 are different from

TABLE 5 | *E. coli* strains tested by real-time PCR.

Group (n° of strains)	Serotypes (n° of strains)
O1 (n = 23)	O1:K1:H7 (n = 17); O1:[H21]; O1:H20 (n = 3); O1:H33; O1:H12
O2 (n = 38)	O2:H32; O2:H25 (n = 4); O2:K5:H4 (n = 2); O2:K1:H4; O2:K2:H1; O2:H1; O2:K5:H1; O2:H27 (n = 3); O2:K1:H6; O2:K1:H7; O2:K1:H4 (n = 2); O2:K1:H1 (n = 2); O2:K1:H5 (n = 3); O2:K1:H6 (n = 3); O2:K1:H48; O2:H34; O2:H49 (n = 3); O2:H29 (n = 2); O2:H8; O2:H40 (n = 4)
K1 (non O1, non O2) (n = 4)	ONT:K1:H4; [O134]:K1:HNT; [O166]:K1:H7; [O13/O135/O129]:K1:H4
Other (n = 88)	O4:H5; O4:H16; O6 (n = 2); O6:H4; O6:H10; O6:H34 (n = 2); O7:H4; O8:H45; O9:H12 (n = 2); O15:H2; O15:H11; O15:H16; O15:H21; O17/77:H41; O20:H9; O20:H30; O20:H33; O20:NM; O23:H15; O28:H28; O33:H12; O40:H6; O41:H7; O45:H20; O45:H31; O51:H49; O55:H6; O55:H7 (n = 3); O55:H19; O55:H21; O55:[H51]; O63:H6; O68:H12; O76:H41; O86:H34; O91:H28; O103:H2; O110:H28; O113:H6 (n = 2); O114:H49; O118:H2; O118:H5; O118:H8; O118:H12; O118:H16; O121:H45; O125:H6; O126:H6; O127:H6; O128:H2; O128:H8; O132:H1; O132:H34; O136:H2; O136:H12; O136:NM; O139:H1; O139:H4; O139:H19; O139:H+; O139; O142:H34; O145:H28; O145:H34; O153:H7; O153:H12; O153:H14; O153:H21; O153:H25; O157:H7 (n = 4); O157:H12 (n = 2); O157:H15; O157:H45; O179:H31; O181:H49; O186:[H45]; OX185:H28; Or:H12

For each serotype, n = 1 unless otherwise stated. Serotypes between brackets were determined by sequencing.

that of the O2 reference strain G1674 suggesting different LPS structures. The O2 polysaccharide contains three L-rhamnose residues, an N-acetylglucosamine residue, and an N-acetyl-D-galactosamine (Jansson et al., 1987). Strain CB11127 most likely lacks the L-rhamnose residues as the absence of the *rmlC* and *rmlD* genes prevents full transformation of glucose-1-phosphate into dTDP-L-Rhamnose. Instead, presence of the *manB* and *manC* genes suggests the presence of residue(s) derived from GDP-D-Mannose. Presence of *rmlA*, *rmlB*, *fdtA*, *fdtB*, and *fdtC* suggest that the N-acetyl-D-galactosamine residue is conserved. Strain CB15123 conserved both the ability to synthesize L-rhamnose residues and N-acetyl-D-galactosamine residues and also acquired the GDP-D-Mannose pathway. Once again, presence of other residues whose biosynthesis genes are located outside of the O-AGC cannot be ruled out.

Based on the nucleotide sequence of the new O1-variant and the two O2-like serogroups, specific qPCR assays were developed, which proved to be 100% sensitive and 100% specific, and able to clearly distinguish the different genetic variants identified within each serogroup.

Overall, the various qPCR assays designed in this study can be used for detection and rapid identification of the different types of *E. coli* strains of serogroups O1 and O2. They provide a significantly improved alternative to classical O-serotyping by their high specificity and sensitivity to detect virtually all types of O1 and O2 strains and their respective associated variants. They are therefore potentially useful for direct identification of clinically important O1 and O2 strains in food and clinical specimens. These different qPCR assays can thus serve as tools for epidemiological investigations and source attribution of human infections. Molecular typing by use of specific qPCRs provides a more specific detection of clinically important O1 and O2 ExPEC strains. Ambiguous results obtained by serotyping such as cross-reactions with other *E. coli* O-groups and agglutination reactions

with O-rough strains are avoided. Moreover, the qPCR approach described in this study provides a better discrimination between O1 and O2 ExPEC and non-ExPEC strains. In combination with the detection of the *neuB* gene (K1 capsule), life-threatening O1 and O2 ExPEC infections in risk groups such as newborn infants can be more rapidly identified. Asymptomatic carriers, animals and food contaminated with O1 and O2 ExPEC strains can be more easily identified and the spread of pathogens to human risk groups (new born-infants, elderly, and intensive-care patients) can thus be prevented. The qPCR detection of ExPEC O1 and O2 strains is a reliable and easy approach to elucidate the significance of zoonotic and food-borne infection of humans with ExPEC O1 and O2 strains in epidemiological investigations.

AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: LB, SD, and PF. Performed the experiments: SD, LB, PM, AF, and SB. Analyzed the data: SD, LB, PM, AF, SB, and PF. Contributed reagents/materials/analysis tools: SD, LB, PM, SB, and PF. Wrote the paper: SD, LB, and PF. Critical revision of the paper for important intellectual content: SD, LB, PM, AF, SB, and PF.

ACKNOWLEDGMENTS

We thank Yannick Marie from the iGenSeq Genotyping and Sequencing Core facility at the ICM Brain and Spine Institute for generating the library and sequencing the O-AGC of strain U18-41 (O50).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fcimb.2017.00030/full#supplementary-material>

REFERENCES

- Achtman, M., Heuzenroeder, M., Kusecek, B., Ochman, H., Caugant, D., Selander, R. K., et al. (1986). Clonal analysis of *Escherichia coli* O2:K1 isolated from diseased humans and animals. *Infect. Immun.* 51, 268–276.
- Achtman, M., Mercer, A., Kusecek, B., Pohl, A., Heuzenroeder, M., Aaronson, W., et al. (1983). Six widespread bacterial clones among *Escherichia coli* K1 isolates. *Infect. Immun.* 39, 315–335.
- Allison, G. E., and Verma, N. K. (2000). Serotype-converting bacteriophages and O-antigen modification in *Shigella flexneri*. *Trends Microbiol.* 8, 17–23. doi: 10.1016/S0966-842X(99)01646-7
- Alonso, M. Z., Sanz, M. E., Irino, K., Krüger, A., Lucchesi, P. M., and Padola, N. L. (2016). Isolation of atypical enteropathogenic *Escherichia coli* from chicken and chicken-derived products. *Br. Poult. Sci.* 57, 161–164. doi: 10.1080/00071668.2015.1135502
- Bettelheim, K. A., Beutin, L., Gleier, K., Pearce, J. L., Luke, R. K., and Zimmermann, S. (2003). Serotypes of *Escherichia coli* isolated from healthy infants in Berlin, Germany and Melbourne, Australia. *Comp. Immunol. Microbiol. Infect. Dis.* 26, 55–63. doi: 10.1016/S0147-9571(02)00015-2
- Beutin, L., Delannoy, S., and Fach, P. (2016). Genetic analysis and detection of *fliC_{H11}* and *fliC_{H12}* genes coding for serologically closely related flagellar antigens in human and animal pathogenic *Escherichia coli*. *Front. Microbiol.* 7:135. doi: 10.3389/fmicb.2016.00135
- Beutin, L., Geier, D., Steinrück, H., Zimmermann, S., and Scheutz, F. (1993). Prevalence and some properties of verotoxin (Shiga-like toxin)-producing *Escherichia coli* in seven different species of healthy domestic animals. *J. Clin. Microbiol.* 31, 2483–2488.
- Beutin, L., Krause, G., Zimmermann, S., Kaulfuss, S., and Gleier, K. (2004). Characterization of Shiga toxin-producing *Escherichia coli* strains isolated from human patients in Germany over a 3-year period. *J. Clin. Microbiol.* 42, 1099–1108. doi: 10.1128/JCM.42.3.1099-1108.2004
- Beutin, L., Miko, A., Krause, G., Pries, K., Haby, S., Steege, K., et al. (2007). Identification of human-pathogenic strains of Shiga toxin-producing *Escherichia coli* from food by a combination of serotyping and molecular typing of Shiga toxin genes. *Appl. Environ. Microbiol.* 73, 4769–4775. doi: 10.1128/AEM.00873-07
- Bidet, P., Mahjoub-Messai, F., Blanco, J., Blanco, J., Dehem, M., Aujard, Y., et al. (2007). Combined multilocus sequence typing and O serogrouping distinguishes *Escherichia coli* subtypes associated with infant urosepsis and/or meningitis. *J. Infect. Dis.* 196, 297–303. doi: 10.1086/518897
- Blanco, J. E., Blanco, M., Alonso, M. P., Mora, A., Dahbi, G., Coira, M. A., et al. (2004). Serotypes, virulence genes, and intimin types of Shiga toxin (verotoxin)-producing *Escherichia coli* isolates from human patients: prevalence in Lugo, Spain, from 1992 through 1999. *J. Clin. Microbiol.* 42, 311–319. doi: 10.1128/JCM.42.1.311-319.2004
- Bugarel, M., Beutin, L., and Fach, P. (2010). Low-density microarray targeting non-locus of enterocyte effacement effectors (nle genes) and major virulence factors of Shiga toxin-producing *Escherichia coli* (STEC): a new approach for molecular risk assessment of STEC isolates. *Appl. Environ. Microbiol.* 76, 203–211. doi: 10.1128/AEM.01921-09
- Chen, Y., Chen, X., Zheng, S., Yu, F., Kong, H., Yang, Q., et al. (2014). Serotypes, genotypes and antimicrobial resistance patterns of human diarrhoeagenic *Escherichia coli* isolates circulating in southeastern China. *Clin. Microbiol. Infect.* 20, 52–58. doi: 10.1111/1469-0691.12188
- Ciesielczuk, H., Jenkins, C., Chattaway, M., Doumith, M., Hope, R., Woodford, N., et al. (2016). Trends in ExPEC serogroups in the UK and their significance. *Eur. J. Clin. Microbiol. Infect. Dis.* 35, 1661–1666. doi: 10.1007/s10096-016-2707-8
- Clermont, O., Johnson, J. R., Menard, M., and Denamur, E. (2007). Determination of *Escherichia coli* O types by allele-specific polymerase chain reaction: application to the O types involved in human septicemia. *Diagn. Microbiol. Infect. Dis.* 57, 129–136. doi: 10.1016/j.diagmicrobio.2006.08.007
- Coimbra, R. S., Grimont, F., Lenormand, P., Burguière, P., Beutin, L., and Grimont, P. A. (2000). Identification of *Escherichia coli* O-serogroups by restriction of the amplified O-antigen gene cluster (rfb-RFLP). *Res. Microbiol.* 151, 639–654. doi: 10.1016/S0923-2508(00)00134-0
- Damborg, P., Nielsen, S. S., and Guardabassi, L. (2009). *Escherichia coli* shedding patterns in humans and dogs: insights into within-household transmission of phylotypes associated with urinary tract infections. *Epidemiol. Infect.* 137, 1457–1464. doi: 10.1017/S095026880900226X
- DeRoy, C., Fratamico, P. M., Yan, X., Baranzoni, G., Liu, Y., Needleman, D. S., et al. (2016). Comparison of O-Antigen gene clusters of all O-Serogroups of *Escherichia coli* and proposal for adopting a new nomenclature for O-Typing. *PLoS ONE* 11:e0147434. doi: 10.1371/journal.pone.0147434
- Diarra, M. S., Giguere, K., Malouin, F., Lefebvre, B., Bach, S., Delaquis, P., et al. (2009). Genotype, serotype, and antibiotic resistance of sorbitol-negative *Escherichia coli* isolates from feedlot cattle. *J. Food Prot.* 72, 28–36. doi: 10.4315/0362-028X-72.1.28
- Döpfer, D., Sekse, C., Beutin, L., Solheim, H., Van Der Wal, F. J., De Boer, A., et al. (2010). Pathogenic potential and horizontal gene transfer in ovine gastrointestinal *Escherichia coli*. *J. Appl. Microbiol.* 108, 1552–1562. doi: 10.1111/j.1365-2672.2009.04575.x
- Dziva, F., and Stevens, M. P. (2008). Colibacillosis in poultry: unravelling the molecular basis of virulence of avian pathogenic *Escherichia coli* in their natural hosts. *Avian Pathol.* 37, 355–366. doi: 10.1080/03079450802216652
- Edwards, P. R., and Ewing, W. H. (1986). *Edwards and Ewing's Identification of Enterobacteriaceae*. New York, NY: Elsevier Science Publishing Co. Inc.
- Ewers, C., Bethe, A., Stamm, I., Grobbel, M., Kopp, P. A., Guerra, B., et al. (2014). CTX-M-15-D-ST648 *Escherichia coli* from companion animals and horses: another pandemic clone combining multiresistance and extraintestinal virulence? *J. Antimicrob. Chemother.* 69, 1224–1230. doi: 10.1093/jac/dkt516
- Fratamico, P. M., Debroy, C., Liu, Y., Needleman, D. S., Baranzoni, G. M., and Feng, P. (2016). Advances in molecular serotyping and subtyping of *Escherichia coli*. *Front. Microbiol.* 7:644. doi: 10.3389/fmicb.2016.00644
- Fröhlicher, E., Krause, G., Zweifel, C., Beutin, L., and Stephan, R. (2008). Characterization of attaching and effacing *Escherichia coli* (AEEC) isolated from pigs and sheep. *BMC Microbiol.* 8:144. doi: 10.1186/1471-2180-8-144
- Geser, N., Stephan, R., Korczak, B. M., Beutin, L., and Hachler, H. (2012). Molecular identification of extended-spectrum- β -lactamase genes from Enterobacteriaceae isolated from healthy human carriers in Switzerland. *Antimicrob. Agents Chemother.* 56, 1609–1612. doi: 10.1128/AAC.05539-11
- Gross, R. J., Cheasty, T., and Rowe, B. (1977). Isolation of bacteriophages specific for the K1 polysaccharide antigen of *Escherichia coli*. *J. Clin. Microbiol.* 6, 548–550.
- Guo, H., Feng, L., Tao, J., Zhang, C., and Wang, L. (2004). Identification of *Escherichia coli* O172 O-antigen gene cluster and development of a serogroup-specific PCR assay. *J. Appl. Microbiol.* 97, 181–190. doi: 10.1111/j.1365-2672.2004.02305.x
- Guo, S., Wakeham, D., Brouwers, H. J., Cobbold, R. N., Abraham, S., Mollinger, J. L., et al. (2015). Human-associated fluoroquinolone-resistant *Escherichia coli* clonal lineages, including ST354, isolated from canine feces and extraintestinal infections in Australia. *Microbes Infect.* 17, 266–274. doi: 10.1016/j.micinf.2014.12.016
- Gupta, D. S., Shashkov, A. S., Jann, B., and Jann, K. (1992). Structures of the O1B and O1C lipopolysaccharide antigens of *Escherichia coli*. *J. Bacteriol.* 174, 7963–7970. doi: 10.1128/jb.174.24.7963-7970.1992
- Houdouin, V., Bonacorsi, S., Bidet, P., Blanco, J., De La Rocque, F., Cohen, R., et al. (2008). Association between mortality of *Escherichia coli* meningitis in young infants and non-virulent clonal groups of strains. *Clin. Microbiol. Infect.* 14, 685–690. doi: 10.1111/j.1469-0691.2008.02019.x
- Hussein, H. S., and Bollinger, L. M. (2005). Prevalence of Shiga toxin-producing *Escherichia coli* in beef cattle. *J. Food Prot.* 68, 2224–2241. doi: 10.4315/0362-028X-68.10.2224
- Hutchinson, J. P., Cheney, T. E., Smith, R. P., Lynch, K., and Pritchard, G. C. (2011). Verocytotoxin-producing and attaching and effacing activity of *Escherichia coli* isolated from diseased farm livestock. *Vet. Rec.* 168:536. doi: 10.1136/vr.d915
- Iguchi, A., Iyoda, S., Kikuchi, T., Ogura, Y., Katsura, K., Ohnishi, M., et al. (2015a). A complete view of the genetic diversity of the *Escherichia coli* O-antigen biosynthesis gene cluster. *DNA Res.* 22, 101–107. doi: 10.1093/dnares/dsu043
- Iguchi, A., Iyoda, S., Seto, K., Morita-Ishihara, T., Scheutz, F., Ohnishi, M., et al. (2015b). *Escherichia coli* O-Genotyping PCR: a comprehensive and practical platform for molecular O Serogrouping. *J. Clin. Microbiol.* 53, 2427–2432. doi: 10.1128/JCM.00321-15

- Jakobsen, L., Spangholm, D. J., Pedersen, K., Jensen, L. B., Emborg, H. D., Agerso, Y., et al. (2010). Broiler chickens, broiler chicken meat, pigs and pork as sources of ExPEC related virulence genes and resistance in *Escherichia coli* isolates from community-dwelling humans and UTI patients. *Int. J. Food Microbiol.* 142, 264–272. doi: 10.1016/j.ijfoodmicro.2010.06.025
- Jansson, P. E., Lennholm, H., Lindberg, B., Lindquist, U., and Svenson, S. B. (1987). Structural studies of the O-specific side-chains of the *Escherichia coli* O2 lipopolysaccharide. *Carbohydr. Res.* 161, 273–279. doi: 10.1016/S0008-6215(00)90084-3
- Joensen, K. G., Tetzschner, A. M., Iguchi, A., Aarestrup, F. M., and Scheutz, F. (2015). Rapid and easy in silico serotyping of *Escherichia coli* isolates by use of whole-genome sequencing data. *J. Clin. Microbiol.* 53, 2410–2426. doi: 10.1128/JCM.00008-15
- Johnson, J. R., Owens, K. L., Clabots, C. R., Weissman, S. J., and Cannon, S. B. (2006). Phylogenetic relationships among clonal groups of extraintestinal pathogenic *Escherichia coli* as assessed by multi-locus sequence analysis. *Microbes Infect.* 8, 1702–1713. doi: 10.1016/j.micinf.2006.02.007
- Johnson, T. J., Kariyawasam, S., Wannemuehler, Y., Mangiamale, P., Johnson, S. J., Doetkott, C., et al. (2007). The genome sequence of avian pathogenic *Escherichia coli* strain O1:K1:H7 shares strong similarities with human extraintestinal pathogenic *E. coli* genomes. *J. Bacteriol.* 189, 3228–3236. doi: 10.1128/JB.01726-06
- Johnson, T. J., Logue, C. M., Johnson, J. R., Kuskowski, M. A., Sherwood, J. S., Barnes, H. J., et al. (2012a). Associations between multidrug resistance, plasmid content, and virulence potential among extraintestinal pathogenic and commensal *Escherichia coli* from humans and poultry. *Foodborne Pathog. Dis.* 9, 37–46. doi: 10.1089/fpd.2011.0961
- Johnson, T. J., Wannemuehler, Y., Kariyawasam, S., Johnson, J. R., Logue, C. M., and Nolan, L. K. (2012b). Prevalence of avian-pathogenic *Escherichia coli* strain O1 genomic islands among extraintestinal and commensal *E. coli* isolates. *J. Bacteriol.* 194, 2846–2853. doi: 10.1128/JB.06375-11
- Kobayashi, Y., Maeyama, K., Shiki, Y., Tsukamoto, T., Nagai, T., Peina, Y., et al. (1993). [Sporadic case of hemolytic uremic syndrome associated with verotoxin producing *Escherichia coli* O2:K1:H7]. *Kansenshogaku Zasshi* 67, 116–121. doi: 10.11150/kansenshogakuzasshi1970.67.116
- Krause, G., Zimmermann, S., and Beutin, L. (2005). Investigation of domestic animals and pets as a reservoir for intimin- (*eae*) gene positive *Escherichia coli* types. *Vet. Microbiol.* 106, 87–95. doi: 10.1016/j.vetmic.2004.11.012
- Li, D., Liu, B., Chen, M., Guo, D., Guo, X., Liu, F., et al. (2010). A multiplex PCR method to detect 14 *Escherichia coli* serogroups associated with urinary tract infections. *J. Microbiol. Methods* 82, 71–77. doi: 10.1016/j.mimet.2010.04.008
- Lu, S., Zhang, X., Zhu, Y., Kim, K. S., Yang, J., and Jin, Q. (2011). Complete genome sequence of the neonatal-meningitis-associated *Escherichia coli* strain CE10. *J. Bacteriol.* 193:7005. doi: 10.1128/jb.06284-11
- Manges, A. R., and Johnson, J. R. (2015). Reservoirs of extraintestinal pathogenic *Escherichia coli*. *Microbiol. Spectr.* 3. doi: 10.1128/microbiolspec.UTI-0006-2012
- Martin, A., and Beutin, L. (2011). Characteristics of Shiga toxin-producing *Escherichia coli* from meat and milk products of different origins and association with food producing animals as main contamination sources. *Int. J. Food Microbiol.* 146, 99–104. doi: 10.1016/j.ijfoodmicro.2011.01.041
- Mekata, H., Iguchi, A., Kawano, K., Kirino, Y., Kobayashi, I., and Misawa, N. (2014). Identification of O serotypes, genotypes, and virulotypes of Shiga toxin-producing *Escherichia coli* isolates, including non-O157 from beef cattle in Japan. *J. Food Prot.* 77, 1269–1274. doi: 10.4315/0362-028X.JFP-13-506
- Mitchell, A., Chang, H. Y., Daugherty, L., Fraser, M., Hunter, S., Lopez, R., et al. (2015). The InterPro protein families database: the classification resource after 15 years. *Nucleic Acids Res.* 43, D213–221. doi: 10.1093/nar/gku1243
- Moll, A., Kusecek, B., Pluschke, G., Morelli, G., Kamke, M., Jann, B., et al. (1986). A reexamination of the O1 lipopolysaccharide antigen group of *Escherichia coli*. *Infect. Immun.* 53, 257–263.
- Mora, A., López, C., Dabhi, G., Blanco, M., Blanco, J. E., Alonso, M. P., et al. (2009). Extraintestinal pathogenic *Escherichia coli* O1:K1:H7/NM from human and avian origin: detection of clonal groups B2 ST95 and D ST59 with different host distribution. *BMC Microbiol.* 9:132. doi: 10.1186/1471-2180-9-132
- Morato, E. P., Leomil, L., Beutin, L., Krause, G., Moura, R. A., and Pestana De Castro, A. F. (2009). Domestic cats constitute a natural reservoir of human enteropathogenic *Escherichia coli* types. *Zoonoses Public Health* 56, 229–237. doi: 10.1111/j.1863-2378.2008.01190.x
- Moreno, E., Johnson, J. R., Pérez, T., Prats, G., Kuskowski, M. A., and Andreu, A. (2009). Structure and urovirulence characteristics of the fecal *Escherichia coli* population among healthy women. *Microbes Infect.* 11, 274–280. doi: 10.1016/j.micinf.2008.12.002
- Moulin-Schouleur, M., Répérant, M., Laurent, S., Brée, A., Mignon-Grasteau, S., Germon, P., et al. (2007). Extraintestinal pathogenic *Escherichia coli* strains of avian and human origin: link between phylogenetic relationships and common virulence patterns. *J. Clin. Microbiol.* 45, 3366–3376. doi: 10.1128/JCM.00037-07
- Nandanwar, N., Janssen, T., Kuhl, M., Ahmed, N., Ewers, C., and Wieler, L. H. (2014). Extraintestinal pathogenic *Escherichia coli* (ExPEC) of human and avian origin belonging to sequence type complex 95 (STC95) portray indistinguishable virulence features. *Int. J. Med. Microbiol.* 304, 835–842. doi: 10.1016/j.ijmm.2014.06.009
- Olesen, B., Kolmos, H. J., Orskov, F., and Orskov, I. (1995). A comparative study of nosocomial and community-acquired strains of *Escherichia coli* causing bacteraemia in a Danish University Hospital. *J. Hosp. Infect.* 31, 295–304. doi: 10.1016/0195-6701(95)90208-2
- Orskov, F. (1978). Virulence factors of the bacterial cell surface. *J. Infect. Dis.* 137, 630–633. doi: 10.1093/infdis/137.5.630
- Orskov, F., and Orskov, I. (1984). “Serotyping of *Escherichia coli*,” in *Methods in Microbiology*, ed T. Bergan (London: Academic Press), 43–113.
- Orskov, F., and Orskov, I. (1992). *Escherichia coli* serotyping and disease in man and animals. *Can. J. Microbiol.* 38, 699–704. doi: 10.1139/m92-115
- Orskov, I., Orskov, F., Jann, B., and Jann, K. (1977). Serology, chemistry, and genetics of O and K antigens of *Escherichia coli*. *Bacteriol. Rev.* 41, 667–710.
- Orskov, I., Orskov, F., and Rowe, B. (1984). Six new *E. coli* O groups: O165, O166, O167, O168, O169 and O170. *Acta Pathol. Microbiol. Immunol. Scand. B* 92, 189–193.
- Peigne, C., Bidet, P., Mahjoub-Messai, F., Plainvert, C., Barbe, V., Médigue, C., et al. (2009). The plasmid of *Escherichia coli* strain S88 (O45:K1:H7) that causes neonatal meningitis is closely related to avian pathogenic *E. coli* plasmids and is associated with high-level bacteremia in a neonatal rat meningitis model. *Infect. Immun.* 77, 2272–2284. doi: 10.1128/IAI.01333-08
- Perelle, S., Dilasser, F., Grout, J., and Fach, P. (2005). Detection of *Escherichia coli* serogroup O103 by real-time polymerase chain reaction. *J. Appl. Microbiol.* 98, 1162–1168. doi: 10.1111/j.1365-2672.2005.02545.x
- Perry, L. L., Sanmiguel, P., Minocha, U., Terekhov, A. I., Shroyer, M. L., Farris, L. A., et al. (2009). Sequence analysis of *Escherichia coli* O157:H7 bacteriophage PhiV10 and identification of a phage-encoded immunity protein that modifies the O157 antigen. *FEMS Microbiol. Lett.* 292, 182–186. doi: 10.1111/j.1574-6968.2009.01511.x
- Reeves, P. R., Hobbs, M., Valvano, M. A., Skurnik, M., Whitfield, C., Coplin, D., et al. (1996). Bacterial polysaccharide synthesis and gene nomenclature. *Trends Microbiol.* 4, 495–503. doi: 10.1016/S0966-842X(97)82912-5
- Riley, L. W. (2014). Pandemic lineages of extraintestinal pathogenic *Escherichia coli*. *Clin. Microbiol. Infect.* 20, 380–390. doi: 10.1111/1469-0691.12646
- Sánchez, S., García-Sánchez, A., Martínez, R., Blanco, J., Blanco, J. E., Blanco, M., et al. (2009). Detection and characterisation of Shiga toxin-producing *Escherichia coli* other than *Escherichia coli* O157:H7 in wild ruminants. *Vet. J.* 180, 384–388. doi: 10.1016/j.tvjl.2008.01.011
- Scheutz, F., Cheasty, T., Woodward, D., and Smith, H. R. (2004). Designation of O174 and O175 to temporary O groups OX3 and OX7, and six new *E. coli* O groups that include Verocytotoxin-producing *E. coli* (VTEC): O176, O177, O178, O179, O180 and O181. *APMIS* 112, 569–584. doi: 10.1111/j.1600-0463.2004.apm1120903.x
- Scheutz, F., Teel, L. D., Beutin, L., Piérard, D., Buvens, G., Karch, H., et al. (2012). Multicenter evaluation of a sequence-based protocol for subtyping Shiga toxins and standardizing Stx nomenclature. *J. Clin. Microbiol.* 50, 2951–2963. doi: 10.1128/JCM.00860-12
- Scott, L., Mcgee, P., Walsh, C., Fanning, S., Sweeney, T., Blanco, J., et al. (2009). Detection of numerous verotoxigenic *E. coli* serotypes, with multiple antibiotic resistance from cattle faeces and soil. *Vet. Microbiol.* 134, 288–293. doi: 10.1016/j.vetmic.2008.08.008
- Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30, 2068–2069. doi: 10.1093/bioinformatics/btu153

- Singer, R. S. (2015). Urinary tract infections attributed to diverse ExPEC strains in food animals: evidence and data gaps. *Front. Microbiol.* 6:28. doi: 10.3389/fmicb.2015.00028
- Stenutz, R., Weintraub, A., and Widmalm, G. (2006). The structures of *Escherichia coli* O-polysaccharide antigens. *FEMS Microbiol. Rev.* 30, 382–403. doi: 10.1111/j.1574-6976.2006.00016.x
- Sugiyama, T., Kido, N., Kato, Y., Koide, N., Yoshida, T., and Yokochi, T. (1998). Generation of *Escherichia coli* O9a serotype, a subtype of *E. coli* O9, by transfer of the wb* gene cluster of *Klebsiella* O3 into *E. coli* via recombination. *J. Bacteriol.* 180, 2775–2778.
- Vincent, C., Boerlin, P., Daignault, D., Dozois, C. M., Dutil, L., Galanakis, C., et al. (2010). Food reservoir for *Escherichia coli* causing urinary tract infections. *Emerg. Infect. Dis.* 16, 88–95. doi: 10.3201/eid1601.091118
- Wang, L., Huskic, S., Cisterne, A., Rothmund, D., and Reeves, P. R. (2002). The O-antigen gene cluster of *Escherichia coli* O55:H7 and identification of a new UDP-GlcNAc C4 epimerase gene. *J. Bacteriol.* 184, 2620–2625. doi: 10.1128/JB.184.10.2620-2625.2002
- Wullenweber, M., Beutin, L., Zimmermann, S., and Jonas, C. (1993). Influence of some bacterial and host factors on colonization and invasiveness of *Escherichia coli* K1 in neonatal rats. *Infect. Immun.* 61, 2138–2144.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Delannoy, Beutin, Mariani-Kurkdjian, Fleiss, Bonacorsi and Fach. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.