



OPEN ACCESS

EDITED AND REVIEWED BY

Enrico Cherubini,
European Brain Research Institute, Italy

*CORRESPONDENCE

Zila Martinez-Lozada
martinezlz@chop.edu

SPECIALTY SECTION

This article was submitted to
Cellular Neurophysiology,
a section of the journal
Frontiers in Cellular Neuroscience

RECEIVED 28 July 2022

ACCEPTED 04 August 2022

PUBLISHED 17 August 2022

CITATION

Martinez-Lozada Z, Hewett SJ, Zafra F
and Ortega A (2022) Editorial: The
known, the unknown, and the future
of glutamate transporters.
Front. Cell. Neurosci. 16:1005834.
doi: 10.3389/fncel.2022.1005834

COPYRIGHT

© 2022 Martinez-Lozada, Hewett,
Zafra and Ortega. This is an
open-access article distributed under
the terms of the [Creative Commons
Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which
does not comply with these terms.

Editorial: The known, the unknown, and the future of glutamate transporters

Zila Martinez-Lozada^{1*}, Sandra J. Hewett², Francisco Zafra^{3,4}
and Arturo Ortega⁵

¹Department of Pediatrics, The Children's Hospital of Philadelphia, Philadelphia, PA, United States, ²Program in Neuroscience, Department of Biology, Syracuse University, Syracuse, NY, United States, ³Center of Molecular Biology Severo Ochoa, School of Science, Consejo Superior de Investigaciones Científicas, Universidad Autónoma de Madrid, Madrid, Spain, ⁴Instituto de Investigación Sanitaria del Hospital Universitario La Paz — IdiPAZ, Madrid, Spain, ⁵Department of Toxicology, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Ciudad de México, Mexico

KEYWORDS

glutamate, glutamate transport, excitatory amino acid transporter (EAAT), vesicular glutamate transport proteins, system x_c^-

Editorial on the Research Topic

The known, the unknown, and the future of glutamate transporters

Glutamate is the main excitatory neurotransmitter in the central nervous system (Fonnum, 1984). It is required for essentially all cognitive functions, however, is also a neurotoxin. Therefore, maintenance of the glutamate extracellular concentration involves tight control of its release and uptake. Several glutamate transport proteins contribute to this regulation. Vesicular glutamate transporters (VGLUTs) package glutamate into synaptic vesicles (Omote et al., 2011). The excitatory amino acid transporters (EAATs), under physiological conditions, remove glutamate from the synaptic cleft (Danbolt et al., 2016). Lastly, the cystine/glutamate exchanger, also known as system x_c^- , exports glutamate in exchange for cystine (Jabaudon et al., 1999; Warr et al., 1999; Featherstone and Shippy, 2008). In this Research Topic, we assemble a review of current literature and new research on these transporters.

VGLUTs

Glutamate is packaged into synaptic vesicles *via* one of three VGLUTs. As part of our collection, Hori and Takamori describe a novel method to monitor glutamate transport *in living nerve terminals* using the rodent giant synapse, the calyx of Held. In addition, they discuss what is presently known about factors that alter the amount and rate of glutamate refilling of synaptic vesicles and the relevance of these findings to central nervous system disorders (Hori and Takamori).

EAATs

Termination of glutamate signaling is mediated *via* uptake by one of five EAATs (Hediger et al., 2013). Not surprisingly, impaired expression and/or activity of EAATs have negative repercussions on health. EAAT2 (GLT-1a,b) is predominately expressed by astrocytes, although a small portion of GLT-1a can be found on axon terminals (Chen et al., 2004; Furness et al., 2008; Melone et al., 2009; Zhou et al., 2018).

In this article collection, Yeung et al. examined the expression of EAAT2 in several brain regions from *postmortem* tissue of patients with Alzheimer's disease (AD) or control (Yeung et al.). The authors found no significant change in EAAT2 density but did observe spatial differences in EAAT2 expression in AD tissue with less immunoreactivity detected in main astrocyte branches, especially on those surrounding neuronal cell bodies (Yeung et al.). Whether this altered expression pattern has implications for glutamate recycling in AD remains to be determined. Direct evidence for pathological loss of function of EAAT2 in a mouse model of Huntington's disease (HD) comes from the paper of Hirschberg et al. Therein they report that abnormal protein-protein interactions of mutant huntingtin (mHTT) with EAAT2 binding partners reduce glutamate uptake in striatal astrocytes and mediate some of the HD-associated deficits studied. Abnormalities in uptake and motor function were alleviated *via* over-expression of a C-terminal truncated EAAT2 protein (Hirschberg et al.).

Because most of the glutamate clearance (80–90%) is mediated by astrocytic EAAT2/GLT-1, the function of GLT-1 in other cell types has remained largely unexplored. In this collection, two groups report interesting findings concerning the physiological function of GLT-1 in neurons and oligodendrocytes using cell-specific knockout mouse lines. The Rosenberg group found that hippocampal slices prepared from neuronal GLT-1 KO (synGLT-1 KO) mice are more vulnerable to excitotoxicity than slices from wild-type mice (Rimmele et al.). Whether this is due to metabolic compromise, or a disturbance of glutamate homeostasis is still unknown. Meanwhile, the Fuss group reports that deletion of GLT-1 (*GLT1^{Plp1}icKO* mice) in maturing oligodendrocytes leads to hypomyelination in the *corpus callosum* of male, but not female, mice (Thomason et al.). The exact mechanism responsible for these abnormalities is at present unknown.

Finally, the Robinson group used pharmacological inhibition of EAAT function to demonstrate their importance to arteriole patency (Jackson et al.), highlighting another important function of EAATs, which is to act as a bridge between local neuronal activity and increases in blood flow. One can speculate that the dysfunction of EAATs observed in several neuropathologies could negatively affect neurovascular coupling.

It is worth nothing that EAATs not only transport glutamate but act as anion channels (Wadiche et al., 1995; Bergles et al., 2002; Jen et al., 2005). Herein, Kovermann

et al. review what is known about this function of the EAATs in both health and disease (Kovermann et al.). For example, they discuss the relationship between EAAT mutations that alter anion channel activity (but not glutamate uptake) resulting in chloride dyshomeostasis and symptoms of neurological disease.

System x_c^-

System x_c^- (Sx_c^-) is a Na^+ -independent, Cl^- -dependent heteromeric amino acid transporter — formed by two polypeptides, xCT, the subunit responsible for the transport function, and 4F2hc, the subunit required for membrane localization — that functions physiologically to export glutamate while importing cystine in a 1:1 ratio (Bannai and Kitamura, 1980; Bannai, 1986). Astrocytes appear to be the main cell type expressing Sx_c^- in the mature brain (Zhang et al., 2014; Ottestad-Hansen et al., 2018). Sx_c^- activity contributes to the maintenance of redox homeostasis (Banjac et al., 2008), is important for the synthesis of glutathione (Sato et al., 1998), and is a major source of ambient extracellular glutamate *in vivo* (Baker et al., 2002; De Bundel et al., 2011). However, under pathological conditions, glutamate release through Sx_c^- contributes to neurological diseases/disorders [for review see Lewerenz et al. (2013)].

In this article collection, the Hermans group describes and validates a method using tritiated glutamate as a substrate for reversed transport to evaluate the activity of system x_c^- (Beckers et al.) both in cultured cells and in synaptosomal preparations. Meanwhile, Bentea et al. investigated the effect of genetic deletion of xCT on two models of Parkinson's disease (PD) (Bentea et al.). They found protection against proteasome inhibition-induced nigrostriatal degeneration, — but not MPTP-induced striatal toxicity — in $xCT^{-/-}$ mice. Last but not least, He and Hewett evaluated the contribution of Sx_c^- to ischemic stroke (He and Hewett). Their results demonstrate that Sx_c^- contributes to cortical ischemic damage when blood flow is moderately but not severely reduced.

Altogether, in this Research Topic, the authors review current literature or provide original research centered on understanding more fully the physiological and/or pathophysiological function of glutamate transport. The work presented here highlights the need for further research on the myriad of roles glutamate transporters play in health and pathology. Therefore, we hope that this collection encourages additional research in this field.

Author contributions

ZM-L produced the initial draft. SH, FZ, and AO further edited and provided additional content and references. All authors approved the final version for publication.

Acknowledgments

We want to thank all the authors of this Research Topic for their contributions, and the reviewers for their insightful comments that helped maintain the articles at the highest standards.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships

References

- Baker, D. A., Xi, Z. X., Shen, H., Swanson, C. J., and Kalivas, P. W. (2002). The origin and neuronal function of in vivo nonsynaptic glutamate. *J. Neurosci.* 22, 9134–9141. doi: 10.1523/JNEUROSCI.22-20-09134.2002
- Banjac, A., Perisic, T., Sato, H., Seiler, A., Bannai, S., Weiss, N., et al. (2008). The cystine/cysteine cycle: a redox cycle regulating susceptibility versus resistance to cell death. *Oncogene* 27, 1618–1628. doi: 10.1038/sj.onc.1210796
- Bannai, S. (1986). Exchange of cystine and glutamate across plasma membrane of human fibroblasts. *J. Biol. Chem.* 261, 2256–2263. doi: 10.1016/S0021-9258(17)35926-4
- Bannai, S., and Kitamura, E. (1980). Transport interaction of L-cystine and L-glutamate in human diploid fibroblasts in culture. *J. Biol. Chem.* 255, 2372–2376. doi: 10.1016/S0021-9258(19)85901-X
- Bergles, D. E., Tzingounis, A. V., and Jahr, C. E. (2002). Comparison of coupled and uncoupled currents during glutamate uptake by GLT-1 transporters. *J. Neurosci.* 22, 10153–10162. doi: 10.1523/JNEUROSCI.22-23-10153.2002
- Chen, W., Mahadomrongkul, V., Berger, U. V., Bassan, M., DeSilva, T., Tanaka, K., et al. (2004). The glutamate transporter GLT1a is expressed in excitatory axon terminals of mature hippocampal neurons. *J. Neurosci.* 24, 1136–1148. doi: 10.1523/JNEUROSCI.1586-03.2004
- Danbolt, N., Furness, D., and Zhou, Y. (2016). Neuronal vs glial glutamate uptake: resolving the conundrum. *Neurochem. Int.* 98, 29–45. doi: 10.1016/j.neuint.2016.05.009
- De Bundel, D., Schallier, A., Loyens, E., Fernando, R., Miyashita, H., Van Liefferinge, J., et al. (2011). Loss of system xc⁻ does not induce oxidative stress but decreases extracellular glutamate in hippocampus and influences spatial working memory and limbic seizure susceptibility. *J. Neurosci.* 31, 5792–5803. doi: 10.1523/JNEUROSCI.5465-10.2011
- Featherstone, D. E., and Shippy, S. A. (2008). Regulation of synaptic transmission by ambient extracellular glutamate. *Neuroscientist* 14, 171–181. doi: 10.1177/1073858407308518
- Fonnum, F. (1984). Glutamate: a neurotransmitter in mammalian brain. *J. Neurochem.* 42, 1–11. doi: 10.1111/j.1471-4159.1984.tb09689.x
- Furness, D., Dehnes, Y., Akhtar, A., Rossi, D., Hamann, M., Grutle, N., et al. (2008). A quantitative assessment of glutamate uptake into hippocampal synaptic terminals and astrocytes: new insights into a neuronal role for excitatory amino acid transporter 2 (EAAT2). *Neuroscience* 157, 80–94. doi: 10.1016/j.neuroscience.2008.08.043
- Hediger, M. A., Cléménçon, B., Burrier, R. E., and Bruford, E. A. (2013). The ABCs of membrane transporters in health and disease (SLC series): introduction. *Molec. Aspects Med.* 34, 95–107. doi: 10.1016/j.mam.2012.12.009
- Jabaudon, D., Shimamoto, K., Yasuda-Kamatani, Y., Scanziani, M., Gahwiler, B. H., and Gerber, U. (1999). Inhibition of uptake unmasks rapid extracellular turnover of glutamate of nonvesicular origin. *Proc. Natl. Acad. Sci. U S A.* 96, 8733–8738. doi: 10.1073/pnas.96.15.8733
- Jen, J. C., Wan, J., Palos, T. P., Howard, B. D., and Baloh, R. W. (2005). Mutation in the glutamate transporter EAAT1 causes episodic ataxia, hemiplegia, and seizures. *Neurology* 65, 529–534. doi: 10.1212/01.WNL.0000172638.58172.5a
- Lewerenz, J., Hewett, S. J., Huang, Y., Lambros, M., Gout, P. W., Kalivas, P. W., et al. (2013). The cystine/glutamate antiporter system x(c)⁻ in health and disease: from molecular mechanisms to novel therapeutic opportunities. *Antioxid Redox Signal* 18, 522–555. doi: 10.1089/ars.2011.4391
- Melone, M., Belleli, M., and Conti, F. (2009). Synaptic localization of GLT-1a in the rat somatic sensory cortex. *Glia* 57, 108–117. doi: 10.1002/glia.20744
- Omote, H., Miyaji, T., Juge, N., and Moriyama, Y. (2011). Vesicular neurotransmitter transporter: bioenergetics and regulation of glutamate transport. *Biochemistry* 50, 5558–5565. doi: 10.1021/bi200567k
- Ottestad-Hansen, S., Hu, Q. X., Follin-Arbelet, V. V., Bentea, E., Sato, H., Massie, A., et al. (2018). The cystine-glutamate exchanger (xCT, Slc7a11) is expressed in significant concentrations in a subpopulation of astrocytes in the mouse brain. *Glia* 66, 951–970. doi: 10.1002/glia.23294
- Sato, Kuriyama-Matsumura, K., Siow, R., Ishii, T., Bannai, S., and Mann, G. E. (1998). Induction of cystine transport via system xc⁻ and maintenance of intracellular glutathione levels in pancreatic acinar and islet cell lines. *Biochim. Biophys. Acta (BBA)-Biomembranes* 1414, 85–94. doi: 10.1016/S0005-2736(98)00159-X
- Wadiche, J. I., Arriza, J. L., Amara, S. G., and Kavanaugh, M. P. (1995). Kinetics of a human glutamate transporter. *Neuron* 14, 1019–1027. doi: 10.1016/0896-6273(95)90340-2
- Warr, O., Takahashi, M., and Attwell, D. (1999). Modulation of extracellular glutamate concentration in rat brain slices by cystine-glutamate exchange. *J. Physiol.* 514, 783–793. doi: 10.1111/j.1469-7793.1999.783ad.x
- Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O’Keefe, S., et al. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J. Neurosci.* 34, 11929–11947. doi: 10.1523/JNEUROSCI.1860-14.2014
- Zhou, Y., Hassel, B., Eid, T., and Danbolt, N. C. (2018). Axon-terminals expressing EAAT2 (GLT-1; Slc1a2) are common in the forebrain and not limited to the hippocampus. *Neurochem. Int.* 123, 101–113. doi: 10.1016/j.neuint.2018.03.006

that could be construed as a potential conflict of interest.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.