



## OPEN ACCESS

EDITED BY  
Yao Sun,  
Tongji University, China

REVIEWED BY  
Vehid Salih,  
UCL Eastman Dental Institute,  
United Kingdom

\*CORRESPONDENCE  
Francisco Wanderley Garcia  
Paula-Silva  
franciscogarcia@forp.usp.br

SPECIALTY SECTION  
This article was submitted to  
Regenerative Dentistry,  
a section of the journal  
Frontiers in Dental Medicine

RECEIVED 12 May 2022  
ACCEPTED 15 July 2022  
PUBLISHED 02 August 2022

CITATION  
Lorencetti-Silva F, Sales LS, Lamarque  
GCC, Caixeta GA, Arnez MFM,  
Faccioli LH and Paula-Silva FWG (2022)  
Effects of inflammation in dental pulp  
cell differentiation and reparative  
response. *Front. Dent. Med.* 3:942714.  
doi: 10.3389/fdmed.2022.942714

COPYRIGHT  
© 2022 Lorencetti-Silva, Sales,  
Lamarque, Caixeta, Arnez, Faccioli and  
Paula-Silva. This is an open-access  
article distributed under the terms of  
the [Creative Commons Attribution  
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution  
or reproduction in other forums is  
permitted, provided the original  
author(s) and the copyright owner(s)  
are credited and that the original  
publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or  
reproduction is permitted which does  
not comply with these terms.

# Effects of inflammation in dental pulp cell differentiation and reparative response

Francine Lorencetti-Silva<sup>1</sup>, Larissa Sthefani Sales<sup>1</sup>,  
Giuliana de Campos Chaves Lamarque<sup>1</sup>,  
Giovana Amorim Caixeta<sup>1</sup>, Maya Fernanda Manfrin Arnez<sup>1</sup>,  
Lúcia Helena Faccioli<sup>2</sup> and  
Francisco Wanderley Garcia Paula-Silva<sup>1\*</sup>

<sup>1</sup>Department of Pediatric Clinics, School of Dentistry of Ribeirão Preto, University of São Paulo, São Paulo, Brazil, <sup>2</sup>Departamento de Análises Clínicas, Toxicológicas e Bromatológicas, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, São Paulo, Brazil

The responsiveness of the dentin-pulp complex is possible due to the stimulation of dental pulp cells, which begin to synthesize and secrete dentin matrix. The inflammatory process generated by harmful stimuli should be understood as a natural event of the immune response, resulting in the recruitment of hematopoietic cells, which cross the endothelial barrier and reach the site affected by the injury in order to eliminate the damage and provide an appropriate environment for the restoration of homeostasis. The repair process occurs in the presence of adequate blood supply, absence of infection, and with the participation of pro-inflammatory cytokines, growth factors, extracellular matrix components, and other biologically active molecules. Prostaglandins and leukotrienes are bioactive molecules derived from the metabolism of arachidonic acid, as a result of a variable range of cellular stimuli. The aim of this review is to describe the process of formation and biomineralization of the dentin-pulp complex and how pro-inflammatory events can modify this response, with emphasis on the lipid mediators prostaglandins and leukotrienes derived from arachidonic acid metabolism.

## KEYWORDS

dental pulp, biomineralization, inflammatory mediators, prostaglandins, leukotrienes

## Introduction

The responsiveness of the dentin-pulp complex to injury is possible by the stimulation of odontoblasts, which begin to synthesize and secrete reactionary dentin matrix, Dentin matrix secretion by primary odontoblasts that survived the tissue injury occurs via signaling molecules similar to those involved in the dentinogenesis phase (1–3). Odontoblasts therefore represent the first line of defense against bacterial invasion in this environment (4, 5).

The inflammatory process is important to fight infection and is essential to protect the tissue from injury and restore its physiological function. Cytokines and various signaling molecules are synthesized and secreted by the host cells of the dentin-pulpal complex prior to the recruitment and activation of immune system cells, which reveals that the dentin-pulpal complex generates a molecular immune response pattern prior to the cellular immune response (6).

Repair, therefore, occurs in the face of adequate blood supply and with the participation of pro-inflammatory cytokines, growth factors, components of the extracellular matrix and other biologically active molecules (7, 8). The aim of this review is to describe the process of formation and biomineralization of the dentino-pulpal complex and how pro-inflammatory events can modify this response, with emphasis on the lipid mediators prostaglandins and leukotrienes derived from arachidonic acid metabolism.

## Formation of the dentin-pulp complex

Deciduous teeth begin their formation between the 3rd and 8th weeks of intrauterine life, whose development involves a series of complex steps regulated by interactions between epithelial and mesenchymal tissue derived from the ectoderm (ameloblasts) and the neural crest (odontoblasts) (9, 10).

For tooth development, ectomesenchymal cells derived from the first branchial arch and neural crest migrate and form cell aggregations (11). The ectoderm thickens and generates sprouts that invade the neural crest-derived mesenchyme. The adjacent epithelium then starts sending signals to the mesenchyme that undergoes condensation around the epithelial band, which undergoes proliferation and surrounds the mesenchyme of the dental papilla (9). The epithelial cells that undergo the differentiation process start secreting enamel matrix and are called ameloblasts, while the differentiated mesenchymal cells, now called odontoblasts, secrete dentin (11).

After complete differentiation, odontoblasts are characterized by a tall columnar shape, polarized nuclear and cytoplasmic organelles, and are united to each other by junctional complexes (12). These are therefore post-mitotic cells, organized in the form of a peripheral cell layer, present along the dentin-pulp interface that have cellular processes extending within tubular structures surrounded by dentin, called dentinal tubules (8, 12, 13). After the cell differentiation process, odontoblasts synthesize organic matrix consisting of type I collagen, and play an important role in the mineralization of this matrix by secreting proteoglycans and non-collagen proteins that participate in nucleation and control of mineral phase growth (8, 14).

During dentin formation, odontoblasts secrete a matrix rich in type I collagen, which is called pre-dentin, thus constituting the organic phase, which in turn is mineralized through the incorporation of hydroxyapatite (HA) crystals through the biomineralization process, which involves mechanisms that control both the sites and the deposition rate of these crystals (14, 15).

## Molecules involved in dentin biomineralization

Alkaline phosphatase (ALPL) is an indispensable enzyme for mineralization of the secreted matrix, as it provides phosphate ions that generate the precipitation of apatite minerals, and hydrolyzes inorganic pyrophosphate, a phosphate ester that inhibits mineralization (16). Alkaline phosphatases is found in four isoforms (isozymes), including Tissue Non-Specific Alkaline Phosphatase (TNAP) (17). The mineralization of the dentin matrix itself starts in plasma membrane-derived matrix vesicles composed of proteins and lipids, which allows the accumulation of high concentrations of calcium ions ( $\text{Ca}^{2+}$ ) and phosphate ( $\text{PO}_4^{3-}$ ), and the participation of Non-Specific Tissue Alkaline Phosphatase (TNAP). This enzyme encoded by the ALPL gene leads to modification of the extracellular matrix and expression of Phosphatase Orfan 1 (PHOSPHO 1), this being the initiating factor for HA deposition inside matrix vesicles (14). Collagen fibers also participate in the enucleation and growth of apatite crystals (18), as do non-collagen proteins characterized by acidity due to their high doses of aspartic and glutamic acids, and phosphorylated serine residues (19).

Among the proteins involved during the mineralization phase are Dentin Matrix Protein 1 (DMP-1) and Dentin Sialophosphoprotein (DSPP) (12, 20). DMP-1 is expressed in both pulp and odontoblastic cells and plays an important regulatory role in odontoblast differentiation, as well as participates in tubular dentin formation and mineralization, proving essential in both the earliest and most advanced stages of odontogenesis (15). DMP-1-mediated mineral deposition begins when this protein binds to calcium ions, because the peptide arrangement of DMP-1 presents domains corresponding to the structure of HA crystals, which in turn reduces activation energy and favors the formation of structurally functional crystalline nuclei (21).

The DSPP, once secreted is rapidly cleaved into COOH-terminal and NH<sub>2</sub>-terminal fragments. The latter is encoded by the 5' portion of the Dspp gene, resulting Dentin Sialoprotein (DSP) and proteoglycans, while the 3' portion is responsible for encoding the NH<sub>2</sub>-terminal portion, which generates Dentin Phosphoprotein (DPP), these two being the most commonly found non-collagen proteins in the dentin matrix (22–24).

DSPP is a member of a family of proteins called SIBLINGs, an acronym for Small Integrin-Binding Ligand N-linked

Glycoproteins, which also includes Bone Sialoprotein (BSP or IBSP), Dentin Matrix Protein-1 (DMP-1), Osteopontin (OPN), and Extracellular Matrix Phosphoglycoprotein (MEPE) (25, 26). These proteins play an important role in dentinogenesis and can be considered markers of differentiated odontoblasts (27).

DPP contains a large amount of aspartic acid and phosphoserines, which allows it to be characterized as a polyanionic molecule. The negative charge distributed along this protein increases its affinity for calcium ions and exposes them, in this way, to the collagen fibers present in front of the mineralized layer, allows the growth of HA crystals, which gives it importance during the maturation phase of mineralized dentin (26, 28).

The increase in *Dspp* gene expression occurs through signaling pathways whose participation involves Bone Morphogenetic Protein-2 (BMP-2) and the Runt-related Transcription Factor (Runx2) (29).

Bone Morphogenetic Proteins (BMPs) are signaling molecules that are part of the Transforming Growth Factor  $\beta$  (TGF- $\beta$ ) superfamily (30). In addition to increasing *Dspp* gene expression, BMP-2 also plays a very important role in regulating the differentiation process of dental pulp cells into odontoblasts, and they are capable of both producing and cleaving this protein (31).

Runx2, in turn, is a transcription factor expressed in the papilla and dental sac, and is involved in the differentiation of odontoblasts and osteoblasts. This transcription factor has the ability to increase DSPP expression in immature odontoblasts, unlike in cases of fully differentiated cells, which reveals that the effect of Runx2 is influenced by the differentiation state of the odontoblast cell (32). During tooth development, Runx2 is expressed in the dental mesenchyme until the cap stage and then has its expression ceased in the dental papilla during the odontoblast differentiation stage, suggesting that this gene presents importance in tooth morphogenesis (32).

The process of terminal differentiation of odontogenic cells results from molecular interactions that occur between dental epithelium and ectomesenchymal cells, involving BMPs, fibroblast growth factors (FGF) and transcriptional factors such as *Msx*, which support the epithelium-mesenchymal interactions crucial for the onset of tooth development (33). The *Msx* gene is a member of the homeobox gene family expressed during the early stages of craniofacial formation, including the condensation of the ectomesenchymal tissue of the tooth germ (34). The *MSX1* transcriptional factor induces mesenchymal cell proliferation and prevents odontoblast differentiation at the hood stage by inhibiting the expression of BMPs, including BMP-2 (35).

A non-collagenous protein present in the dentinal matrix secreted by odontoblasts is integrin-bound sialoprotein (IBSP) (36). It is an acidic glycoprotein expressed by osteoblasts, odontoblasts, and cementoblasts during the early stages of mineralization, and has the ability to bind to HA via sequential

polyglutamine acidic bonds (37). IBSP is present in matrix vesicles and constitutes one of the initiating proteins of HA crystal deposition. This process is possible thanks to the synergistic action with the ALPL enzyme, because in the presence of IBSP, high levels of this enzyme are able to induce the initiation of mineral deposition (29).

Osteocalcin (BGLAP) is revealed as the main non-collagenous protein produced by odontoblasts and osteoclasts, whose role is to regulate the organization of the extracellular matrix through interaction with HA, protein matrix and surface receptors (38).

In summary, odontoblasts secrete specific dentin matrix proteins, such as DSP and DPP derived from DSPP. Dentin also presents in its composition collagen proteins, IBSP, DMP-1, BGLAP, and ALPL that share common regulatory pathways through transcription factors RUNX-2 and MSX-1 (39, 40).

## Immune system in the dentin-pulp complex

In addition to the important role of odontoblasts in the process of dentin matrix synthesis, as described earlier, these cells are also important sensory structures of the pulp organ, as they have the ability to detect bacterial invasion during the development of dental caries and sequentially initiate the immune response in the pulp (8).

The cells of the innate immune system possess receptors that recognize pathogen-associated molecular patterns (PAMPs), which includes bacterial components such as lipopolysaccharide (LPS) and lipoteichoic acid (LTA) (41, 42). Through pattern recognition receptors (PRRs), odontoblasts are able to respond to the invasion of pathogens Toll-like receptors (TLRs), specifically TLR2 and TLR4, and Leucine Rich Nucleotides (NLRs), the most prominent being the NOD2 receptors, capable of recognizing peptidoglycans present in Gram-positive and Gram-negative bacteria and thus activate the MAPK and NF- $\kappa$ B signaling pathways, in order to produce pro-inflammatory cytokines (43). The interaction of TLR-2 with LTA, a structure present in the cell wall of Gram-positive bacteria, promotes nuclear translocation of the transcription factor NF- $\kappa$ B, and generates the production of chemokines, including CCL2, CXCL1, CXCL2, CXCL8, and CXCL10, recruits immature dendritic cells (4, 44, 45) and reduces the expression of type 1 collagen and DSPP, components of the dentinal matrix (44). The TLR4 receptor recognizes LPS present in Gram-negative bacteria and consequently increases the expression of important pro-inflammatory cytokines such as IL-1 $\beta$ , TNF- $\alpha$ , and IL-8 (46). Furthermore, TLR4 receptors may play a crucial role in the immune response by activating and regulating pulp stem cell proliferation and migration processes (47).

The inflammatory process generated under harmful stimuli should be understood as a natural event of the immune response, resulting in the recruitment of hematopoietic cells, which cross the endothelial barrier and reach the site affected by the injury in order to eliminate the damage and provide a suitable environment for the restoration of homeostasis (48). This requires the activation of PRRs, the release of mediators, such as Leukotriene B<sub>4</sub> (LTB<sub>4</sub>) and Prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), being LTB<sub>4</sub> the main mediator in the recruitment of polymorphonuclear cells (49–52).

## Inflammatory lipid mediators - prostaglandins and leukotrienes

PGs and LTs are bioactive molecules derived from the metabolism of arachidonic acid, a polyunsaturated fatty acid derived from cell membrane phospholipids, by action of the enzyme Phospholipase A<sub>2</sub> (PLA<sub>2</sub>). PLA<sub>2</sub> are a group of proteins that have the ability to hydrolyze the fatty acid at the sn-2 position of glycerophospholipids, especially the PLA<sub>2</sub> IV group (cPLA<sub>2</sub>), as a result of a variable range of cellular stimuli (53–55). Because these are molecules generated from the oxidation of carbon 20 of the polyunsaturated fatty acid they are called eicosanoids (from the Greek eikosi = twenty) (54, 56, 57).

Free arachidonic acid can be metabolized via the Cyclooxygenase-1 (COX-1) and Cyclooxygenase-2 (COX-2) pathways and generate Prostaglandins (PGs) or Thromboxanes (TX), or it can be oxidized along the Lipoxygenase (LO) pathway, which includes the enzyme 5-Lipoxygenase (5-LO), to produce different classes of LTs and Lipoxins (54).

One of the pathways of AA metabolism is the 5-LO pathway. In the presence of FLAP, a membrane-associated nuclear protein, the 5-LO enzyme is activated and oxidizes AA, converting it to 5S-hydroperoxyeicosatetraenoic acid (5S-HpETE), which is further reduced by the enzyme peroxidase to 5S-hydroxyeicosatetraenoic acid (5S-HETE) or is converted to LTA<sub>4</sub>, which by action of LTA<sub>4</sub> hydrolase, results in LTB<sub>4</sub> (58). These mediators are involved in chemotaxis of neutrophils, dendritic cells and T cells, increase vascular permeability and act directly on antigen presenting cells (59). Chemotaxis is correlated to the activation of the BLT1 receptor in addition to the BLT2 receptor, with the latter showing low affinity for LTB<sub>4</sub> (60).

PGs, in turn, are produced by a sequence of events involving the actions of the COX-1 and COX-2 enzymes and have their structural basis consisting of prostanoid acid, which is composed of a cyclopentane ring and two carbon chains, and are known as prostanoids (61).

Importantly, the enzyme COX-2 has its expression induced by a range of stimuli related to the inflammatory response, such as growth factors and cytokines, and is therefore considered

the inducible isoform of COX. This enzyme is responsible for the synthesis of PGs involved in the inflammatory response, although they are expressed in organs such as the brain and kidneys under physiological conditions. PGs derived from COX-1 are involved in the maintenance of biological functions (62). The enzyme COX metabolizes AA and converts it into the intermediate isoform of PGE<sub>2</sub>: PGH<sub>2</sub>, which is converted to PGE<sub>2</sub> by means of the microsomal Prostaglandin E synthases 1 and 2 (mPGE-1 and mPGE-2), and this mediator acts on 4 different types of membrane receptors (EP1, EP2, EP3, and EP4) coupled to G proteins (G<sub>αs</sub>, G<sub>i</sub> and G<sub>q</sub>) and, depending on the type of receptor stimulated, different cellular pathways are activated (63).

Thus, the induction of the inflammatory process generates PGs release (64), which causes increased local blood flow, increased vascular permeability (when associated with other soluble factors, such as leukotrienes), and sensitization of afferent nerve fibers, generating hyperalgesia by acting on peripheral sensory neurons, in sites of the spinal cord and brain. These mediators contribute to the amplification of the pattern of inflammatory response, in order to promote both the increase and prolongation of the effects and signals produced by pro-inflammatory agents (62).

It is known that experimentally, the presence of LPS generates in the pulp an increase in AA metabolism, resulting in increased gene and protein expression of COX-2 and increased production of PGE<sub>2</sub> in the inflammatory environment, which is involved in the increased vascular permeability of the pulp (65–67). LTB<sub>4</sub> production and its receptors expression (BLT1 and BLT2) was shown to be almost parallel to neutrophil infiltration, which reveals involvement of this mediator in the infiltration of these cells in experimental pulp inflammation (66, 68).

In intense inflammatory pulp conditions, the deposition of reactive dentin by odontoblasts can be stopped (8, 69), and dentin repair can be performed from stem cells present in the dental pulp. By modifying the local environment, these cells have their behavior and differentiation potential affected, because when primary odontoblasts die, they are able to undergo differentiation into cells called odontoblast-like cells, which start to secrete dentin matrix and deposit it in the form of repair dentin (1, 2, 70, 71). It should be noted that mesenchymal stem cells are potentially immunoregulatory structures, endowed with anti-inflammatory function, with the capacity for self-renewal and multilineage differentiation, and capable of producing structures similar to the original ones in the dental pulp (72). Thus, the immune response directed to infectious processes involves complex molecular mechanisms with coordinated actions, being the eicosanoids lipid mediators derived from arachidonic acid capable of regulating homeostatic and inflammatory processes. However,

the amount of eicosanoids produced is in dependence on the activation state and the physiological condition of the tissue (73).

Eicosanoids regulate the innate immune response by presenting immunomodulatory properties, highlighting the production of PGE<sub>2</sub> (64). PGE<sub>2</sub> is able to bind to 4 different types of receptors (EP1, EP2, EP3, and EP4, also known as PTGR1, PTGER2, PTGER3, and PTGER4). The EP1 receptor is coupled to the G<sub>q</sub> protein, and once activated, increases the intracellular calcium concentration. EP2 and EP4 receptors are coupled to the G<sub>s</sub> protein and, upon activation, are able to increase intracellular cyclic AMP (cAMP). On the other hand, the EP3 receptor is coupled to the G<sub>i</sub> protein and, unlike the EP2 and EP4 receptors, decreases cAMP formation. However, PGs generated from COX-2 and localized to the nuclear membrane can control nuclear pathways through interaction with peroxisome proliferator-activated receptor (PPAR), which regulates nuclear events of cell growth and survival (74, 75). This event occurs in some cases when PGs and their metabolites bind on these nuclear receptors, and PGE<sub>2</sub> can indirectly activate the PPAR $\delta$  receptor (75).

## Involvement of lipid inflammatory mediators in biomineralization

Not only the modulation of cell proliferation was affected by lipid inflammatory mediators, but also the expression of important genes inducing dentinal matrix mineralization (76, 77). Previously, PGE<sub>2</sub> has already been shown to have an anabolic effect on osteoblast proliferation and differentiation and induced IBSP transcription (78).

Growth factors such as BMP-2 and bioactive molecules from the dentinal matrix are important signaling molecules for dental pulp stem cells, stimulating them to differentiate into odontoblast-like cells (79). BMP2 protein is essential for the control of dentinal matrix mineralization, and is correlated to the differentiation of dental pulp cells (80).

PGE<sub>2</sub> induces BMP-2 production in culture of stem cells extracted from tendon, and consequently promote differentiation into cells of the osteoblastic lineage (81). On the other hand, LTB<sub>4</sub> is known to favor the resorption process in bone tissue, as it recruits clastic cells and inhibits the blast differentiation process, even in the presence of stimulation with BMP-2 (82).

In addition to the expression of *Ibsp* and *Bmp2*, the transcriptional factor *Runx2* also had gene expression stimulated by PGE<sub>2</sub> within the first 6 h of stimulation (76). *Runx2* is the main gene controlling the differentiation process of odontoblasts and is expressed in odontoblast-like cells and in dental pulp stem cells in the region of reparative dentin deposition (83), which makes it the transcriptional factor promoting the differentiation

of pulpal stem cells so that they are able to form reparative dentin (84).

The literature reports that this mediator has its production increased in cases of experimentally induced pulpal inflammation (65) and also was able to increase *Runx2* gene expression (76). This data turns out to be a great finding, because during the inflammatory process PGE<sub>2</sub> has its production increased through pro-inflammatory cytokines (85) which reveals that this mediator is present in the inflammatory environment in the very first hours. This suggests that PGE<sub>2</sub> is an early inducer mediator of the mineralizing response, because by increasing the expression of *Runx2* it tends to favor the deposition of mineralizing matrix, since it is an important transcriptional factor in the regulation of stem cell differentiation and formation of the dental organ (86). PGE<sub>2</sub>, in fact, has already revealed a dual role by participating in both resorption and formation of this bone tissue (87). Similar to what was observed in undifferentiated dental pulp cells, human periodontal ligament cells cultured in osteogenic medium and stimulated with PGE<sub>2</sub> at different molarities had increased expression of *RUNX2*, which demonstrates that PGE<sub>2</sub> is able to modulate the expression of this gene involved in osteogenic regulation (76, 88).

## Conclusion

It is well-established that pulp and dentin constitute a single unit, capable of responding to external stimuli, which makes the dentin-pulp complex an important strategic and dynamic barrier to the various injuries suffered by teeth. Given the above, odontoblasts are cells of fundamental importance for the initiation and amplification of innate immune response events in the search for protection of the pulp organ in the presence of pathogens, as well as undifferentiated cells present in the pulp and precursors of odontoblasts, both being able to respond to injury by modulating the immune response.

## Author contributions

FL-S, GAC, and LSS wrote the first draft of the manuscript. GCCL, MFMA, LHF, and FWGP-S revised the manuscript. All authors approved the final version of this review.

## Funding

This study was supported by The São Paulo Research Foundation (FAPESP grants #2019/00204-1 to FWGP-S and #2014/07125-6 to LHF) and Conselho Nacional de

Desenvolvimento Científico e Tecnológico (CNPq grant #303259/2020-5 to LHF and PIBIC fellowship to GAC).

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## References

- Téclès O, Laurent P, Zygouritsas S, Burger AS, Camps J, Dejou J, et al. (2005). Activation of human dental pulp progenitor/stem cells in response to odontoblast injury. *Arch Oral Biol.* (2004) 50: 103–8. doi: 10.1016/j.archoralbio.2004.11.009
- Chogle SM, Goodis HE, Kinaia BM. Pulpal and periradicular response to caries: current management and regenerative options. *Dent Clin North Am.* (2012) 57:1–10. doi: 10.1016/j.cden.2012.05.003
- Chen H, Fu H, Wu X, Duan Y, Zhang S, Hu H, et al. Regeneration of pulp-dentinal-like complex by a group of unique multipotent CD24a<sup>+</sup> stem cells. *Sci Adv.* (2020) 6:16154. doi: 10.1126/sciadv.aay1514
- Farges JC, Keller JF, Carrouel F, Durand SH, Romeas A, Bleicher F, et al. Odontoblasts in the dental pulp immune response. *J Exp Zool B Mol Dev Evol.* (2009) 312B 425–36. doi: 10.1002/jez.b.21259
- Cooper PR, Holder MJ, Smith AJ. Inflammation and regeneration in the dentin-pulp complex: a double-edged sword. *J Endod.* (2014) 24:S46–51. doi: 10.1016/j.joen.2014.01.021
- Cooper PR, McLachlan JL, Simon S, Graham LW, Smith AJ. Mediators of inflammation and regeneration. *Adv Dent Res.* (2011). 23:290–5. doi: 10.1177/0022034511405389
- Smith AJ, Lesot H. Induction and regulation of crown dentinogenesis: embryonic events as a template for dental tissue repair? *Crit Rev Oral Biol Med.* (2001) 12:425–37. doi: 10.1177/10454411010120050501
- Bleicher F. Odontoblast physiology. *Exp Cell Res.* (2014) 325(2):65–71. doi: 10.1016/j.yexcr.2013.12.012
- Thesleff I. Epithelial-mesenchymal signalling regulating tooth morphogenesis. *J Cell Sci.* (2003) 116:1647–8. doi: 10.1242/jcs.00410
- Larmas MA, Sándor GK. Solid nomenclature: the bedrock of science. Similarities and dissimilarities in phenomena and cells of tooth and bone ontogeny. *Anat Rec.* (2013) 296:564–7. doi: 10.1002/ar.22671
- Cunha da, da Costa-Neves JM, Kerkis A, da Silva IMC. Pluripotent stem cell transcription factors during human odontogenesis. *Cell Tissue Res.* (2013) 353:435–41. doi: 10.1007/s00441-013-1658-y
- Couve E, Osorio R, Schmachtenberg O. The amazing odontoblast: activity, autophagy, and aging. *J Dent Res.* (2013) 92:765–72. doi: 10.1177/0022034513495874
- Horst OV, Tompkins KA, Coats SR, Brahm PH, Darveau RP, Dale BA, et al. TGF-beta1 Inhibits TLR-mediated odontoblast responses to oral bacteria. *J Dent Res.* (2009) 88:333–8. doi: 10.1177/0022034509334846
- Kuzynski M, Goss M, Bottini M, Yadav MC, Mobley C, Winters T, et al. Dual role of the Trps1 transcription factor in dentin mineralization. *J Biol Chem.* (2014) 289:27481–93. doi: 10.1074/jbc.M114.550129
- Qin C, D'Souza R, Feng JQ. Dentin matrix protein 1 (DMP1): new and important roles for biomineralization and phosphate homeostasis. *J Dent Res.* (2007) 86:1134–41. doi: 10.1177/154405910708601202
- Woltgens JH, Lyaruu DM, Bronckers AL, Bervoets TJ, Van Duin M. Biomineralization during early stages of the developing tooth in vitro with special reference to secretory stage of amelogenesis. *Int J Dev Biol.* (1995) 39: 203–12.
- Hoylaerts ME, Van Kerckhoven S, Kiffer-Moreira T, Sheen C, Narisawa S, Millán JL, et al. Functional significance of calcium binding to tissue-nonspecific alkaline phosphatase. *PLoS ONE.* (2015) 10:e0119874. doi: 10.1371/journal.pone.0119874

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Chen L, Jacquet R, Lowder E, Landis WJ. Refinement of collagen-mineral interaction: a possible role for osteocalcin in apatite crystal nucleation, growth and development. *Bone.* (2015) 5:21. doi: 10.1016/j.bone.2014.09.021
- Moradian-Oldak J, George A. Biomineralization of enamel and dentin mediated by matrix proteins. *J Dent Res.* (2021) 100:1020–9. doi: 10.1177/00220345211018405
- George A, Gui J, Jenkins NA, Gilbert DJ, Copeland NG, Veis A, et al. In situ localization and chromosomal mapping of the AG1 (Dmp1) gene. *J Histochem Cytochem.* (1994) 42:1527–31. doi: 10.1177/42.12.7983353
- He G, Dahl T, Veis A, George A. Nucleation of apatite crystals in vitro by self-assembled dentin matrix protein 1. *Nat Mater.* (2003) 2:552–8. doi: 10.1038/nmat945
- von Marschall Z, Fisher LW. Dentin sialophosphoprotein (DSPP) is cleaved into its two natural dentin matrix products by three isoforms of bone morphogenetic protein-1 (BMP1). *Matrix Biol.* (2010) 29:295–303. doi: 10.1016/j.matbio.2010.01.002
- Tsuchiya S, Simmer JP, Hu JC, Richardson AS, Yamakoshi F, Yamakoshi Y, et al. Astacin proteases cleave dentin sialophosphoprotein (Dspp) to generate dentin phosphoprotein (Dpp). *J Bone Miner Res.* (2011) 26:220–8. doi: 10.1002/jbmr.202
- Zhu Q, Gibson MP, Liu Q, Liu Y, Lu Y, Wang X, et al. Proteolytic processing of dentin sialophosphoprotein (DSPP) is essential to dentinogenesis. *J Biol Chem.* (2012) 287:30426–35. doi: 10.1074/jbc.M112.388587
- Suzuki S, Sreenath T, Haruyama N, Honeycutt C, Terse A, Cho A, et al. Dentin sialoprotein and dentin phosphoprotein have distinct roles in dentin mineralization. *Matrix Biol.* (2009) 28:221–9. doi: 10.1016/j.matbio.2009.03.006
- Suzuki S, Haruyama N, Nishimura F, Kulkarni AB. Dentin sialophosphoprotein and dentin matrix protein-1: Two highly phosphorylated proteins in mineralized tissues. *Arch Oral Biol.* (2012) 57:1165–75. doi: 10.1016/j.archoralbio.2012.03.005
- Gallorini M, Krifka S, Widbiller M, Schröder A, Brochhausen C, Cataldi A, et al. Distinguished properties of cells isolated from the dentin-pulp interface. *Ann Anat.* (2021) 234:151628. doi: 10.1016/j.aanat.2020.151628
- Prasad M, Butler WT, Qin C. Dentin sialophosphoprotein in biomineralization. *Connect Tissue Res.* (2010) 51:404–17. doi: 10.3109/03008200903329789
- Staines KA, MacRae VE, Farquharson C. The importance of the SIBLING family of proteins on skeletal mineralisation and bone remodelling. *J Endocrinol.* (2012) 214:241–55. doi: 10.1530/JOE-12-0143
- Yamashiro T, Tummers M, Thesleff I. Expression of bone morphogenetic proteins and Msx genes during root formation. *J Dent Res.* (2003) 82:172–6. doi: 10.1177/154405910308200305
- Yang J, Ye L, Hui TQ, Yang DM, Huang DM, Zhou XD, et al. Bone morphogenetic protein 2-induced human dental pulp cell differentiation involves p38 mitogen-activated protein kinase-activated canonical WNT pathway. *Int J Oral Sci.* (2015) 7:95–102. doi: 10.1038/ijos.2015.7
- Camilleri S, McDonald, F. (2006). Runx2 and dental development. *Eur J Oral Sci.* doi: 10.1111/j.1600-0722.2006.00399.x
- Lézet F, Thomas B, Hotton D, Forest N, Orestes-Cardoso S, Robert B, et al. Biomineralization, life-time of odontogenic cells and differential expression of the two homeobox genes MSX-1 and DLX-2 in transgenic mice. *J Bone Miner Res.* (2000) 15:430–41. doi: 10.1359/jbmr.2000.15.3.430

34. Cobourne MT. Familial human hypodontia—is it all in the genes? *Br Dent J*. (2007) 203:203–8. doi: 10.1038/bdj.2007.732
35. Feng XY, Zhao YM, Wang WJ, Ge LH. Mx1 regulates proliferation and differentiation of mouse dental mesenchymal cells in culture. *Eur J Oral Sci*. (2013) 121:412–20. doi: 10.1111/eos.12078
36. Chen J, Sasaguri K, Sodek J, Aufdemorte TB, Jiang H, Thomas HF, et al. Enamel epithelium expresses bone sialoprotein (BSP). *Eur J Oral Sci*. (1998) 1:331–6. doi: 10.1111/j.1600-0722.1998.tb02194.x
37. Takai H, Matsumura H, Matsui S, Kim KM, Mezawa M, Nakayama Y, et al. Unliganded estrogen receptor alpha stimulates bone sialoprotein gene expression. *Gene*. (2014) 539:50–7. doi: 10.1016/j.gene.2014.01.063
38. Lombardi G, Perego S, Luzi L, Banfi G. A four-season molecule: osteocalcin. Updates in its physiological roles. *Endocrine*. (2015) 48:394–404. doi: 10.1007/s12020-014-0401-0
39. Papagerakis P, Bernal A, Mesbah M, Peuchmaur M, Malaval L, Nydegger J, et al. Investigation of osteocalcin, osteonectin, and dentin sialophosphoprotein in developing human teeth. *Bone*. (2002) 30:377–85. doi: 10.1016/S8756-3282(01)00683-4
40. Yang F, Xu N, Li D, Guan L, He Y, Zhang Y, et al. A feedback loop between RUNX2 and the E3 ligase SMURF1 in regulation of differentiation of human dental pulp stem cells. *J Endod*. (2014) 40:1579–86. doi: 10.1016/j.joen.2014.04.010
41. Pääkkönen V, Rusanen P, Hagström J, Tjäderhane L. Mature human odontoblasts express virus-recognizing toll-like receptors. *Int Endod J*. (2014) 47:934–41. doi: 10.1111/iej.12238
42. Widbillier M, Schmalz G. Endodontic regeneration: hard shell, soft core. *Odontology*. (2021) 109:303–12. doi: 10.1007/s10266-020-00573-1
43. Staquet MJ, Carrouel F, Keller JF, Baudouin C, Msika P, Bleicher F, et al. Pattern-recognition receptors in pulp defense. *Adv Dent Res*. (2011) 23:296–301. doi: 10.1177/0022034511405390
44. Durand SH, Flacher V, Roméas A, Carrouel F, Colomb E, Vincent C, et al. Lipoteichoic acid increases TLR and functional chemokine expression while reducing dentin formation in vitro differentiated human odontoblasts. *J Immunol*. (2006) 176:2880–7. doi: 10.4049/jimmunol.176.5.2880
45. Keller JF, Carrouel F, Colomb E, Durand SH, Baudouin C, Msika P, et al. Toll-like receptor 2 activation by lipoteichoic acid induces differential production of pro-inflammatory cytokines in human odontoblasts, dental pulp fibroblasts and immature dendritic cells. *Immunobiology*. (2009) 215:53–9. doi: 10.1016/j.imbio.2009.01.009
46. Veerayuthwilai O, Byers MR, Pham TT, Darveau RP, Dale BA. Differential regulation of immune responses by odontoblasts. *Oral Microbiol Immunol*. (2007) 22:5–13. doi: 10.1111/j.1399-302X.2007.00310.x
47. Liu Y, Gao Y, Zhan X, Cui L, Xu S, Ma D, et al. TLR4 activation by lipopolysaccharide and Streptococcus mutans induces differential regulation of proliferation and migration in human dental pulp stem cells *J Endod*. 40:1375–81. (2014) 015. doi: 10.1016/j.joen.2014.03.015
48. Farges JC, Alliot-Licht B, Baudouin C, Msika P, Bleicher F, Carrouel F, et al. Odontoblast control of dental pulp inflammation triggered by cariogenic bacteria. *Front Physiol*. (2013) 4:326. doi: 10.3389/fphys.2013.00326
49. Crean D, Godson C. (2015). Specialised lipid mediators and their targets. *Semin Immunol* 27:169–76. doi: 10.1016/j.smim.2015.05.002
50. An S, Chen Y, Yang T, Huang Y, Liu, Y. A role for the calcium-sensing receptor in the expression of inflammatory mediators in LPS-treated human dental pulp cells. *Mol Cell Biochem*. (2022) 3:4486. doi: 10.1007/s11010-022-04486-1
51. Funk CD. Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science*. (2001) 294:1871–5. doi: 10.1126/science.294.5548.1871
52. Muñoz-Carrillo JL, Vargas-Barboza JM, Villalobos-Gutiérrez PT, Flores-De La Torre JA, Vazquez-Alcaraz SJ, Gutiérrez-Coronado O. Effect of treatment with resiniferatoxin in an experimental model of pulpal inflammatory in mice. *Int Endod J*. (2021) 54:2099–112. doi: 10.1111/iej.13606
53. Uozumi N, Kume K, Nagase T, Nakatani N, Ishii S, Tashiro F, et al. Role of cytosolic phospholipase A2 in allergic response and parturition. *Nature*. (1997) 390:618–22. doi: 10.1038/37622
54. Haeggström JZ, Rinaldo-Matthis A, Wheelock CE, Wetterholm A. Advances in eicosanoid research, novel therapeutic implications. *Biochem Biophys Res Commun*. (2010) 396:135–9. doi: 10.1016/j.bbrc.2010.03.140
55. Guijas C, Rodríguez JP, Rubio JM, Balboa MA, Balsinde J. Phospholipase A2 regulation of lipid droplet formation. *Biochim Biophys Acta*. 1841:1661–71. doi: 10.1016/j.bbali.2014.10.004
56. Harizi H, Corcuff JB, Gualde N. Arachidonic-acid-derived eicosanoids: roles in biology and immunopathology. *Trends Mol Med*. (2008) 14:461–469. doi: 10.1016/j.molmed.2008.08.005
57. Hammond VJ, O'Donnell VB. Esterified eicosanoids: generation, characterization, and function. *Biochim Biophys Acta*. (2011) 1818:2403–12. doi: 10.1016/j.bbame.2011.12.013
58. Powell WS, Rokach J. Biosynthesis, biological effects, and receptors of hydroxyeicosatetraenoic acids (HETEs) and oxo-eicosatetraenoic acids (oxo-ETEs) derived from arachidonic acid. *Biochim Biophys Acta*. (2014) 1851:340–55. doi: 10.1016/j.bbali.2014.10.008
59. Rådmark O, Werz O, Steinhilber D, Samuelsson B. 5-Lipoxygenase, a key enzyme for leukotriene biosynthesis in health and disease. *Biochim Biophys Acta*. (2014) 1851:331–9. doi: 10.1016/j.bbali.2014.08.012
60. Back M, Powell WS, Dahlen SE, Drazen JM, Evans JF, Serhan CN, et al. Update on leukotriene, lipoxin and oxo-eicosanoid receptors: IUPHAR Review 7. *Br J Pharmacol*. (2014) 171:3551–74. doi: 10.1111/bph.12665
61. Kawahara K, Hohjoh H, Inazumi T, Tsuchiya S, Sugimoto Y. Prostaglandin E2-induced inflammation: Relevance of prostaglandin E receptors. *Biochim Biophys Acta*. (2015) 1851:414–21. doi: 10.1016/j.bbali.2014.07.008
62. Clària J. Cyclooxygenase-2 biology. *Curr Pharm Des*. (2003) 9:2177–90. doi: 10.2174/1381612033454054
63. Legler, D. F., and Bruckner, M. Uetz-von Allmen, E., Krause, P. Prostaglandin E2 at new glance: novel insights in functional diversity offer therapeutic chances. *Int J Biochem Cell Biol*. (2010) 42:198–201. doi: 10.1016/j.biocel.2009.09.015
64. Rodríguez M, Domingo E, Municio C, Alvarez Y, Hugo E, Fernández N, et al. Polarization of the innate immune response by prostaglandin E2: a puzzle of receptors and signals. *Mol Pharmacol*. (2014) 85:187–97. doi: 10.1124/mol.113.089573
65. Okiji T, Morita I, Kobayashi C, Sunada I, Murota S. Arachidonic-acid metabolism in normal and experimentally-inflamed rat dental pulp. *Arch Oral Biol*. (1987) 32:723–7. doi: 10.1016/0003-9969(87)90116-6
66. Okiji T, Morita I, Suda H, Murota S. Pathophysiological roles of arachidonic acid metabolites in rat dental pulp. *Proc Finn Dent Soc*. (1992) 88(Suppl 1):433–8.
67. Ribeiro-Santos FR, Silva GGD, Petean IBF, Arnez MFM, Silva LABD, Faccioli LH, et al. Periapical bone response to bacterial lipopolysaccharide is shifted upon cyclooxygenase blockage. *J Appl Oral Sci*. (2019) 27:e20180641. doi: 10.1590/1678-7757-2018-0641
68. Paula-Silva FWG, Ribeiro-Santos FR, Petean IBF, Manfrin Arnez ME, Almeida-Junior LA, Carvalho FK, et al. Root canal contamination or exposure to lipopolysaccharide differentially modulate prostaglandin E 2 and leukotriene B 4 signaling in apical periodontitis. *J Appl Oral Sci*. (2020) 28:699. doi: 10.1590/1678-7757-2019-0699
69. Cooper PR, Takahashi Y, Graham LW, Simon S, Imazato S, Smith AJ, et al. Inflammation-regeneration interplay in the dentine-pulp complex. *J Dent Sep*. (2010) 38:687–97. doi: 10.1016/j.jdent.2010.05.016
70. Yalvac ME, Ramazanoglu M, Rizvanov AA, Sahin F, Bayrak OF, et al. Isolation and characterization of stem cells derived from human third molar tooth germs of young adults: implications in neo-vascularization, osteo-, adipo- and neurogenesis. *Pharmacogenomics J*. (2010) 10:105–13. doi: 10.1038/tpj.2009.40
71. Chmielewski F, Jeanneau C, Dejou J, About I. Sources of dentin-pulp regeneration signals and their modulation by the local microenvironment. *J Endod*. (2014) 9:519–25. doi: 10.1016/j.joen.2014.01.012
72. Li Z, Jiang CM, An S, Cheng Q, Huang YF, Wang YT, et al. Immunomodulatory properties of dental tissue-derived mesenchymal stem cells. *Oral Dis*. (2014) 1:25–34. doi: 10.1111/odi.12086
73. Dennis EA, Norris PC. Eicosanoid storm in infection and inflammation. *Nat Rev Immunol*. (2015) 15:511–23. doi: 10.1038/nri3859
74. Romano M, Claria J. Cyclooxygenase-2 and 5-lipoxygenase converging functions on cell proliferation and tumor angiogenesis: implications for cancer therapy. *FASEB J*. (2003) 17:1986–95. doi: 10.1096/fj.03-0053rev
75. Wang D, Dubois RN. Eicosanoids and cancer. *Nat Rev Cancer*. (2010) 10:181–93. doi: 10.1038/nrc2809
76. Lorencetti-Silva F, Pereira PAT, Meirelles AFG, Faccioli LH, Paula-Silva FWG. Prostaglandin E2 induces expression of mineralization genes by undifferentiated dental pulp cells. *Braz Dent J30*. (2019) 201–7. doi: 10.1590/0103-6440201902542
77. da Silva FL, de Campos Chaves Lamarque G, de Oliveira FMMPC, Nelson-Filho P, da Silva LAB, Segato RAB, et al. Leukotriene B4 loaded in microspheres regulate the expression of genes related to odontoblastic differentiation and biomineralization by dental pulp stem cells. *BMC Oral Health*. (2022) 23:22:45. doi: 10.1186/s12903-022-02083-8
78. Samoto H, Shimizu E, Matsuda-Honjyo Y, Saito R, Nakao S, Yamazaki M, et al. Prostaglandin E2 stimulates bone sialoprotein (BSP) expression

- through cAMP and fibroblast growth factor 2 response elements in the proximal promoter of the rat BSP gene. *J Biol Chem.* (2003) 278:28659–67. doi: 10.1074/jbc.M300671200
79. Lin LM, Rosenberg PA. Repair and regeneration in endodontics. *Int Endod J.* (2011) 44:889–906. doi: 10.1111/j.1365-2591.2011.01915.x
80. Zhang W, Zhang X, Ling J, Liu W, Zhang X, Ma J, et al. Proliferation and odontogenic differentiation of BMP2 gene transfected stem cells from human tooth apical papilla: an in vitro study. *Int J Mol Med.* (2014) 34:1004–12. doi: 10.3892/ijmm.2014.1862
81. Zhang J, Wang JH. BMP-2 mediates PGE(2) -induced reduction of proliferation and osteogenic differentiation of human tendon stem cells. *J Orthop Res.* (2012) 30:47–52. doi: 10.1002/jor.21485
82. Traianedes K, Dallas MR, Garrett IR, Mundy GR, Bonewald LF. 5-Lipoxygenase metabolites inhibit bone formation in vitro. *Endocrinology.* (1998) 139:3178–84. doi: 10.1210/endo.139.7.6115
83. Daltoé MO, Paula-Silva FW, Faccioli LH, Gatón-Hernández PM, Rossi De, Bezerra Silva A, et al. Expression of mineralization markers during pulp response to biodentine and mineral trioxide aggregate. *J Endod.* (2016) 42:596–603. doi: 10.1016/j.joen.2015.12.018
84. Han N, Zheng Y, Li R, Li X, Zhou M, Niu Y, et al.  $\beta$ -catenin enhances odontoblastic differentiation of dental pulp cells through activation of Runx2. *PLoS ONE.* (2014) 9:e88890. doi: 10.1371/journal.pone.0088890
85. Yu KR, Lee JY, Kim HS, Hong IS, Choi SW, Seo Y, et al. A p38 MAPK-mediated alteration of COX-2/PGE2 regulates immunomodulatory properties in human mesenchymal stem cell aging. *PLoS ONE.* (2014) 9:2426. doi: 10.1371/journal.pone.0102426
86. Wen J, Tao R, Ni L, Duan Q, Lu Q. Immunolocalization and expression of Runx2 in tertiary dentinogenesis. *Hybridoma.* (2010) 29:195–9. doi: 10.1089/hyb.2009.0120
87. Yoshida K, Oida H, Kobayashi T, Maruyama T, Tanaka M, Katayama T, et al. Stimulation of bone formation and prevention of bone loss by prostaglandin E EP4 receptor activation. *Proc Natl Acad Sci USA.* (2002) 99:4580–5. doi: 10.1073/pnas.062053399
88. Manokawinchoke J, Pimkhaokhum A, Everts V, Pavasant P. Prostaglandin E2 inhibits in-vitro mineral deposition by human periodontal ligament cells via modulating the expression of TWIST1 and RUNX2. *J Periodontol Res.* (2014) 49:777–84. doi: 10.1111/jre.12162