



# Geologic Controls on Erosion Mechanism on the Alaska Beaufort Coast

Thomas M. Ravens<sup>1\*†</sup> and Sasha Peterson<sup>2‡</sup>

<sup>1</sup>Department of Civil Engineering, University of Alaska Anchorage, Anchorage, AK, United States, <sup>2</sup>Department of Geological Sciences, University of Alaska Anchorage, Anchorage, AK, United States

## OPEN ACCESS

### Edited by:

Scott Raymond Dallimore,  
Geological Survey of Canada, Canada

### Reviewed by:

Henry Patton,  
Arctic University of Norway, Norway  
Jennifer Frederick,  
Sandia National Laboratories,  
United States

### \*Correspondence:

Thomas M. Ravens  
tmravens@alaska.edu

### †ORCID

Thomas M. Ravens  
orcid.org/0000-0002-4613-4632

### ‡Present address:

Sasha Peterson,  
Department of Environmental Science  
and Engineering, University of Texas  
at El Paso, El Paso, TX, United States

### Specialty section:

This article was submitted to  
Cryospheric Sciences,  
a section of the journal  
Frontiers in Earth Science

**Received:** 12 April 2021

**Accepted:** 24 September 2021

**Published:** 15 October 2021

### Citation:


Ravens TM and Peterson S (2021)  
Geologic Controls on Erosion  
Mechanism on the Alaska  
Beaufort Coast.  
Front. Earth Sci. 9:693824.  
doi: 10.3389/feart.2021.693824

Two prominent arctic coastal erosion mechanisms affect the coastal bluffs along the North Slope of Alaska. These include the niche erosion/block collapse mechanism and the bluff face thaw/slump mechanism. The niche erosion/block collapse erosion mechanism is dominant where there are few coarse sediments in the coastal bluffs, the elevation of the beach below the bluff is low, and there is frequent contact between the sea and the base of the bluff. In contrast, the bluff face thaw/slump mechanism is dominant where significant amounts of coarse sediment are present, the elevation of the beach is high, and contact between the sea and the bluff is infrequent. We show that a single geologic parameter, coarse sediment areal density, is predictive of the dominant erosion mechanism and is somewhat predictive of coastal erosion rates. The coarse sediment areal density is the dry mass (g) of coarse sediment (sand and gravel) per horizontal area (cm<sup>2</sup>) in the coastal bluff. It accounts for bluff height and the density of coarse material in the bluff. When the areal density exceeds 120 g cm<sup>-2</sup>, the bluff face thaw/slump mechanism is dominant. When the areal density is below 80 g cm<sup>-2</sup>, niche erosion/block collapse is dominant. Coarse sediment areal density also controls the coastal erosion rate to some extent. For the sites studied and using erosion rates for the 1980–2000 period, when the sediment areal density exceeds 120 g cm<sup>-2</sup>, the average erosion rate is low or 0.34 ± 0.92 m/yr. For sediment areal density values less than 80 g cm<sup>-2</sup>, the average erosion rate is higher or 2.1 ± 1.5 m/yr.


**Keywords:** arctic, coastal erosion, mechanism, coarse sediment, areal density

## INTRODUCTION

The Arctic is experiencing high and accelerating coastal erosion rates. For example, Mars and Houseknecht (2007) used remote sensing techniques to study coastal erosion-derived land loss on a 60-km segment of the Beaufort Sea coast (between Drew Point and Cape Halkett, Alaska, **Figure 1**) and found that the amount of land loss was significantly greater in 1985–2005 (1.08 km<sup>2</sup> yr<sup>-1</sup>) relative to the loss in 1955–1985 (0.48 km<sup>2</sup> yr<sup>-1</sup>). Jones et al. (2009) working in the same area determined that the average rate of erosion increased from 6.8 m yr<sup>-1</sup> (1955–1979), to 8.7 m yr<sup>-1</sup> (1979–2002), and to 13.6 m yr<sup>-1</sup> (2002–2007). Erosion rates are high in this location because of the high ice content of the coastal bluffs and the absence of coarse material (sand and gravel). At other locations, erosion rates are often lower but still accelerating. For example, on Barter Island, where coastal bluffs contain significant amounts of coarse material, bluff retreat rate averaged 1.8 myr<sup>-1</sup> between 1955 and 2004 and 3.8 m yr<sup>-1</sup> between 2004 and 2010 (Gibbs et al., 2010). Erosion rates are generally accelerating



**FIGURE 1** | Map of the north coast of Alaska showing color-coded shoreline change rates for the period circa-1940s (1947 and 1949) to circa-2000s (1997–2012, Gibbs and Richmond, 2015).




**FIGURE 2** | Conceptual model of the niche erosion/block collapse erosion mechanism (from Ravens et al., 2012).

because of 1) greater spatial extent of open water, which allows for the generation of larger waves, 2) greater open water period, and 3) increased rate of coastal permafrost thaw (Barnhart et al., 2014a; Barnhart et al., 2014b; Frederick et al., 2016). Erosion threatens coastal infrastructure throughout the Arctic including governmental assets and community infrastructure. The US Army Corps of Engineers (2009) has designated 26 Alaska communities (including Barrow) “Priority Action Communities” due to the threat of erosion.

A number of arctic coastal erosion mechanisms affecting high coastal bluffs in the Arctic have been identified including niche erosion/block collapse (prevalent in the Drew Point area (Ravens et al., 2012; Barnhart et al., 2014a)) and bluff face thaw/slump (also referred to as translational-shear ice-thaw, Gibbs et al., 2013, and thermal denudation, Barankaya et al., 2021). The erosion mechanisms affecting Arctic coastal bluffs differ from the erosion of non-Arctic bluffs (e.g., Carter and Guy 1988) because of the role played by thermal processes in



**FIGURE 3** | Photos of (A) an erosional niche from Elson Lagoon Alaska and (B) a fallen block by Drew Point, Alaska (image courtesy of Christopher Arp of the Alaska Science Center, U.S. Geological Survey).



**FIGURE 4** | Sketch of the bluff cross-section assumed by Hoque and Pollard (2016) in their analysis of overturning failure.




**FIGURE 5** | Photo showing material that has slumped onto the beach face following bluff face thaw at Barter Island (2011 image courtesy of Li Erikson, U.S. Geological Survey). The bluff height is about 10 m and the sediment areal density is about 600 g/cm<sup>2</sup>, based on USGS data. Interestingly, the photo was taken soon after the 2008 niche erosion/block collapse event and the niche is still in evidence.

the Arctic. With the niche erosion/block collapse erosion mechanism, typically a small beach is present before the bluff (Figures 2, 3). During a storm surge event, waters rise allowing contact between sea and the base of the bluff. Waves

and currents thermally and mechanically carve a niche at the base of the bluff (Kobayashi 1985). Niche growth undermines the bluffs leading to block collapse due to an overturning failure (Hoque and Pollard 2016). The lower failure plane intersects with a shore parallel ice wedge (Figure 4). The upper failure plane is at interface of the ice wedge and the soil. The failure is governed by the tensile strength of the frozen soil, as well as the niche depth, the ice wedge location, and the depth of the ice wedge. Niche erosion/block collapse is the predominant erosion mechanism in settings where the coastal bluffs have high ice content (~70%, Ping et al., 2011), and where the bluffs lack significant amounts of coarse material (sand and gravel). The lack of coarse material leads to a low elevation beach at the base of the bluff and frequent contact between the sea and the coastal bluffs (Ravens et al., 2011; Ravens and Peterson 2018).

Bluff face thaw/slump is the predominant erosion mechanism in settings where significant amounts of coarse sediments are common (e.g., at Barter Island, Ravens et al., 2011; Ravens and Peterson 2018). With significant amounts of coarse sediments in the coastal bluffs, the elevation of the beach before the bluff is relatively high (1–2 m above mean sea level) and contact between the sea and the base of the bluff—and niche erosion—is infrequent. For example, data provided by the USGS (Ann Gibbs, personal communication) indicates that only a single significant niche erosion/block collapse event occurred in the 1955–2010 time period at Barter Island which has significant amounts of coarse sediments (Figure 5). The bluff face warms due to the combined effect of a number of heat transfer processes including solar (shortwave) radiation, longwave radiation emission from the Earth's surface, absorption of downward longwave radiation from the atmosphere, sensible heat flux, and latent heat flux (Westermann et al., 2009; Ravens and Ulmgren, 2020). When the bluff face is warmed sufficiently, it thaws and material slumps to the beach face (Figures 5,6). Relatively small storms (e.g., the 1-year return period storm) are sufficient to remove the sediment that accumulates on the beach (Ravens et al., 2011).

Ravens et al. (2011) defined a parameter, the “coarse sediment areal density”, and they hypothesized that this parameter




**TABLE 1** | Photographic and geologic data used in the analysis.

Photo ID	Photo location		Erosion mechanism	Barrier island present	Ping et al. (2021) site	Average sediment density	Bluff height	Coarse material (sand) content	Coarse sediment areal density	Erosion rate
—	Latitude	Longitude	—	—	—	(g cm <sup>-3</sup> )	(cm)	(%)	(g cm <sup>-2</sup> )	(m/year)
IMG_9510	70.899	-153.367	Niche/block	N	BSC17	0.69	50	21.9	7.5	3.47
IMG_8113	70.1629	-145.845	Niche/block	N	BSC39	0.64	250	54.1	86.0	0.35
IMG_0238	71.02287	-154.623	Niche/block	N	BSC15	0.37	40	48.4	7.1	2.14
IMG_9428	70.78902	-152.271	Niche/block	N	BSC20	0.61	250	30.8	46.8	2.7
IMG_8136	70.04606	-145.447	Niche/block	Y	BSC40	0.57	280	47.3	76.1	0.12
IMG_0065	71.33132	-156.566	Niche/block	Y	BSC01	1.03	40	53.2	22.0	0.31
IMG_0087	71.29122	-156.438	Niche/block	Y	BSC02	0.52	230	32.6	39.3	1.56
IMG_0124	71.21429	-156.047	Niche/block	Y	BSC03	0.38	140	34.6	18.6	4.57
IMG_0184	71.12589	-155.548	Niche/block	Y	BSC04	0.62	160	54.3	53.9	2.25
IMG_8366	70.03766	-142.72	Bluff face thaw	N	BSC46	0.60	300	88.4	158.9	0.54
IMG_8210	69.99457	-144.546	Bluff face thaw	N	BSC42	0.66	200	62.7	87.7	0.26
IMG_8385	69.98949	-142.556	Bluff face thaw	N	BSC47	0.70	320	74.4	238.0	0.96
IMG_8470	69.65694	-141.039	Bluff face thaw	N	BSC50	0.54	350	48.7	91.5	3.88
IMG_8772	70.00185	-144.828	Bluff face thaw	N	BSC41b	0.81	400	11.6	82.7	0.36
IMG_9327	70.55583	-151.709	Bluff face thaw	N	BSC24	1.53	320	90.1	441.7	0.24
IMG_7869	70.4919	-149.226	Bluff face thaw	Y	BSC31	1.12	200	90.3	202.1	-1.55
IMG_7924	70.40772	-148.778	Bluff face thaw	Y	BSC32	0.66	260	71.3	122.6	1.24
IMG_8225	70.03146	-144.319	Bluff face thaw	Y	BSC42	0.66	200	62.7	87.7	0.26
IMG_8241	70.08234	-144.002	Bluff face thaw	Y	BSC43	1.32	170	83.0	186.9	-0.24
IMG_7571	70.33116	-148.08	Bluff face thaw	Y	BSC34	1.03	300	62.5	193.2	0.33


determined whether the bluffs at a given coastal site were controlled by niche erosion/block collapse or by bluff face thaw/slumping. The sediment areal density is the dry mass of coarse sediment (sand and gravel) contained in a column of bluff sediment/soil per unit horizontal area (g cm<sup>-2</sup>). If there was a virtual column in the bluff extending from mean sea level to the

bluff top, the coarse sediment areal density would be the dry mass of coarse sediment (sand and gravel) per unit horizontal area in the column. In this paper, we test this hypothesis by examining the extent to which coarse sediment areal density can predict coastal erosion mechanism. We also examine the relationship between coarse sediment areal density and coastal erosion rate.



## METHODOLOGY


Coastal locations with both sediment data and aerial photo data from the north coast of Alaska between Utqiagvik (formerly




Barrow) and the Canadian border were sought. Data on sediment grain size distribution (percent sand, silt, and clay) as a function of depth into the bluffs, sediment bulk density, and bluff height were obtained from 22 coastal sites according to Ping et al. (2011). Note, Ping et al. (2011) did not report on the presence of gravel so we concluded that it was negligible in their samples. However, the USGS, working at their Barter Island site, found significant gravel (Gibbs et al., 2010). The samples were collected from undisturbed areas between ice wedges after removal of slumped material. We examined oblique aerial photos from Gibbs and Richmond (2009) at locations proximal to the sites with sediment data to determine if the coastal erosion mechanism was niche erosion/block collapse or bluff face thaw/slump (**Table 1**). On average, the distance between location with sediment data and photos was about 6 km. For each photo, sand and gravel content data from one proximal core or bluff sample was used to determine the sediment areal density (**Figure 7**). Locations experiencing niche erosion/block collapse were readily determined based on the characteristic erosional blocks (**Figure 8**). Locations dominated by bluff face thaw/slump were evident based on the presence of a high elevation beach before the coastal bluff and the presence of material (e.g., vegetation) that was slumping on the bluff face (**Figure 9**). The coarse sediment areal density ( $\text{g cm}^{-2}$ ) was calculated as the product of the coarse sediment (sand and gravel) content (%), sediment bulk density ( $\text{g cm}^{-3}$ ) and the bluff height (cm), using data from Ping et al., 2011. The ice content of the bluffs was implicitly included in the sediment bulk density.

## RESULTS AND DISCUSSION

The locations of the 19 coastal sites subject to analysis, as well as the erosion mechanisms attributed to those sites based on the analysis of the aerial photos, are shown in **Figure 10**. It is noteworthy that the majority of the sites experiencing niche erosion/block collapse are on the western side of the study




**FIGURE 10 |** Map of the north coast of Alaska showing the locations of the coastal sites studied as well as the erosion mechanism attributed to those sites. Base map imagery courtesy of Esri.



**FIGURE 11 |** A histogram showing the frequency of occurrence of the niche erosion/block collapse erosion mechanism and bluff face thaw/slump mechanism as a function of coarse sediment areal density.

domain, whereas the sites experiencing bluff face thaw/slump are mainly on the eastern side. Note also that there was relatively little variation of erosion mechanism with position according to our analysis. The frequency of occurrence of the niche erosion/block collapse mechanism and the bluff face thaw/slump mechanism relative to the coarse sediment areal density ( $\text{g cm}^{-2}$ , **Figure 11**)



**FIGURE 12 |** Dependence of coastal erosion rates for the 1980–2000 time period on coarse sediment areal density, for sites experiencing niche erosion/block collapse and bluff face thaw/slump. Note, the figure provides data on coastal sites that are protected by barrier islands as well as ones without protection as indicated in the legend. Trend lines are provided for sites with niche erosion/block collapse (orange line,  $R^2 = 0.37$ ) as well as considering all sites (black line,  $R^2 = 0.25$ ). For bluff face thaw/slump sites, the correlation was negligible ( $R^2 = 0.09$ ) and no trend line is provided.

shows that with sediment areal density greater than  $120 \text{ g cm}^{-2}$ , the dominant erosion mechanism was bluff face thaw/slumping. With sediment areal density less than  $80 \text{ g cm}^{-2}$ , the dominant erosion mechanism was niche erosion/block collapse. One might wonder whether the erosion mechanism at specific sites, inferred based on the 2006 areal photos, might vary over time. It is noteworthy that, for example, Elson Lagoon, Drew Point, and

**TABLE 2** | Average erosion rates (for 1980–2000 period) for different ranges of coarse sediment areal density.

Range of sediment areal density (g cm <sup>-2</sup> )	Average erosion rate (m/yr) considering all locations	Average erosion rate (m/yr) considering sites with barrier island protection
>100 g cm <sup>-2</sup>	0.22±0.92	-0.06±1.17
<80 g cm <sup>-2</sup>	2.1±1.5	1.8±1.8

Barter Island have been subject to numerous research papers over the past few decades, and there has been no mention of a change in erosion mechanism although there are some caveats. First, Barter Island has eroded mainly due to bluff face thaw/slump (as expected due to its high sediment areal density), but it was subject to a significant niche erosion/block collapse event during a large 2008 storm (Gibbs et al., 2010; Ravens et al., 2011). Also, Gibbs et al. (2019) point out the seasonality of erosion mechanism. In early to mid-summer, there tends to be more bluff face thaw/slumping because of the high levels of solar (short wave) radiation. In the second half of the summer, after the thaw of sea ice, storm surges and wave action bring aggressive mechanical forces to the coast removing previously thawed and deposited material, and potentially causing niche erosion if the beach elevation is sufficiently low.

Erosion rates for the 1980–2000 period (from Ping et al., 2011) are plotted relative to coarse sediment areal density (Figure 12). For sediment areal density values greater than 120 g cm<sup>-2</sup> (coincident with the bluff face thaw/slump mechanism), erosion rates ranged from 1.24 m/yr to -1.55 m/yr (i.e., an accretion of 1.55 m/yr) with an average erosion rate of 0.34 ± 0.92 m/yr, Table 2). For sediment areal density values less than 80 g cm<sup>-2</sup> (coincident with the niche erosion/block collapse mechanism), erosion rates ranged from 4.57 to 0.12 m/yr with an average of 2.1 ± 1.5 m/yr. Thus, the presence of elevated coarse sediment areal density appears to control (or reduce) the coastal erosion rate.

Analysis was also performed to determine whether the presence of barrier island protection translated to reduced erosion rates for the two ranges of sediment areal density and the associated erosion mechanisms. For locations with coarse sediment areal density above 120 g cm<sup>-2</sup> (i.e., bluff face thaw/slump sites), the average erosion rate was reduced from 0.34 ± 0.92 m/yr (considering all sites) to -0.06 ± 1.17 m/yr, when only sites protected by barrier islands were considered (Table 2). For locations with coarse sediment areal density less than 80 g cm<sup>-2</sup> (i.e., the niche erosion/block collapse sites), the average erosion rate was reduced from 2.1 ± 1.5 m/yr (considering all sites) to 1.8 ± 1.8 m/yr, when only sites protected by barrier islands were considered (Table 1). Thus, barrier island protection appeared to provide a small reduction in erosion rate for all levels of coarse sediment areal density (and for both erosion mechanisms) though the reduction was less than the standard deviation. When all of the data (Figure 11) was subject to linear regression, the erosion rate (*ER*, m/yr, 1980–2000 period) was found to be somewhat correlated with coarse sediment areal density ( $\rho_{areal}$ , g cm<sup>-2</sup>) with an

$R^2$  of 0.20:  $ER = -0.0068 \rho_{areal} + 1.88$ . This indicates that the erosion rate is negatively correlated with sediment areal density.

A significant amount of the variance in the measured erosion rate could not be explained using the coarse sediment areal density alone. Various explanations for the unexplained variance exist. First, we had to work with a significant distance (order 1 km) between the location of the erosion measurement and the borehole from which the sediment areal density was derived. Given spatial non-uniformity in the coastal stratigraphy, it is reasonable to suggest that the sediment areal density at the location of the erosion measurement differed from the density at the borehole. Second, there are many environmental variables that affect erosion but were not included in the regression including: nearshore water surface elevation, nearshore wave condition, and nearshore water and air temperature. Third, the way in which environmental variables affect arctic coastal erosion can be quite complex as indicated by process-based approaches to determine erosion rate (Ravens et al., 2012; Barnhart et al., 2014a).

The analysis presented above focuses on the predictability of Arctic coastal erosion mechanism based on sediment areal density. However, once this relationship has been established, it is noteworthy that sediment character can be inferred to some extent based on the erosion mechanism. For example, in locations where niche erosion/block collapse is dominant, we can infer that the coarse sediment in the eroding bluffs is limited. Such insights could be used in sediment transport and other studies.

## CONCLUSION

The research presented here suggests that a single geologic parameter, the coarse sediment areal density, controls the dominant arctic coastal erosion mechanism of coastal bluffs on the North Slope (i.e., north coast) of Alaska. The coarse sediment areal density is the dry mass (g) of coarse sediment (sand) per horizontal area (cm<sup>2</sup>) in the coastal bluff. When the coarse sediment areal density exceeds 120 g cm<sup>-2</sup>, the bluff face thaw/slump erosion mechanism is dominant. When the coarse sediment areal density is below 80 g cm<sup>-2</sup>, the niche erosion/block collapse erosion mechanism is dominant. The coarse sediment areal density also has some influence on coastal erosion rates. Considering the 22 sites addressed in this study, the sediment areal density was found to have a controlling effect on erosion rate. Using erosion rates for the 1980–2000 period, when the sediment areal density exceeds 120 g cm<sup>-2</sup>, the average erosion rate was of 0.34 ± 0.92 m/yr. For sediment areal density values less than 80 g cm<sup>-2</sup>, the average erosion rate was as high as 2.1 ± 1.5 m/yr. Linear regression between coarse sediment areal density and erosion rate found that ~20% of the variance in erosion rate was explainable by coarse sediment areal density.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding author.

## AUTHOR CONTRIBUTIONS

TR conceived the overall manuscript and did the majority of the writing. SP did the analysis of the aerial photos and sediment data.

## REFERENCES

- Baranskaya, A., Novikova, A., Shabanova, N., Belova, N., MaznevJones, S. B., Ogorodov, S., et al. (2021). The Role of Thermal Denudation in Erosion of Ice-Rich Permafrost Coasts in an Enclosed Bay (Gulf of Kruzenstern, Western Yamal, Russia). *Front. Earth Sci.* 8, 2021. doi:10.3389/feart.2020.566227
- Barnhart, K. R., Anderson, R. S., Overeem, I., Wobus, C., Clow, G. D., and Urban, F. E. (2014a). Modeling Erosion of Ice-Rich Permafrost bluffs along the Alaskan Beaufort Sea Coast. *J. Geophys. Res. Earth Surf.* 119, 1155–1179. doi:10.1002/2013jf002845
- Barnhart, K. R., Overeem, I., and Anderson, R. S. (2014b). The Effect of Changing Sea Ice on the Physical Vulnerability of Arctic Coasts. *The Cryosphere* 8, 1777–1799. doi:10.5194/tc-8-1777-2014
- Carter, C. H., and E. Guy, D., Jr. (1988). Coastal Erosion: Processes, Timing and Magnitudes at the bluff Toe. *Mar. Geology*. 84, 1–17. Available at: <https://pubs.er.usgs.gov/publication/70014360>. doi:10.1016/0025-3227(88)90121-1
- Frederick, J. M., Thomas, M. A., BullJones, D. L. C. A., and Roberts, J. (2016). “The Arctic Coastal Erosion Problem,”. SAND2016-9762. doi:10.2172/1431492
- Gibbs, A. E., Erikson, L. H., Jones, B. M., and Richmond, B. M. (2010). *Characterizing Morphology and Erosional Trends of Permafrost Bluffs, Barter Island*. Alaska: AGU Fall Meeting. EP23A-0772.
- Gibbs, A. E., and Richmond, B. M. (2009). Oblique Aerial Photography of the Arctic Coast of Alaska, Nulavik to Demarcation Point, August 7 -10, 2006. *U.S. Geol. Surv. Data Ser. 436 6* (databases), 4, 2009. Available at: <http://pubs.usgs.gov/ds/436/>. doi:10.3133/ds436
- Gibbs, A. E., Richmond, B. M., Palaseanu-Lovejoy, M., Erikson, L. H., Jones, B. M., and Brock, J. (2013). *Remote Sensing of the Arctic Coast of Alaska Using Airborne Lidar Data*. San Francisco, CA: American Geophysical Union Fall Meeting. B51H-0406.
- Gibbs, A. E., and Richmond, B. M. (2015). National Assessment of Shoreline Change: Historical Change along the north Coast of Alaska, U.S.-Canadian Border to Icy Cape. *Geol. Surv. Open-File Rep. 2015 1048*, 96. doi:10.3133/ofr20151048
- Hoque, M. A., and Pollard, W. H. (2016). Stability of Permafrost Dominated Coastal Cliffs in the Arctic. *Polar Sci.* 10, 79–88. doi:10.1016/j.polar.2015.10.004
- Jones, B. M., Arp, C. D., Jorgenson, M. T., Hinkel, K. M., Schmutz, J. A., and Flint, P. L. (2009). Increase in the Rate and Uniformity of Coastline Erosion in Arctic Alaska. *Geophys. Res. Lett.* 36, L03503. doi:10.1029/2008gl036205
- Kobayashi, N. (1985). Formation of Thermoerosional Niches into Frozen bluffs Due to Storm Surges on the Beaufort Sea Coast. *J. Geophys. Res.* 90 (C6), 11983–11988. doi:10.1029/jc090ic06p11983
- Mars, J. C., and Houseknecht, D. W. (2007). Quantitative Remote Sensing Study Indicates Doubling of Coastal Erosion Rate in Past 50 Yr along a Segment of the Arctic Coast of Alaska. *Geol* 35 (7), 583–586. doi:10.1130/g23672a.1
- Ping, C.-L., Michaelson, G. J., Guo, L., Jorgenson, M. T., Kanevskiy, M., Shur, Y., et al. (2011). Soil Carbon and Material Fluxes across the Eroding Alaska Beaufort Sea Coastline. *J. Geophys. Res.* 116, G02004. doi:10.1029/2010JG001588
- Ravens, T., Kartezhnikova, M., Ulmgren, M., Yager, G., Jones, B., Erikson, L., et al. (2011). *Arctic Coastal Erosion Modeling*. San Francisco, CA: Presented at the American Geophysical Union Fall Meeting.
- Ravens, T. M., Jones, B. M., Zhang, J., Arp, C. D., and Schmutz, J. A. (2012). Process-based Coastal Erosion Modeling for Drew Point, North Slope, Alaska. *J. Waterway, Port, Coastal, Ocean Eng.* 138 (2), 122–130. doi:10.1061/(asce)ww.1943-5460.0000106
- Ravens, T. M., and Peterson, S. (2018). “Arctic Coastal Erosion Modeling,” in *A Chapter in Advances In Coastal Hydraulics*. Editors P. Vijay and K. Jim (London Singapore: World Scientific). doi:10.1142/9789813231283\_0002
- Ravens, T., and Ulmgren, M. (2020). *Arctic Coastal Geomorphic Change Modeling with Arctic Xbeach*. Anchorage, AK: AGU Fall Meeting.
- U.S. Army Corps of Engineers. (2009). Alaska Baseline Erosion Assessment.65
- Westermann, S., Lüers, J., Langer, M., Piel, K., and Boike, J. (2009). The Annual Surface Energy Budget of a High-Arctic Permafrost Site on Svalbard, Norway. *The Cryosphere* 3, 245–263. doi:10.5194/tc-3-245-2009

## FUNDING

Funding for this research came from the National Science Foundation (Award # 7416374) and from the Bureau of Ocean Energy Management (NSL# AK-17-01).

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher’s Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Ravens and Peterson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.