



The Impact of Ice Caps on the Mechanical Stability of Magmatic Systems: Implications for Forecasting on Human Timescales

Lilian C. Lucas¹, John A. Albright¹, Patricia M. Gregg^{1*} and Yan Zhan^{1,2}

¹Department of Geology, University of Illinois at Urbana-Champaign, Urbana, IL, United States, ²Earth and Planets Laboratory, Carnegie Institution for Science, Washington, DC, United States

OPEN ACCESS

Edited by:

Simona Petrosino,
Istituto Nazionale di Geofisica e
Vulcanologia, Italy

Reviewed by:

Luis E. Lara,
Servicio Nacional de Geología y
Minería de Chile (SERNAGEOMIN),
Chile
Lucía Capra,
National Autonomous University of
Mexico, Mexico

*Correspondence:

Patricia M. Gregg
pgregg@illinois.edu

Specialty section:

This article was submitted to
Volcanology,
a section of the journal
Frontiers in Earth Science

Received: 02 February 2022

Accepted: 13 April 2022

Published: 29 April 2022

Citation:

Lucas LC, Albright JA, Gregg PM and
Zhan Y (2022) The Impact of Ice Caps
on the Mechanical Stability of
Magmatic Systems: Implications for
Forecasting on Human Timescales.
Front. Earth Sci. 10:868569.
doi: 10.3389/feart.2022.868569

Monitoring the activity of subglacial volcanoes along the Aleutian Arc in Alaska is important to the safety of local populations, as well as air traffic flying through the region. However, observations of volcanic unrest are limited by accessibility and resources, particularly at glacier-covered systems, making investigations of their stability challenging. Westdahl Peak, a subglacial volcano on Unimak Island in the Aleutian Arc has experienced significant unrest and uplift since its most recent VEI three eruption in 1991-1992. Given the magnitude of observed uplift, previous investigations suggested the potential for eruption by 2010, but no such event has occurred. One hypothesis to explain this prolonged unrest is that the 1-km thick glacier may increase the stability of the magma system. However, the impact of ice caps and glaciers on the short-term stability of volcanoes is not well understood. In this study, thermomechanical finite element models are used to evaluate how the stability of a glaciated volcano is impacted by variations in ice cap thickness, magma chamber depth, geometry, magma flux rate, and seasonal changes in ice cover thickness. Our numerical experiments indicate that the presence of an ice cap (1–3 km thick) increases the average repose interval for a magma system. Among models with different magma chamber geometries, depths, and flux rates, the greatest increases in repose interval are observed in prolate systems where the increase is up to 57% for a chamber located at 5 km-depth. Spherical and oblate also experience smaller, yet significant, increases in repose interval. Additionally, the percentage increase in repose interval is not impacted by variations in magma flux rate for a given ice cap thickness and magma chamber geometry. However, flux rates do influence the timing of eruptions when the system is experiencing seasonal variations in ice thickness. Our results show that systems with low flux rates are more likely to fail when the ice thickness is at its lowest. The numerical estimates further suggest that the ice cap on Westdahl Peak, which is ~1 km, may slightly increase the stability of the magma system. In general, given flux rates and magma chamber geometries estimated for the Westdahl system, the repose interval can increase by ~7 years due to the Westdahl glacier. This increase is small on a geologic scale but is significant on human time scales and the impact of glaciers must be considered in future forecasting efforts.

Keywords: volcano, subglacial, flux rate, westdahl, seasonality, viscoelastic, finite element method

INTRODUCTION

Constraining the impact of glaciation on magma system stability is critical for evaluating potential hazards and the eruption potential of active volcanoes located in high latitudes. Previous studies have investigated the effects of deglaciation on magma systems through elastic axisymmetric models, and conclude that a decrease in surface load pressure may either enhance or inhibit conditions for dike initiation dependent upon magma chamber geometry and depth (Albino et al., 2010; Sigmundsson et al., 2010). In particular, these studies evaluate how decreasing surface loads impact both the pressure within the magma chamber and the threshold conditions for tensile failure and dike initiation in the host rock. Albino et al. (2010) apply their models to Grímsvötn and Katla Volcanoes in Iceland. Both applications conclude that surface load variations, such as jökulhlaups (large lake water discharge events) and annual snow cover melting, may enhance the potential for dike initiation and/or eruption when the magma chamber is already near failure. Additionally, recent numerical studies have shown that viscoelastic effects in the host rock impact the timing and conditions for failure of long-lived magma systems (Del Negro et al., 2009; Gregg et al., 2012; Cabaniss et al., 2018; Zhan and Gregg, 2019). However, the combined effects of both viscosity and ice cap loading have not yet been studied. It is therefore critical to investigate the roles that viscoelasticity and long-term stress accumulation play in glaciated systems.

In this study, we run a series of numerical simulations to investigate the impact of ice caps on the stability and eruption potential of long-lived magma systems. Viscoelastic, temperature-dependent models are developed to evaluate the overall effect of ice caps on magma systems. Particular care is taken to assess the effect of variables such as ice cap thickness, magma chamber size, geometry, and the flux of new material into the magma system. Finally, we apply our generic model to Westdahl Volcano, a caldera system on Unimak Island in the Aleutian Arc of Alaska (Figure 1). Westdahl is covered by ~1 km

of ice and has an active magma chamber at a depth of ~7.2 km (Gong, 2015). Since its last eruption, in 1991-1992, Westdahl has experienced significant and ongoing inflation that previous studies interpreted as an indication of imminent eruption (Lu et al., 2004). However, no eruption has occurred despite continuing inflation well past the time scale considered in these studies. One potential explanation for this lack of eruption is that the mechanical loading of the ice cap is prolonging the magma reservoir's stability. In applying our generic model, we investigate the contribution of Westdahl's ice cap to the mechanical stress state of the volcano and the ongoing likelihood of eruption.


METHODS

Building upon previous models (Grosfils, 2007; Gregg et al., 2012; Albright et al., 2019), we use COMSOL Multiphysics 5.5 to create viscoelastic finite element models that incorporate temperature-dependent rheology (Figure 2). In these models, ice caps of various thicknesses are then implemented to simulate glaciated volcanoes. In a 50 by 25 km, two-dimensional model space, the magma reservoir is constructed as a pressurized elliptical void along a central axis of rotational symmetry. Roller boundaries are implemented on the right lateral and bottom walls. A layer with variable thickness is included to simulate an ice cap covering the volcano. The rock and ice domains are gravitationally loaded to impose an initial lithostatic stress.

In this study, we use a generalized Maxwell model to describe a viscoelastic rheology whose relaxation time, τ , is given by:

$$\tau = \left(\frac{\eta}{G_{0td}u_1} \right) * \left(\frac{3K_{td} + G_{0td}}{3K_{td} + G_{0td}u_0} \right) \quad (1)$$

In Eq (1), the shear modulus, G_{0td} , is the instantaneous elastic response of the material when under pressure. The time dependent viscous response is represented by η (Gregg et al.,






TABLE 1 | Model parameters.

Parameter	Description	Value
a, b	Magma chamber radius, m	1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000
C_p	Heat Capacity, J/(g*°C)	1×10^{-5}
ΔV	Flux rate, m ³ /yr	$1 \times 10^5, 5 \times 10^5, 1 \times 10^6, 2.4 \times 10^6, 5 \times 10^6, 6.7 \times 10^6, 1 \times 10^7, 5 \times 10^7$
d	Depth to center of magma chamber, m	2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000
G_0	Shear Modulus of rock, Pa	2×10^{10}
G_i	Shear Modulus of ice, Pa	2×10^{10}
g	Gravity, m/s ²	9.8
η_i	Viscosity of ice, Pa*s	2×10^{13}
η_r	Viscosity of rock, Pa*s	2×10^{16}
ih	Ice thickness, m	0, 1000, 2000, 3000
K	Bulk Modulus, Pa	3.333×10^{10}
k	Thermal Conductivity, W/(m*K)	3
μ_1	Fractional Shear	0.5
N_i	Bulk Modulus, Pa	9.8029×10^8
ν	Poisson Ratio	0.25
ρ_i	Density of ice, g/m ³	916000
ρ_r	Density of rock, g/m ³	2700000
T_{grad}	Temperature Gradient, °C/m	0.03
T_m	Magma chamber temperature °C	100

2012). K_{td} is the bulk modulus, and u_0 and u_1 are the fractional moduli (see **Table 1**).


The COMSOL solid mechanics and heat transfer modules are used to implement both time- and temperature-dependence within the viscoelastic model space. **Eq (2)** represents the temperature-dependent host rock viscosity:

$$\eta = A_D e^{\frac{E_a}{R_{gas} T}} \quad (2)$$

where A_D is the Dorn parameter, E_a is activation energy, R_{gas} is the gas constant, and T is temperature. Our model also

incorporates a geothermal gradient of 30°C/km as well as a constant magma chamber temperature of 1,000°C, which is appropriate for a basaltic magma system.

Throughout the modeling process, we measure the change in repose interval of the magma systems in response to changing parameters including different thicknesses of ice, magma chamber geometries, magma chamber depths, and magma flux rates. Repose interval is defined as the time period for the tensile stress along the reservoir wall to progress from a state of compression to tension. In this investigation, we focus on



tensile failure along the reservoir boundary as a proxy for dike initiation and eruption, as opposed to shear failure in the surrounding host rock, which has also been shown to potentially catalyze eruptions (Gregg et al., 2012). Tensile rupture of the chamber and the onset of eruption is assumed to occur when tensile stress becomes positive:

$$\sigma_t > 0 \quad (3)$$

FACTORS CONTROLLING THE HOST ROCK STABILITY IN ICE-COVERED VOLCANOES

A series of numerical experiments are conducted to investigate the impact of ice cap loading on the stability of magma systems. In particular, we examine the relationship between the stability of a magma system replenished by a constant magma flux and the presence an ice cap. The repose interval is defined as the time from the start of new magma injection, which remains constant throughout the simulation, to when tensile failure is observed at the reservoir boundary, as described above. Specifically, repose intervals of magma systems with ice caps of 1–3 km are compared with repose intervals of otherwise identical magma systems without ice to calculate percent change in time to eruption due to the ice cap. We should note that the repose interval described here represents a minimum, and natural systems may have protracted periods with no new magma injection. Results, including both the absolute timescales as well as the percent



change in repose for all of the experiments described below can be found in the **Supplementary Material S1**.

Ice Cap Thickness

In general, an additional load above a magma chamber, such as from ice, increases the confining stress along the reservoir wall, making it more difficult for the chamber to rupture (Grosfils, 2007). In all experiments, we found that ice cap thickness increases the repose interval, but different parameters of the magma system affect how much the repose interval is changed relative to an ice-free baseline. In the following sections, we investigate the effects of magma flux (3.2), depth to reservoir crest (3.3), and reservoir geometry (3.4).

Magma Flux

Magma flux is thought to be a primary driver of inflation and ultimately controls the timing of failure for magma systems (Cabaniss et al., 2018). As such, a wide range of fluxes is investigated from 0.0001 to 0.05 km^3/yr (Wilson et al., 2006; Annen, 2009; Gelman et al., 2013). Model results indicate that the percent increase of repose interval due to glacial loading is independent of the flux rate (**Figure 3**). For instance, the repose interval of a spherical chamber at 5 km depth will increase by $\sim 4.8\%$, independent of varying the flux rate. Similarly, a prolate chamber located at a depth to center of 5 km underneath 3 km of ice, will see an increase in repose of $\sim 23\%$, regardless of the flux rate (**Figure 3B**). A similar response is observed for chambers with an oblate geometry (**Figure 3C**). Additionally, the timing of tensile failure in systems with identical



depth, shape, and ice cap thickness occurs independent of flux rate for similar volumes of magma added. Although the modeled magma systems indicate an insensitivity to flux rate, their repose intervals are highly dependent on the total magnitude of volume

change (Figure 4). For each simulation, tensile failure is produced for the same total volume of magma added, regardless of the rate of intrusion, for the flux rates investigated (0.0001–0.05 km³/yr). For example, tensile failure for a spherical magma chamber with a




FIGURE 6 | Magma chamber radius versus percent change in repose interval of volcanic systems with spherical, prolate, and oblate magma chamber geometries. In each case, a volcanic system with a depth of 5 km to the crest of the magma chamber and a $0.0024 \text{ km}^3/\text{year}$ flux rate is modeled. The magma chamber radii change for each of the spherical, prolate and oblate chamber systems. **(A)** In the spherical system, the horizontal and vertical radii are identical lengths and range from 0.5 to 4 km. The percent increase of repose interval from the presence of the glacier slightly decreases when the radius is increased from 0.5 to 2 km length and increases when the radius is increased from 2 to 4 km. **(B)** In the prolate system, the horizontal radius remains ~ 0.49 km while the vertical radius ranges between 0.5 and 4 km. **(C)** The vertical radius for an oblate magma chamber system is held constant at ~ 0.39 km while the horizontal radius ranges between 0.5 and 4 km. Repose interval percent change increases as the horizontal radius of an oblate magma chamber is increased in length.

depth of 7 and 1 km of ice will occur when $\sim 0.24 \text{ km}^3$ of magma is added to the chamber (Figure 4A).

Magma Chamber Depth

Previous studies have interpreted magma chamber depth to be an influential factor on the stability of a magma system (Albino et al., 2010; Geyer and Bindeman, 2011; Satow et al., 2021). A suite of simulations is run to examine the impact of magma chamber depths to center, ranging from 2–9 km (e.g., Huber et al., 2019; Rasmussen et al., 2019; Kent et al., 2022). Results indicate that the ice cap has a decreased effect on the relative repose interval for deeper magma sources, as is expected given the overall contribution to the surface load. For instance, the repose interval of a system with a spherical chamber at a depth of 2 km will increase by $\sim 55\%$ when loaded with a 3 km ice cap (Figure 5A). Identical loading conditions will increase the repose interval by only $\sim 11\%$ for a system with a magma chamber at 9 km depth to center (Figure 5A). This effect is greater for non-spherical chambers, with prolate chambers being most sensitive to variations in magma chamber depth (Figure 5B). Specifically, a prolate chamber located at 2 km depth beneath 3 km of ice will experience an $\sim 86\%$ increase in its repose interval. Whereas the same chamber at 9 km depth will have an increase in repose of $\sim 12\%$ (Figure 5B).

Magma Chamber Geometry

The final parameter we investigate in this study is magma chamber geometry. In accordance with previous studies, we

model three simplified magma chamber shapes: spherical, prolate, and oblate (Lisowski, 2007). Starting off, as the radius of a spherical chamber increases, we find glacial loading will ultimately have an increased effect on the repose interval. Our models show that the presence of a 3 km glacier increases the relative repose interval from 17.6% for a chamber with a 0.5 km radius to 34.6% for a chamber with a 4 km radius (Figure 6A). Oblate systems experience a similar, but amplified change in relative repose interval as the size of the horizontal radius is increased (Figure 6C). For example, a 3 km-thick ice cap will increase the repose interval from 22.3% for a 0.5 km horizontal radius to 103% for a 4 km horizontal radius (Figure 6C). Finally, as a prolate chamber becomes more prolate, there is not a significant change in the repose interval given a specific ice cap thickness.

DISCUSSION

The Role of Glaciers in Maintaining Magma System Stability

Previous investigations have focused on the twofold impact of active regional ice cap withdrawal on both the flux of magma into the magma chamber (Sigmundsson et al., 2010) and the threshold conditions for driving tensile failure and dike initiation (Albino et al., 2010). In particular, the removal of ice caps may promote additional melt generation from the mantle source due to decompression, which may increase flux to the crustal magma

system (Sigmundsson et al., 2010). Furthermore, the decreased load from the ice cap withdrawal will reduce confining pressure, promoting tensile failure and the likelihood of dike initiation if a magma system is near failure (Albino et al., 2010). In contrast, this investigation focuses purely on the static, long-term effects of the presence or absence of ice caps and does not consider the dynamic processes of their active removal. Motivated by high latitude, glacially covered systems such as Westdahl Volcano in Alaska, we seek to determine whether the presence of a large glacier or ice cap can account for unexpectedly long repose intervals (Lu et al., 2004).

Our numerical experiments indicate that glaciers modulate eruption timescales. In particular, the additional load from an ice cap increases the confining stress around the magma reservoir, which must subsequently be overcome to initiate tensile failure. For a given reservoir size and shape, this additional stress is dependent only on the thickness of the glacier, requiring a fixed volume change within the reservoir, and by extension time, to reach failure. In most cases, the delay due to ice loading is relatively small compared to the system's baseline repose interval, which is predominantly controlled by the lithostatic stress. However, for shallow reservoirs (<5 km depth to center), an ice cap contributes a greater proportion of the overlying load, and as such has a larger relative impact on the potential repose interval. Regardless, even for deep systems where the relative change in repose interval due to the presence of ice is small, the impact is significant on human timescales and must be considered in future forecasting efforts. For example, a spherical chamber (radius of 0.62 km and flux of 0.0024 km³/yr) at 9 km depth will experience an increase in repose of ~3 months~15 years with a 1 km or 3 km-thick ice cap respectively.

The Role of Flux

In many previous numerical modeling studies that use elastic rheology, the main driver of reservoir tensile failure has been the cumulative critical volume change or a critical change in overpressure of the magma reservoir (Grosfils, 2007; Gerbault et al., 2012; Gregg et al., 2012). This effect is somewhat modulated in viscoelastic rheology where the host rock can relax to dissipate some of the reservoir boundary stresses (Gregg et al., 2012; Gregg et al., 2013; Degruyter and Huber, 2014). In our results, however, the viscoelasticity does not appear to play a major role in system stability. If significant viscoelastic relaxation were present, we would expect to see a greater relative change in repose interval at lower fluxes. Specifically, in the time it takes to overcome the additional load of the ice cap, the system would have relaxed even further, requiring yet more volume change. In contrast, we observe a near constant relative change in repose interval, regardless of flux rate (Figure 3).

Our numerical results agree with previous studies, which indicate that volcanic unrest on year to decadal timescales mostly behaves in an elastic manner (Zhan and Gregg, 2019). With higher flux rates (>0.05 km³/yr), repose timescales will be even shorter, bringing the system even closer to the elastic endmember. Alternatively, at lower flux rates (<0.0001 km³/yr), the likelihood of solidification prior to eruption increases

(Annen, 2009). Additionally, on longer timescales there is more time for viscoelastic relaxation to take effect, prolonging stable storage and inhibiting eruption even further.

The existence of an ice cap linearly increases the confining pressure around a magma reservoir, which must then be overcome by changing the volume of magma by a set amount, ΔV_q . The time needed to accumulate this critical volume of magma, in other words the total delay before an eruption (t_d), will therefore vary inversely with the flux rate (q) into the system:

$$t_d = \Delta V_q / q$$

For example, the total volume of magma required for the tensile failure of a volcanic system with a spherical magma chamber at 3 km depth will be ~0.1 km³ of magma, independent of flux rate (Figure 4).

Seasonal Impacts on the Timing of Failure

In understanding the potential effects of loading on the stability of a magma system, previous studies have additionally investigated the effects of a seasonal surface load change. Annual unloading events, such as snow melt represented in numerical models, have suggested that a decrease in a load as thin as 6.5 m may trigger a dike initiation event, if the system is initially near failure conditions (Albino et al., 2010). This effect has been suggested in systems such as Katla Volcano, a subglacial system with annual snow cover, whose last nine major eruptions have been initiated between May and November when the yearly snow had melted (Larsen, 2000).

The results of our study suggest the presence of an annual snow cover increases the repose interval of the volcano, but the strength of this effect is strongly dependent on the thickness of the ice cover and parameters of the magma system. In particular, annual changes in snow and ice thickness at Katla and Westdahl volcanoes are on the order of 6.5 m (Albino et al., 2010; Littell et al., 2018), significantly less than the km-scale ice masses modeled here. Compared to the total overlying load that the magma chamber must overcome in order to reach tensile failure, small seasonal variations are unlikely to play a major role for most systems. However, seasonal ice loss may impact the timing of eruption for systems close to failure.

Using our modeling results above as well as additional simulations with 10 m of ice, we compare the impact of short-term ice thickness variations on the seasonal timing of magma system failure (Figure 7). Four end-member models are evaluated to determine the relative impacts of flux-rate and seasonal ice variability on when magma system failure is more likely to occur. In each case, the reservoir failure threshold is interpolated as a sinusoid that alternates yearly between the ice-free and the 1 km ice cap results. For lower seasonality, we assume a linear relationship between ice thickness and the reservoir failure threshold and proportionally reduce the amplitude of the sinusoid.

Our results indicate that low flux rate systems are much more likely to fail when the ice thickness is at its lowest, August - October in the northern hemisphere (Figure 7). In these systems ice mass variations impact the failure conditions more than




FIGURE 7 | The relationship between flux rate and seasonality on the timing of reservoir tensile failure. Each subplot compares the seasonally varying tensile failure threshold of a 3 km deep (depth to center, red line) and 5 km deep (depth to center, green line) spherical reservoir (radius = 0.62 km) against the continuous volume flux into the magma system (black line). The subplots are arranged within a set of larger axes, determining the amount of seasonal ice variation (10 m vs. 1 km) and the flux rate (0.0001 km³/yr vs. 0.05 km³/yr) into the system. In each case, tensile failure occurs at the point where the volume in the magma reservoir first intersects the failure threshold (black stars). The solid, dashed, and dotted black lines indicate an example trajectory of the volume change before the initial rupture of the 3 km deep chamber, before the rupture of the 5 km chamber, and after the rupture of the 5 km chamber, respectively. When flux is low (left subplots), seasonal ice variations outpace the system’s long-term accumulation of magma volume, consistently causing failure when ice thickness reaches its minimum. On the other hand, high-flux systems (right subplots) can produce tensile failure in any season, although the distribution may be skewed depending on the magnitude of ice variation.

increasing volume change in the magma reservoir. In contrast, for magma systems with high flux and low seasonal ice variability, the onset of failure is most sensitive to the accumulation of magma, rather than the timing of the lowest level of ice mass, and so can occur at any time of year. Finally, systems with both high magma flux and high-seasonal ice variability display a slight skew in the specific timing of reservoir failure; eruptions are possible at any time, but are more likely to occur during periods of low ice mass.

We should note, however, that our results are entirely based on the stresses within the host rock and do not take into account the pressure changes within the magma reservoirs during ice loss that may also affect the onset of eruptions, such as decompression melting in the magma reservoir (Mora and Tassara, 2019) or hydrologic effects from the influx of meltwater (Albino et al., 2010). Such effects would likely push eruptions to earlier in the warm season.

Is Westdahl’s Protracted Unrest due to Glacial Loading?

Previous investigations of the unrest of Westdahl have observed a prolonged period of surface inflation, but without an accompanying eruption (Lu and Dzurisin, 2014). To evaluate the effect Westdahl’s 1 km-thick ice cap has on this long-term stability, we apply our aforementioned numerical modeling




FIGURE 8 | Westdahl tensile stress versus time with and without a 1 km glacier. The figure has been zoomed in to highlight the timing of failure. Failure condition is represented by the black horizontal line and repose intervals are represented by the dotted lines. Depth to reservoir center is 7.2 km, flux rate is 0.0024 km³/yr, and the radius of the magma chamber is 1.39 km (based on geodetic inversions conducted by Gong, 2015).

approach to replicate conditions at this specific volcano. Based on previous geodetic inversions (Lu and Dzurisin, 2014; Gong, 2015), we simulate a 1.39 km-radius spherical reservoir centered 7.2 km below the ground surface with a flux of 0.0024 km³/yr, and then compare the system’s repose interval with and without an ice cover of 1 km (Figure 8).

Based on our numerical results, we find that the presence of the glacier would at most delay the onset of eruption by ~7 years as compared to models of Westdahl with no glacier (99 vs. 92 years from the start of constant inflation). This is significant on monitoring time scales, but relatively small compared to the modeled system's repose interval of ~100 years. However, the true system's absolute repose times will vary as the system geometry and underlying flux rates evolve. Based on our more general results, Westdahl's long-term stability is most likely controlled by the system's relatively deep reservoir (Figure 5) and spherical geometry (Figure 6). Regardless, future models attempting to forecast Westdahl's unrest need to consider all parameters that may significantly affect the onset of eruption on human time scales, including loading from the ice cap.

Westdahl's 5–10 m seasonal ice variations (Littell et al., 2018) and recent inflation, which has an estimated flux of ~0.0024 km³/yr (Lu and Dzurisin, 2014), place it as a moderately high flux rate system with low seasonal variability (Figure 7). As such, we would expect the timing of eruption from Westdahl to be only moderately sensitive to seasonal ice loss. However, we do not know how close to failure the system is currently, and recent geodetic data have not yet been investigated to update our estimates of the system's flux rate. If the flux into Westdahl's magma reservoir has slowed in recent years while the system remains close to failure, seasonality may ultimately play a larger role than would be expected from this study. Future investigations would benefit from comparing more up-to-date estimates of Westdahl's unrest and its seasonal ice variations, as demonstrated here.

CONCLUSION

In this study, we investigate how the presence or absence of ice caps impacts the stability of volcanic systems on human timescales. Our numerical results indicate that large ice caps and glaciers with thicknesses of 1–3 km can appreciably delay the onset of eruptions on the order of years to decades. In particular, there is a linear relationship between the thickness of an ice cap and the additional reservoir volume change necessary to induce tensile failure. As such, the specific change in repose interval will depend on the underlying flux rate into the system for a given ice thickness. Therefore, when flux rates are low, seasonal variations in the overlying ice cap thickness may be adequate to trigger tensile failure once a system approaches the critical stress threshold. When we apply our findings to the recent unrest at Westdahl Volcano, Alaska, we find that for the previously modeled flux rate of 0.0024 km³/year, the ~1 km ice cap would delay an eruption by ~7 years. Although the ice cap may greatly

increase stability of the volcano, it is unlikely to fully account for the volcano's long periods of non-eruptive inflation. However, the numerical results presented assume constant flux, which may not capture the full dynamics of systems with long periods of quiescence or sudden spikes in flux rate. Moreover, we focus only on the stress conditions related to ice caps and not additional dynamic effects that may occur during their active emplacement or removal. This study establishes basic relationships between ice caps and volcanic reservoir stability, but future works should incorporate more realistic or dynamic parameters.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding author.

AUTHOR CONTRIBUTIONS

LL led this work as part of her undergraduate research supervised primarily by JA with guidance from PG and YZ. All authors contributed to analyzing the numerical results, drafting figures, and writing the manuscript.

FUNDING

Numerical modeling to investigate magma system stability is funded U.S. National Science Foundation (NSF GRFP-Albright, EAR 2122745 and EAR 1752477-Gregg), NASA (80-NSSC19K-0357-Gregg), and the University of Illinois Department of Geology Summer Undergraduate Research Opportunity Program (Lucas).

ACKNOWLEDGMENTS

The authors would like to thank C. Proistosescu, H. Cabaniss, R. Goldman, and M. Head for helpful conversations that enhanced this investigation.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/feart.2022.868569/full#supplementary-material>

REFERENCES

Albino, F., Pinel, V., and Sigmundsson, F. (2010). Influence of Surface Load Variations on Eruption Likelihood: Application to Two Icelandic Subglacial Volcanoes, Grímsvötn and Katla. *Grímsvötn*, 1510–1524. doi:10.1111/j.1365-246X.2010.04603.x

Albright, J. A., Gregg, P. M., Lu, Z., and Freymueller, J. T. (2019). Hindcasting Magma Reservoir Stability Preceding the 2008 Eruption of Okmok, Alaska. *Geophys. Res. Lett.* 46, 8801–8808. doi:10.1029/2019GL083395

Annen, C. (2009). From Plutons to Magma Chambers: Thermal Constraints on the Accumulation of Eruptible Silicic Magma in the Upper Crust. *Earth Planet. Sci. Lett.* 284, 409–416. doi:10.1016/j.epsl.2009.05.006

- Cabaniss, H. E., Gregg, P. M., and Grosfils, E. B. (2018). The Role of Tectonic Stress in Triggering Large Silicic Caldera Eruptions. *Geophys. Res. Lett.* 45, 3889–3895. doi:10.1029/2018GL077393
- Degruyter, W., and Huber, C. (2014). A Model for Eruption Frequency of Upper Crustal Silicic Magma Chambers. *Earth Planet. Sci. Lett.* 403, 117–130. doi:10.1016/j.epsl.2014.06.047
- Del Negro, C., Currenti, G., and Scandura, D. (2009). Temperature-dependent Viscoelastic Modeling of Ground Deformation: Application to Etna Volcano during the 1993–1997 Inflation Period. *Phys. Earth Planet. Interiors* 172, 299–309. doi:10.1016/j.pepi.2008.10.019
- Gelman, S. E., Gutiérrez, F. J., and Bachmann, O. (2013). On the Longevity of Large Upper Crustal Silicic Magma Reservoirs. *Geology* 41, 759–762. doi:10.1130/G34241.1
- Gerbault, M., Cappa, F., and Hassani, R. (2012). Elasto-plastic and Hydromechanical Models of Failure Around an Infinitely Long Magma Chamber. *Geochem. Geophys. Geosyst.* 13, a–n. doi:10.1029/2011GC003917
- Geyer, A., and Bindeman, I. (2011). Glacial Influence on Caldera-Forming Eruptions. *J. Volcanol. Geotherm. Res.* 202, 127–142. doi:10.1016/j.jvolgeores.2011.02.001
- Gong, W. (2015). Journal of Geophysical Research : Solid Earth. *AGU J. Geophys. Res. Solid Earth* 119, 3678–3699. doi:10.1002/2013JB010641. Received
- Gregg, P. M., De Silva, S. L., Grosfils, E. B., and Parmigiani, J. P. (2012). Catastrophic Caldera-Forming Eruptions: Thermomechanics and Implications for Eruption Triggering and Maximum Caldera Dimensions on Earth. *J. Volcanol. Geotherm. Res.* 241–242, 1–12. doi:10.1016/j.jvolgeores.2012.06.009
- Gregg, P. M., de Silva, S. L., and Grosfils, E. B. (2013). Thermomechanics of Shallow Magma Chamber Pressurization: Implications for the Assessment of Ground Deformation Data at Active Volcanoes. *Earth Planet. Sci. Lett.* 384, 100–108. doi:10.1016/j.epsl.2013.09.040
- Grosfils, E. B. (2007). Magma Reservoir Failure on the Terrestrial Planets: Assessing the Importance of Gravitational Loading in Simple Elastic Models. *J. Volcanol. Geotherm. Res.* 166, 47–75. doi:10.1016/j.jvolgeores.2007.06.007
- Huber, C., Townsend, M., Degruyter, W., and Bachmann, O. (2019). Optimal Depth of Subvolcanic Magma Chamber Growth Controlled by Volatiles and Crust Rheology. *Nat. Geosci.* 12, 762–768. doi:10.1038/s41561-019-0415-6
- Kent, A., Wieser, P. E., and Till, C. (2022). *Geophysical and Geochemical Constraints on Magma Storage Depths along the Cascade Arc: Knowns and Unknowns*. Goldschmidt.
- Larsen, G. (2000). Holocene Eruptions within the Katla Volcanic System, South Iceland: Characteristics and Environmental Impact. *Jökull* 49, 1–28.
- Littell, J., McAfee, S., and Hayward, G. (2018). Alaska Snowpack Response to Climate Change: Statewide Snowfall Equivalent and Snowpack Water Scenarios. *Water* 10, 668. doi:10.3390/w10050668
- Lu, Z., and Dzurisin, D. (2014). *InSAR Imaging of Aleutian Volcanoes: Monitoring a Volcanic Arc from Space*.
- Lu, Z., Rykhus, R., Masterlark, T., and Dean, K. G. (2004). Mapping Recent Lava Flows at Westdahl Volcano, Alaska, Using Radar and Optical Satellite Imagery. *Remote Sens. Environ.* 91, 345–353. doi:10.1016/j.rse.2004.03.015
- Lisowski M., (Editor) (2007). *Analytical Volcano Deformation Source Models* (Bath, UK: Springer), 469. doi:10.1007/978-3-540-49302-0
- Mora, D., and Tassara, A. (2019). Upper Crustal Decompression Due to Deglaciation-Induced Flexural Unbending and its Role on Post-glacial Volcanism at the Southern Andes. *Geophys. J. Int.* 216, 1549–1559. doi:10.1093/gji/ggy473
- Rasmussen, D. J., Plank, T. A., and Roman, D. C. (2019). *The Aleutian Arc through and through: Magmatic Water Content Controls Magma Storage Depth*. New York, NY: Columbia University.
- Satow, C., Gudmundsson, A., Gertisser, R., Ramsey, C. B., Bazargan, M., Pyle, D. M., et al. (2021). Eruptive Activity of the Santorini Volcano Controlled by Sea-Level Rise and Fall. *Nat. Geosci.* 14, 586–592. doi:10.1038/s41561-021-00783-4
- Sigmundsson, F., Pinel, V., Lund, B., Albino, F., Pagli, C., Geirsson, H., et al. (2010). Climate Effects on Volcanism: Influence on Magmatic Systems of Loading and Unloading from Ice Mass Variations, with Examples from Iceland. *Phil. Trans. R. Soc. A* 368, 2519–2534. doi:10.1098/rsta.2010.0042
- Wilson, C. J. N., Blake, S., Charlier, B. L. A., and Sutton, A. N. (2006). The 26.5 Ka Oruanui Eruption, Taupo Volcano, New Zealand: Development, Characteristics and Evacuation of a Large Rhyolitic Magma Body. *J. Pet.* 47, 35–69. doi:10.1093/petrology/egi066
- Zhan, Y., and Gregg, P. M. (2019). How Accurately Can We Model Magma Reservoir Failure with Uncertainties in Host Rock Rheology? *J. Geophys. Res. Solid Earth* 124, 8030–8042. doi:10.1029/2019JB018178

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Lucas, Albright, Gregg and Zhan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.