



Association Between Different Indicators of Obesity and Depression in Adults in Qingdao, China: A Cross-Sectional Study

Jing Cui^{1,2†}, Xiufen Sun^{3†}, Xiaojing Li^{1,2†}, Ma Ke⁴, Jianping Sun^{1,2}, Nafeesa Yasmeen⁵, Jamal Muhammad Khan⁶, Hualei Xin^{1,2}, Shouyong Xue^{7*} and Zulqarnain Baloch^{8*}

¹ Qingdao Municipal Center for Disease Control and Prevention, Qingdao, China, ² Qingdao Institute of Preventive Medicine, Qingdao, China, ³ Qingdao Shi'nan Municipal Center for Disease Control and Prevention, Qingdao, China, ⁴ College of Traditional Chinese Medicine, Shandong University of Traditional Chinese Medicine, Jinan, China, ⁵ Institute of Microbiology, Agriculture University Faisalabad Pakistan, Bahawalpur, Pakistan, ⁶ Department of Patho-biology, The Islamia University of Bahawalpur, University College of Veterinary and Animal Sciences, Bahawalpur, Pakistan, ⁷ Qingdao Shi'bei Municipal Center for Disease Control and Prevention, Qingdao, China, ⁸ College of Veterinary Medicine, South China Agricultural University, Guangzhou, China

OPEN ACCESS

Edited by:

Andrew J. McAinch,
Victoria University, Australia

Reviewed by:

Scott B. Teasdale,
University of New South Wales,
Australia
Guoqin Xu,
Guangzhou Sport University, China

*Correspondence:

Shouyong Xue
sbjxxue@163.com
Zulqarnain Baloch
znbalooch@yahoo.com

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Obesity,
a section of the journal
Frontiers in Endocrinology

Received: 26 May 2018

Accepted: 29 August 2018

Published: 10 October 2018

Citation:

Cui J, Sun X, Li X, Ke M, Sun J, Yasmeen N, Khan JM, Xin H, Xue S and Baloch Z (2018) Association Between Different Indicators of Obesity and Depression in Adults in Qingdao, China: A Cross-Sectional Study. *Front. Endocrinol.* 9:549. doi: 10.3389/fendo.2018.00549

Background: This study was designed to investigate the perceived relationship between body weight and depression risk in a Chinese population in Qingdao, China.

Methods: A population-based cross-sectional survey was performed with 4,573 participants (between 35 and 74 years) from the year 2009 to 2012 in Qingdao, China. We applied the Zung self-rating depression scale to ascertain the level of depression in participants. The associations between different indicators of obesity [body mass index (BMI), waist circumference (WC), and waist-to-hip ratio (WHR)] and depression were assessed by logistic regression based on the Chinese criteria of obesity. Sensitivity analysis was done based on the Asian and WHO criteria of obesity.

Results: The Zung scores for the 243 participants (5.2%) were over 45 and they were entitled as depression. Furthermore, multivariable logistic analyses revealed that being overweight [odds ratios (OR): 1.48, 95% confidence intervals [95% CI]: 1.08–2.03] and having abdominal obesity (WC category in Chinese criteria) (OR: 1.47, 95% CI: 1.08–2.00) were often associated with a higher risk for depression compared to normal weight subjects. Sensitivity analysis revealed that abdominal obesity (Asian criterion) (OR: 1.41, 95% CI: 1.03–1.91) was a significant risk factor for depression. Similarly, being overweight (WHO criterion) (OR: 1.39, 95% CI: 1.03–1.87) was an obvious risk factor for depression.

Conclusion: Being overweight and having abdominal obesity (WC category) were found to be linked with a higher risk of depression. However, abdominal obesity (WHR category) was not associated with depression.

Keywords: depression, body mass index, waist circumference, waist-to-hip ratio, cross-sectional study

Abbreviations: BMI, Body mass index; CI, Confidence intervals; OR, Odds ratio; WC, waist circumference; WHO, World Health Organisation.

INTRODUCTION

Significant progress has indeed been made in improving public health in the world; however, many serious health issues still need to be resolved such as depression and obesity. The global prevalence of being overweight and obese was estimated to be 39% and 13% in adults aged 18 years and over in 2017, respectively (1). In 2018, the World Health Organization (WHO) estimates that depression is seen to affect approximately 300 million people worldwide (2). In China, the prevalence of obesity has been continuously increasing during the last few decades (3–5). Additionally, depression is also frequently diagnosed at public and private clinics. According to health officials, approximately 26 million people are struggling with symptoms of mild depression annually in China (6).

Several population-based studies conducted in Western countries demonstrated that the risk for depression was positively associated with being overweight and obesity (7–12). In another study, a U-shaped association between BMI and depression has been reported (10). However, estimates obtained from the Chinese population are found to be controversial (13–15). Several research activities have reported a negative correlation between the risk for depression and obesity among middle-aged and the elderly Chinese (13, 14). In contrast, Li et al. observed a significant positive association between obesity and increased depression scores in the Chinese elderly (aged ≥ 65 years) (15). Nonetheless, mechanisms underlying the association between obesity and depression are still unclear. Some studies have shown that the brain–reward regions for depression and obesity, such as the dopaminergic pathways, can enhance positive mood. Yet, they also cause increased intake of “comfort” food, which may lead to obesity (16, 17). As a matter of fact, the perceptions regarding obesity are different between the Western world and the Chinese. While obesity is a stigma in Western countries (18), it is considered as a symbol of wealth in China (13–15). Regardless of these differences, it may be due to population diversity, different body weight criteria, and depression standards in China.

The association between obesity and depression has been normally studied in adolescents and the elderly population in China (19–21); yet there is a need to explore various indicators of obesity in detail. Hence, this large population-based cross-sectional survey was carried out to assess the potential relationship between depression and different body weight levels among the general population of Qingdao, China.

METHODS

Ethical Considerations

All the participants voluntarily signed the consent prior to their participation. The security, anonymity, and the privacy of participants were strictly respected. This study was approved by the Ethics Committee of the Qingdao Municipal Center for Disease Control and Prevention.

Study Population

This cross-sectional population-based survey was conducted in the eastern city of Qingdao, Mainland China during 2009 to 2012.

Inclusion and exclusion criteria have already been described in our previous study (22). Briefly, a total of 6,100 Chinese adults (between 35–74 years old) were selected with a stratified, random cluster sampling procedure and invited to participate in the survey. A total of 5,110 individuals agreed to take part in our survey, with a response rate of 83.8%. From the 5,110 individuals, 454 participants were excluded due to an insufficient Zung score or body mass index (BMI) or waist circumference (WC) or waist-to-hip ratio (WHR) baseline information. Therefore, results from the current study were based on data from 4,656 participants, accounting for 76.3% of the initially invited individuals.

Questionnaires

A standard questionnaire was designed to collect basic information which included age, gender, marital status, educational background, occupation, smoking, and alcohol consumption. The marital status was divided into married/cohabiting and unmarried (single, divorced, or widowed). Educational attainment was categorized into three groups: illiterate/ elementary school, junior high school, and senior high school or higher. Occupational physical activity (PA) was categorized into light (housewife, retired, or unemployed), moderate (teacher, doctor, or nurse), and heavy (worker, farmer, or fisherman), according to occupation. Smoking was classified into current smokers (smoking every day and occasional smoking in the past years) and noncurrent smokers (including ex-smokers and nonsmokers). Alcohol consumption was categorized into regular drinkers (drinking every day in the past years) and noncurrent drinkers (including ex-drinkers, rare drinkers, and nondrinkers). Personal monthly income was categorized into ≤ 999 Chinese Yuan (CNY), 1,000–2,999 CNY, and $\geq 3,000$ CNY.

The plasma glucose level was determined by the glucose oxidase method. All subjects (without prediagnosed diabetes) underwent a standard 2-h 75-g oral glucose tolerance test. Diabetes was diagnosed on the basis of the WHO/International Diabetes Federation criteria (23). Subjects with fasting plasma glucose (FPG) level of ≥ 7.0 mmol/l and/or 2-h postload plasma glucose (2-hPG) level of ≥ 11.0 mmol/l were diagnosed with diabetes. The individual was taken as nondiabetic if FPG was < 7.0 mmol/l and/or 2-hPG < 11.1 mmol/l.

Blood pressure was measured using a mercury sphygmomanometer. Measurements were taken three times in 5-min intervals. The mean of the three readings was used for data analysis. Hypertension was defined by a mean systolic blood pressure of ≥ 140 mmHg and/or a mean diastolic blood pressure of ≥ 90 mmHg and/or an established diagnosis of hypertension at the baseline (24).

Anthropometric Measurement

We measured the participant's height and weight while wearing light clothes and no shoes. The BMI was calculated as the weight in kilograms divided by the height in squared meters (kg/m^2). Waist circumference (WC) was measured at the midpoint between the lower rib margin and the iliac crest. Hip circumference (HC) was measured at the maximal horizontal

girth between the waist and the thigh. The WHR was calculated by dividing WC (in cm) by HC (cm).

Assessment of Depression

Depression was assessed using the Zung Depression Rating Scale (ZDRS) (25, 26). The ZDRS questionnaire contains 20 questions divided into 10 positively and 10 negatively phrased questions. Each question was scored as 1 through 4, and the total scores ranged from 20 to 80 based on the participant's feelings (sadness, indifference, or happiness) toward family, work, and living conditions, respectively. Participants were divided into two major groups: normal (20–44) and depressed (≥ 45) (25).

Criteria for Obesity

Overweight and obesity were defined in accordance with the published criteria (26, 27). According to the Chinese criteria, subjects were divided into four BMI categorical groups as: underweight (BMI < 18.5 kg/m²), normal weight (18.5–23.9 kg/m²), overweight (24.0–27.9 kg/m²), and obese (≥ 28.0 kg/m²). According to Asian criteria, subjects were subdivided as: underweight (BMI < 18.5 kg/m²), normal weight (18.5–22.9 kg/m²), overweight (23.0–24.9 kg/m²), and obese (≥ 25.0 kg/m²) (27) and according to the WHO criteria as: underweight (BMI < 18.5 kg/m²), normal weight (18.5–24.9 kg/m²), overweight (25.0–29.9 kg/m²), and obese (≥ 30.0 kg/m²) (28).

Based on the value of WC, subjects were further subdivided into two groups as follows: normal (< 85.0 cm for males and < 80.0 cm for females); abdominal obesity, (≥ 85.0 cm for males and ≥ 80.0 cm for females) according to Chinese criteria (26). Alternatively, subjects were subdivided into: normal (< 90.0 cm for males and < 80.0 cm for females) and abdominal obesity, (≥ 90.0 cm for males and ≥ 80.0 cm for females) according to the Asian guidelines (27). Finally, according to the WHO guidelines, subjects were divided into normal (< 94.0 cm for males and < 80.0 cm for females) and abdominal obesity (≥ 94.0 cm for males and ≥ 80.0 cm for females) (28).

Subjects were divided into two categories based on their WHR cutoff points (Chinese guidelines): normal (< 0.9 for males and < 0.85 for females) and abdominal obesity (≥ 0.90 for males and ≥ 0.85 for females) (26); or the Asian WHR cutoff points: normal (< 0.95 for males and < 0.80 for females) and abdominal obesity (≥ 0.95 for males and ≥ 0.80 for females) (29) or the WHO WHR cutoff points: normal (< 1.00 for males and < 0.85 for females) and abdominal obesity (≥ 1.00 for males and ≥ 0.85 for females) (28).

Statistical Analysis

Statistical analyses were performed using IBM SPSS Statistics 17.0. Continuous variables and categorical variables were presented as the mean \pm standard deviation and the number (percentages), respectively. Using data from the 2010 census in Qingdao, the age-standardized prevalence of depression was calculated according to different obesity criterion for the age group of 35–74 years. Logistic regression was used to determine the possible link between the risk for depression and different indicators of obesity (BMI, WC, and WHR). Odds ratios (ORs) and 95% confidence intervals (CIs) were estimated by logistic

regression to obtain the association between depression and each variable. Sensitivity analysis was done based on the Asian and WHO criteria of obesity. A *P*-value < 0.05 was considered to be statistically significant.

RESULTS

The demographic information, social-economic status, lifestyle information, and anthropometric measurement have been shown in **Table 1**. In this study, a total of 4,656 participants were enrolled; among them, 1,804 were males (38.6%) and 2,852 were females (61.4%). The mean age was 52.3 ± 10.7 years (rang 35–74 years).

TABLE 1 | Characteristics of the population included in the study.

Characteristic	Total N = 4,656
Sex, female, <i>n</i> (%)	2852 (61.3)
Age, years (mean \pm SD)	52.3 \pm 10.7
Zung score (mean \pm SD)	29.8 \pm 7.9
Depression, <i>n</i> (%)	243 (5.2)
BMI (kg/m ² , mean \pm SD)?	25.0 \pm 3.7
Normal, <i>n</i> (%)	1871 (40.2)
Underweight, <i>n</i> (%)	83 (1.8)
Overweight, <i>n</i> (%)	1815 (39.0)
Obese, <i>n</i> (%)	887 (19.1)
WC (cm, mean \pm SD)?	83.8 \pm 10.6
Normal, <i>n</i> (%)	1982 (42.6)
Abdominal obesity, <i>n</i> (%)	2674 (57.4)
WHR (mean \pm SD) ?	0.87 \pm 0.07
Normal, <i>n</i> (%)	2306 (49.5)
Abdominal obesity, <i>n</i> (%)	2350 (50.5)
Hypertension, <i>n</i> (%)	2159 (46.5)
Diabetes, <i>n</i> (%)	741 (15.9)
Urban living, <i>n</i> (%)	1384 (29.7)
Married, <i>n</i> (%)	4361 (93.7)
EDUCATIONAL ATTAINMENT, <i>n</i> (%)	
Illiterate /elementary school	1904 (41.0)
Junior high school	1723 (37.1)
Senior high school or higher	1022 (22.0)
OCCUPATIONAL PA, <i>n</i> (%)	
Light	483 (10.5)
Moderate	346 (7.5)
Heavy	3796 (82.1)
Current smoker, <i>n</i> (%)	1137 (24.8)
Current drinker, <i>n</i> (%)	751 (16.1)
INCOME (CNY/MONTH), <i>n</i> (%)	
≤ 599	2372 (52.4)
600–1999	1820 (40.2)
≥ 2000	332 (7.4)

BMI, body mass index; WC, waist circumference; HC, hip circumference; WHR, waist-to-hip ratio; PA, physical activity; CNY, Chinese Yuan. ? obesity and abdominal obesity is defined according to Chinese criteria. *n*, number.


According to the BMI category, the age-standardized prevalence of depression was higher in underweight (10.5%) and overweight participants (6.0%) compared with the normal weight (4.3%); however, the difference was not statistically significant. Based on the WC category, a slightly higher age-standardized prevalence of depression was observed in the abdominal obesity participants (5.7%) compared with the normal WC participants (4.3%), and the difference was statistically significant. The corresponding age-standardized prevalence of depression was 5.5% in normal WHR participants and 4.7% in the abdominal obesity participants, respectively, and the difference was nonsignificant. The age-standardized prevalence of depression according to BMI, WC, and WHR has shown in **Figure 1**.

In this study, we determined the role of potential risk factors in depression using multivariable regression analysis according to the Chinese criterion (**Table 2**). Overweight indicated higher risk of depression (OR: 1.49, 95%CI: 1.08-2.05) compared with normal weight. We observed a significant relationship between abdominal obesity and depression risk according to WC.

Next, we performed a sensitivity analysis according to the Asian and WHO criteria (model 1 and model 2, respectively; **Table 3**). In model 1, sensitivity analysis based on Asian criterion revealed that overweight and obesity (BMI category) were not significantly associated with depression, while abdominal obesity (according to WC) was significantly associated with depression similar to that of Chinese criterion. In model 2, based on the WHO criterion, abdominal obesity (according to WC) was not significantly associated with depression, while overweight participants in the BMI category were significantly associated with depression.

DISCUSSION

In this study, we investigated the association between depression and body weight among adult people in Qingdao, China. Our results showed that the prevalence of depression was higher among the overweight participants (according to BMI category in China) and abdominal obesity participants (according to WC category in China) compared with other groups. Additionally, we defined prevalence of depression by using a Zung score equal to or greater than 45. In Qingdao, the overall prevalence of depression rate of 5.2% (5.5% in males and 5.0% in females) was lower than the documented rates in South Korea (overall 5.7%, 3.9% in males and 7.0% in females) (30). Additionally, a cross-sectional study of 512 891 Chinese adults, aged 30–79 years, had also revealed a lower pooled prevalence of depression compared with the current study (2.4%) (31). This difference might be due to social and family factors. Residents of the eastern coastal areas could not convincingly adapt to rapid industrialization and urbanization; thus it lead to depression. Secondly, accompanied with China entering the aging society, the “empty nest” phenomenon is the leading cause of unaccompanied, unattended, and psychosocial problems (such as depression) in middle-aged and elderly adults (32, 33). Depression, in turn, affects the middle-aged and the elderly population’s social and physical



activities, which strongly influenced the wellbeing of elderly adults (34).

The prevalence of depression is higher in underweight than normal individuals. However, there was an insignificant positive association between underweight and depression in general adults in Mainland China. Our results are inconsistent with the estimates reported in Japan, Korea, and Taiwan (13, 30, 35, 36). These trends can be observed in different age populations, and underweight elderly adults are far more likely to be depressed (13, 30, 35, 36). Yet, the common population tends to have a distortion of body weight, which entitles thinness as a beauty symbol owing to social standard (37). Our observations also support this fact.

TABLE 2 | Risk factors associated with depression determined by multivariable logistic regression according to the Chinese criterion.

	Model 1	Model 2	Model 3
Female	1.09 (0.72–1.62)	1.05 (0.70–1.56)	1.09 (0.73–1.62)
age	1.01 (1.00–1.03)	1.01 (1.00–1.03)	1.01 (1.00–1.03)
BMI CATEGORY			
Normal	1.00		
Underweight	1.24 (0.44–3.50)		
Overweight	1.48 (1.08–2.03)*		
Obese	1.16 (0.76–1.78)		
WC CATEGORY			
Abdominal obesity		1.47 (1.08–2.00)*	
WHR CATEGORY			
Abdominal obesity			0.86 (0.64–1.17)
Hypertension	0.81 (0.60–1.10)	0.78 (0.58–1.06)	0.86 (0.64–1.17)
Diabetes	0.81 (0.55–1.21)	0.80 (0.54–1.19)	0.84 (0.56–1.24)
Urban living	1.45 (0.95–2.22)	1.44 (0.95–2.20)	1.49 (0.98–2.28)
Unmarried	1.96 (1.26–3.06)*	1.93 (1.24–3.01)*	1.93 (1.24–3.01)*
EDUCATIONAL ATTAINMENT			
Illiterate	1.00	1.00	1.00
/elementary school			
Junior high school	0.81 (0.55–1.19)	0.79 (0.54–1.16)	0.79 (0.54–1.17)
Senior high school or higher	0.69 (0.41–1.14)	0.68 (0.41–1.13)	0.66 (0.40–1.11)
OCCUPATIONAL PHYSICAL ACTIVITY			
Light	1.00	1.00	1.00
Moderate	0.88 (0.48–1.61)	0.86 (0.47–1.56)	0.90 (0.49–1.64)
Heavy	0.71 (0.46–1.11)	0.72 (0.46–1.12)	0.72 (0.46–1.12)
Current smoker	0.76 (0.51–1.15)	0.77 (0.51–1.16)	0.79 (0.52–1.19)
Current drinker	0.85 (0.55–1.31)	0.84 (0.55–1.30)	0.86 (0.56–1.32)
INCOME (CHINESE YUAN/MONTH)			
≤599	1.00	1.00	1.00
600–1999	0.91 (0.49–1.71)	0.92 (0.49–1.71)	0.91 (0.48–1.70)
≥2000	1.14 (0.65–2.00)	1.14 (0.65–2.00)	1.12 (0.64–1.98)

Model 1, model 2 and Model 3, the risk factors listed in the table were entered at the same time. * $P < 0.05$ for factors associated with depression.

The prevalence of depression was higher in overweight and abdominal obesity (WC category) participants compared with the normal participants. In contrast, we did not observe a similar association in obese and abdominal obesity (WHR category) participants. This trend can be explained by the Chinese cultural heritage that usually associates overweight with a higher economic status, since those who can afford to eat more could attain more body weight and vice versa. In China, prevalence of obesity was substantially higher in rich people (38, 39). The overweight carries a social stigma, such as body image, self-esteem, and social life, which can help a bit for depression (40).

In this study, our analysis revealed a significant positive association between being overweight and depression, while a nonsignificant association between being obese and depression. However, the association of BMI with depression is still controversial. A nonsignificant positive relationship between

TABLE 3 | Sensitivity analyses for depression according to the Asian and WHO criteria.

	Model 1	Model 2
BMI CATEGORY		
Normal	1.00	1.00
Underweight	1.20 (0.42–3.42)	1.14 (0.41–3.21)
Overweight	1.14 (0.75–1.71)	1.39 (1.03–1.87)*
Obese	1.33 (0.95–1.88)	0.56 (0.28–1.13)
WC CATEGORY		
Abdominal obesity	1.41 (1.03–1.91)*	1.32 (0.96–1.82)
WHR CATEGORY		
Abdominal obesity	1.19 (0.83–1.70)	1.02 (0.72–1.46)

Model 1 is according to Asia criterion. Model 2 is according to WHO criterion. All models were adjusted for gender, age, hypertension, diabetes, resident districts, marital status, educational attainment, occupational physical activity, smoking status, alcohol drinking status, personal monthly income. BMI category, WC category, and WHR category entered models separately. * $P < 0.05$ for factors associated with depression.

being overweight and depression has been reported in developed countries such as Canada (41). Additionally, Palinkas et al. observed a nonsignificant inverse association between being overweight and depression in women (42). Moreover, a nonsignificant negative association between depression and obesity has also been reported previously (12, 43). In contrast, several studies reported a significant positive association between depression and obesity (11, 12, 20, 21, 44, 45). Young people regarded thinness as a beauty symbol. However, middle-aged and elderly people are still more likely to view obesity as a symbol of wealth and happiness in the traditional Chinese culture. Meanwhile, in this study, participants are mainly middle-aged and the elderly people aged 35 years or over, which is the possible reason for the current observations.

Next, we examined the effect of abdominal obesity on depression. We observed a significant positive association between abdominal obesity and depression. These results are in agreement with those of Takeuchi et al. (46) and Vogelzangs et al. (47). In contrast, Herva et al. (48), Gil et al. (49), and Zavala GA (12) reported a nonsignificant positive association between depression and abdominal obesity. The absence of significant associations between abdominal obesity (WHR category) and depression were inconsistent with the results reported by Ahlberg et al. (50).

Here, for the first time, we reported a significant association between being overweight, abdominal obesity (WC category), and depression in the general population of Qingdao China, but not in obese or abdominal obesity (WHR category) participants, which might be due to the following reasons. First, cultural factors are known to influence the association between depression and body weight. A higher socioeconomic status is generally associated with more concern for body image and abnormal body weight, which can possibly be a stress factor that leads to depression (51, 52). Second, diet is an important factor in the association between depression and body weight status. Emotions of individuals can also affect food

intake. Depression was associated with emotional eating such as negative emotional overeaters (53). Moreover, a negative emotion, such as depression, is further inclined to uncontrolled eating and/or overeating (54). However, this will ultimately lead to a high body weight gain. Finally, we cannot underestimate the roles of neurological mechanisms (16, 17, 55) and/or genetic susceptibility (56), which may also lead to depression and an abnormal weight status.

Previous studies had reported conflicting results in China, partly because different studies had used different criteria to define obesity. Therefore, in this study, we performed sensitivity analyses according to the Asian and WHO criteria of obesity. Sensitivity analyses showed a significant positive association between depression and abdominal obesity (WC category in Asian criteria), and being overweight (BMI category in WHO criteria) after adjustment for gender, age, hypertension, diabetes, resident districts, marital status, educational attainment, occupational PA, smoking status, alcohol-drinking status, and personal monthly income. These results were slightly different from the results produced using the Chinese criteria. Taken together, we suggest that the association between depression and weight status is influenced by the criteria adopted. Additionally, the reciprocal and complex associations of being overweight, abdominal obesity, and depression are strongly intertwined, and these phenomena are likely to be true for the earlier-mentioned results. Hence, further large-scale research is needed to reveal the association between obesity and depression among the whole Chinese population.

This study suffered from a few limitations. First, the present study was a cross-sectional study that does not reflect the underlying mechanisms between depression and weight status. Therefore, a follow-up study is imperative. Second, this study was performed on a relatively small sample size of the adult community in Qingdao, China.

CONCLUSION

We observed that overweight and abdominal obesity (WC category) participants were at a higher risk of depression according to Chinese criterion. However, abdominal obesity according to WHR was not associated with depression.

REFERENCES

1. Available online at: <http://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
2. Available online at: <http://www.who.int/news-room/fact-sheets/detail/depression>
3. Xi B, Liang Y, He T, Reilly KH, Hu Y, Wang Q et al. Secular trends in the prevalence of general and abdominal obesity among Chinese adults, 1993–2009. *Obes Rev.* (2012) 13:287–96. doi: 10.1111/j.1467-789X.2011.00944.x
4. Hu L, Huang X, You C, Li J, Hong K, Li P, et al. Prevalence of overweight, obesity, abdominal obesity and obesity-related risk factors in southern China. *PLoS ONE* (2017) 12:e0183934. doi: 10.1371/journal.pone.0183934
5. Wu J, Xu H, He X, Yuan Y, Wang C, Sun J, et al. Six-year changes in the prevalence of obesity and obesity-related diseases in Northeastern China from 2007 to 2013. *Sci Rep.* (2017) 7:41518. doi: 10.1038/srep41518

Furthermore, Asian and WHO criteria of obesity might influence the association between depression and body weight status. On all accounts, controlling being overweight and having abdominal obesity had a protective impact on depression. Future research will involve a larger multicenter study in China to further investigate the relationship between body weight and depression.

ETHICS AND CONSENT TO PARTICIPATE

All the participants voluntarily signed the informed consent before their participation and the consent of ethics, including three urban districts (Shinan, Shibe, and Sifang) and three rural districts (Huangdao, Jiaonan, and Jimo), was obtained from the ethics committee in Qingdao Municipal Center for Disease Control and Prevention. This study was approved by the local ethics committee at Qingdao Municipal Health and Family planning commission.

AVAILABILITY OF DATA AND MATERIALS

The aggregate data supporting findings contained within this manuscript will be shared upon request submitted by the corresponding author. Identifying patient data will not be shared.

AUTHOR CONTRIBUTIONS

ZB, JC, and JS were the primary authors and leading investigators. NY, JC, MK, ZB, and HX carried out the experiments, analyzed experimental results. ZB, JC, and JS wrote the manuscript. All authors read and approved the final manuscript.

FUNDING

This study was funded by the Bayer Healthcare in China and World Diabetes Foundation [WDF07-308].

ACKNOWLEDGMENTS

We are grateful to the participants, primary care doctors, and nurses who participated in this survey.

6. Cesar C. *Depression Exact a Heavy Toll on Chinese People's Health*. Counter punch (2018).
7. Eidsdottir ST, Kristjansson AL, Sigfusdottir ID, Garber CE, Allegrante JP. Association between higher BMI and depressive symptoms in Icelandic adolescents: the mediational function of body image. *Eur Public Health J.* (2014) 24:888–92. doi: 10.1093/eurpub/ckt180
8. de Wit LM, van Straten A, van Herten M, Penninx BW, Cuijpers P. Depression and body mass index, a u-shaped association. *BMC Public Health* (2009) 9:14. doi: 10.1186/1471-2458-9-14
9. Marijnissen RM, Bus BA, Holewijn S, Franke B, Purandare N, de Graaf J, et al. Depressive symptom clusters are differentially associated with general and visceral obesity. *J Am. Geriatr Soc.* (2011) 59:67–72. doi: 10.1111/j.1532-5415.2010.03228.x
10. Johnston E, Johnson S, McLeod P, Johnston M. The relation of body mass index to depressive symptoms. *Can J Public Health* (2004) 95:179–83.

11. Gibson-Smith D, Bot M, Snijder M, Nicolaou M, Derks EM, Stronks K, et al. The relation between obesity and depressed mood in a multi-ethnic population. The HELIUS study. *Soc Psychiatry Psychiatr Epidemiol.* (2018) 53:629–38. doi: 10.1007/s00127-018-1512-3
12. Zavala GA, Kolovos S, Chiarotto A, Bosmans JE, Campos-Ponce M, Rosado JL et al. Association between obesity and depressive symptoms in Mexican population. *Soc Psychiatry Psychiatr Epidemiol.* (2018) 53:639–46. doi: 10.1007/s00127-018-1517-y
13. Chang HH, Yen ST. Association between obesity and depression: evidence from a longitudinal sample of the elderly in Taiwan. *Aging Ment Health* (2012) 16:173–80. doi: 10.1080/13607863.2011.605053
14. Qian J, Li N, Ren X. Obesity and depressive symptoms among Chinese people aged 45 and over. *Sci Rep.* (2017) 7:45637. doi: 10.1038/srep45637
15. Li ZB, Ho SY, Chan WM, Ho KS, Li MP, Leung GM, et al. Obesity and depressive symptoms in Chinese elderly. *Int J Geriatr Psychiatry* (2004) 19:68–74. doi: 10.1002/gps.1040
16. Tindell AJ, Smith KS, Berridge KC, Aldridge JW. Dynamic computation of incentive salience: “wanting” what was never “liked”. *J Neurosci.* (2009) 29:12220–8. doi: 10.1523/JNEUROSCI.2499-09.2009
17. Nestler EJ. Epigenetics: Stress makes its molecular mark. *Nature* (2012) 490:171–2. doi: 10.1038/490171a
18. Puhl R, Brownell KD. Ways of coping with obesity stigma: review and conceptual analysis. *Eat Behav.* (2003) 4:53–78. doi: 10.1016/S1471-0153(02)00096-X
19. Sun Y, Liu Y, Tao FB. Associations Between Active Commuting to School, Body Fat, and Mental Well-being: Population-Based, Cross-Sectional Study in China. *J Adolesc Health* (2015) 6:679–85. doi: 10.1016/j.jadohealth.2015.09.002.
20. Luo H, Li J, Zhang Q, Cao P, Ren X, Fang A, et al. Obesity and the onset of depressive symptoms among middle-aged and older adults in China: evidence from the CHARLS. *BMC Public Health* (2018) 18:909. doi: 10.1186/s12889-018-5834-6
21. Qin T, Liu W, Yin M, Shu C, Yan M, Zhang J, et al. Body mass index moderates the relationship between C-reactive protein and depressive symptoms: evidence from the China Health and Retirement Longitudinal Study. *Sci Rep.* (2017) 7:39940. doi: 10.1038/srep39940
22. Zhang Y, Sun J, Pang Z, Wang X, Gao W, Ning F, et al. The impact of new screen-detected and previously known type 2 diabetes on health-related quality of life: a population-based study in Qingdao, China. *Qual. Life Res.* (2014) 23:2319–26. doi: 10.1007/s11136-014-0674-z
23. M. I. O. C. *Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycemia.* Geneva World Health Organization (2006).
24. 1999 World Health Organization-International Society of Hypertension Guidelines for the Management of Hypertension. Guidelines Subcommittee. (1999) *J Hypertens.* 17:151–83.
25. Fountoulakis KN, Lacovides A, Samolis S, Kleanthous S, Kaprinis SG, St K, et al. Reliability, validity and psychometric properties of the Greek translation of the Zung Depression Rating Scale. *BMC Psychiatry* (2001) 1:6. doi: 10.1186/1471-244X-1-6
26. Disease Control CM. *Chinese Guidelines on Overweight and Obesity Prevention and Control in Adults.* Beijing: People’s Medical Publishing House (2006).
27. WE C. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* (2004) 363:157–63. doi: 10.1016/S0140-6736(03)15268-3
28. WE C. *Waist Circumference and Waist-Hip Ratio: Report of a WHO Expert Consultation Geneva, 8-11 December 2008.* World Health Organization, (2008).
29. Lean ME, Han TS, Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ* (1995) 311:158–61.
30. Oh J, Chae JH, Kim TS. Age-specific association between body mass index and depression: The Korea National Health and Nutrition Examination Survey 2014. *Int J Obes.* (2018) 42:327–33. doi: 10.1038/ijo.2017.234
31. Chen Y, Bennett D, Clarke R, Guo Y, Yu C, Bian Z, et al. Patterns and correlates of major depression in Chinese adults: a cross-sectional study of 0.5 million men and women. *Psychol Med.* (2017) 47:958–70. doi: 10.1017/S0033291716002889
32. Wang G, Hu M, Xiao SY, Zhou L. Loneliness and depression among rural empty-nest elderly adults in Liuyang, China: a cross-sectional study. *BMJ Open* (2017) 7:e016091. doi: 10.1136/bmjopen-2017-016091
33. Gao YL, Wei YB, Shen YD, Tang YY, Yang JR. China’s empty nest elderly need better care. *J Am Geriatr Soc.* (2014) 62:1821–2. doi: 10.1111/jgs.12997
34. Xie LQ, Zhang JB, Peng F, Jiao NN. Prevalence and related influencing factors of depressive symptoms for empty-nest elderly living in the rural area of Yongzhou, China. *Arch Gerontol Geriatr.* (2010) 50:24–9. doi: 10.1016/j.archger.2009.01.003
35. Kuo SY, Lin KM, Chen CY, Chuang YL, Chen WJ. Depression trajectories and obesity among the elderly in Taiwan. *Psychol. Med.* (2011) 41:1665–76. doi: 10.1017/S0033291710002473
36. Hides S, Asano S, Saito K, Sasayama D, Kunugi H. Association of depression with body mass index classification, metabolic disease, and lifestyle: A web-based survey involving 11,876 Japanese people. *J Psychiatr Res.* (2018) 102:23–8. doi: 10.1016/j.jpsychires.2018.02.009
37. Cash TF. Body image: past, present, and future. *BODY IMAGE* (2004) 1:1–5. doi: 10.1016/S1740-1445(03)00011-1
38. Xiao Y, Zhao N, Wang H, Zhang J, He Q, Su D, et al. Association between socioeconomic status and obesity in a Chinese adult population. *BMC Public Health* (2013) 13:355. doi: 10.1186/1471-2458-13-355
39. Zhao P, Gu X, Qian D, Yang F. Socioeconomic disparities in abdominal obesity over the life course in China. *Int J Equity Health* (2018) 17:96. doi: 10.1186/s12939-018-0809-x
40. Phelan SM. An update on research examining the implications of stigma for access to and utilization of bariatric surgery. *Curr Opin Endocrinol Diabetes Obes.* (2018) 25:321–5. doi: 10.1097/MED.0000000000000431
41. Chen Y, Jiang Y, Mao Y. Association between obesity and depression in Canadians. *J Womens Health* (2009) 18:1687–92. doi: 10.1089/jwh.2008.1175
42. Palinkas LA, Wingard DL, Barrett-Connor E. Depressive symptoms in overweight and obese older adults: a test of the “jolly fat” hypothesis. *J Psychosom Res.* (1996) 40:59–66.
43. Wade TW, Oberhelman SS, Angstman KB, Sawchuk CN, Meunier MR, Angstman GL, et al. Diabetes and obesity not associated with 6-month remission rates for primary care patients with depression. *Psychosomatics* (2015) 56:354–61. doi: 10.1016/j.psym.2014.05.012
44. de Wit L, Luppino F, van Straten A, Penninx B, Zitman F, Cuijpers P. Depression and obesity: a meta-analysis of community-based studies. *Psychiatry Res.* (2010) 178:230–5. doi: 10.1016/j.psychres.2009.04.015
45. Zhong W, Cruickshanks KJ, Schubert CR, Nieto FJ, Huang GH, Klein BE, et al. Obesity and depression symptoms in the Beaver Dam Offspring Study population. *Depress. Anxiety* (2010) 27:846–51. doi: 10.1002/da.20666
46. Takeuchi T, Nakao M, Nomura K, Yano E. Association of metabolic syndrome with depression and anxiety in Japanese men. *Diabetes Metab.* (2009) 35:32–6. doi: 10.1016/j.diabet.2008.06.006
47. Vogelzangs N, Beekman AT, Kritchevsky SB, Newman AB, Pahor M, Yaffe K, et al. Psychosocial risk factors and the metabolic syndrome in elderly persons: findings from the Health, Aging and Body Composition study. *J Gerontol A Biol Sci Med Sci.* (2007) 62:563–9.
48. Herva A, Rasanen P, Miettunen J, Timonen M, Laksy K, Veijola J, et al. Co-occurrence of metabolic syndrome with depression and anxiety in young adults: the Northern Finland 1966 Birth Cohort Study. *Psychosom Med.* (2006) 68:213–6. doi: 10.1097/01.psy.0000203172.02305.ea
49. Gil K, Radzillowicz P, Zdrojewski T, Pakalska-Korcala A, Chwojnicki K, Pivonski J, et al. Relationship between the prevalence of depressive symptoms and metabolic syndrome. Results of the SOPKARD Project. *Kardiol Pol.* (2006) 64:464–9.
50. Ahlberg AC, Ljung T, Rosmond R, McEwen B, Holm G, Akesson HO, et al. Depression and anxiety symptoms in relation to anthropometry and metabolism in men. *Psychiatry Res.* (2002) 112:101–10.
51. Wu YK, Berry DC. Impact of weight stigma on physiological and psychological health outcomes for overweight and obese adults: a systematic review. *J Adv Nurs.* (2018) 74:1030–42. doi: 10.1111/jan.13511
52. Stevens SD, Herbozo S, Morrell HE, Schaefer LM, Thompson JK. Adult and childhood weight influence body image and depression through weight stigmatization. *J Health Psychol.* (2017) 22:1084–93. doi: 10.1177/1359105315624749

53. Bourdier L, Morvan Y, Kotbagi G, Kern L, Romo L, Berthoz S. Examination of emotion-induced changes in eating: A latent profile analysis of the Emotional Appetite Questionnaire. *Appetite* (2018) 123:72–81. doi: 10.1016/j.appet.2017.11.108
54. Bongers P, de Graaff A, Jansen A. 'Emotional' does not even start to cover it: Generalization of overeating in emotional eaters. *Appetite* (2016) 96:611–6. doi: 10.1016/j.appet.2015.11.004
55. Kelley AE, Berridge KC. The neuroscience of natural rewards: relevance to addictive drugs. *J Neurosci.* (2002) 22:3306–11.
56. Stunkard AJ, Faith MS, Allison KC. Depression and obesity. *Biol Psychiatry* (2003) 54:330–7.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Cui, Sun, Li, Ke, Sun, Yasmeen, Khan, Xin, Xue and Baloch. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.