



OPEN ACCESS

APPROVED BY

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE

Frontiers Production Office
production.office@frontiersin.org

SPECIALTY SECTION

This article was submitted to
Clinical Diabetes,
a section of the journal
Frontiers in Endocrinology

RECEIVED 12 September 2022

ACCEPTED 12 September 2022

PUBLISHED 29 September 2022

CITATION

Frontiers Production Office (2022)
Erratum: Dyslipidemia and 10-year
diabetes incidence in Japanese
people: Population-based Panasonic
cohort study 9.
Front. Endocrinol. 13:1042407.
doi: 10.3389/fendo.2022.1042407

COPYRIGHT

© 2022 Frontiers Production Office. This
is an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](#).
The use, distribution or reproduction
in other forums is permitted, provided
the original author(s) and the
copyright owner(s) are credited and
that the original publication in this
journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is
permitted which does not comply with
these terms.

Erratum: Dyslipidemia and 10-year diabetes incidence in Japanese people: Population-based Panasonic cohort study 9

Frontiers Production Office*

Frontiers Media SA, Lausanne, Switzerland

KEYWORDS

incident diabetes, LDL/HDL ratio, LDL cholesterol, HDL cholesterol, cohort study

An Erratum on

Dyslipidemia and 10-year diabetes incidence in Japanese people: Population-based Panasonic cohort study 9.

By Kobayashi G, Okada H, Hamaguchi M, Kurogi K, Murata H, Ito M and Fukui M (2022) *Front. Endocrinol.* 13:957728. doi: 10.3389/fendo.2022.957728

Due to a production error, there was a mistake in **Table 3** as published. The third sub-heading within the table was incorrectly captured as “Body mass index < 25 kg/m²” instead of “Body mass index ≥ 25 kg/m²”.

The corrected **Table 3** appears below.

The publisher apologizes for this mistake.

The original version of this article has been updated.

TABLE 3 The area under the curve and optimal cut-off values.

Whole participants	AUC	Sensitivity	Specificity	Cut-off value
Low-density lipoprotein cholesterol	0.613	64.2%	53.3%	124 mg/dl
High-density lipoprotein cholesterol	0.640	57.7%	63.7%	54 mg/dl
LDL/HDL ratio	0.662	60.7%	63.7%	2.4
Body mass index < 25 kg/m²	AUC	Sensitivity	Specificity	Cut-off value
Low-density lipoprotein cholesterol	0.595	57.6%	57.7%	124 mg/dl
High-density lipoprotein cholesterol	0.600	64.2%	50.6%	61 mg/dl
LDL/HDL ratio	0.624	54.4%	64.4%	2.3
Body mass index ≥ 25 kg/m²	AUC	Sensitivity	Specificity	Cut-off value
Low-density lipoprotein cholesterol	0.553	62.0%	46.0%	130 mg/dl
High-density lipoprotein cholesterol	0.577	64.0%	48.7%	52 mg/dl
LDL/HDL ratio	0.591	57.5%	56.6%	2.8

LDL, low-density lipoprotein; HDL, high-density lipoprotein.