



## OPEN ACCESS

APPROVED BY  
Frontiers Editorial Office,  
Frontiers Media SA, Switzerland

\*CORRESPONDENCE  
Chun-feng Lu  
lcf0413cwl1030@126.com  
Jian-bin Su  
sujbzjx@163.com  
Xue-qin Wang  
wangxueqin108@163.com

†These authors have contributed  
equally to this work

SPECIALTY SECTION  
This article was submitted to  
Clinical Diabetes,  
a section of the journal  
Frontiers in Endocrinology

RECEIVED 03 November 2022  
ACCEPTED 14 November 2022  
PUBLISHED 25 November 2022

CITATION  
Huang H-y, Huang Z-q, Hua L-y,  
Liu W-s, Xu F, Ge X-q, Lu C-f,  
Su J-b and Wang X-q (2022)  
Corrigendum: The association  
between normal serum sodium  
levels and bone turnover in  
patients with type 2 diabetes.  
*Front. Endocrinol.* 13:1088810.  
doi: 10.3389/fendo.2022.1088810

COPYRIGHT  
© 2022 Huang, Huang, Hua, Liu, Xu, Ge,  
Lu, Su and Wang. This is an open-  
access article distributed under the  
terms of the [Creative Commons  
Attribution License \(CC BY\)](#). The use,  
distribution or reproduction in other  
forums is permitted, provided the  
original author(s) and the copyright  
owner(s) are credited and that the  
original publication in this journal is  
cited, in accordance with accepted  
academic practice. No use,  
distribution or reproduction is  
permitted which does not comply with  
these terms.

# Corrigendum: The association between normal serum sodium levels and bone turnover in patients with type 2 diabetes

Hai-yan Huang<sup>1†</sup>, Zhi-qi Huang<sup>2†</sup>, Ling-yan Hua<sup>3†</sup>,  
Wang-shu Liu<sup>1</sup>, Feng Xu<sup>1</sup>, Xiao-qin Ge<sup>1</sup>, Chun-feng Lu<sup>1\*</sup>,  
Jian-bin Su<sup>1\*</sup> and Xue-qin Wang<sup>1\*</sup>

<sup>1</sup>Department of Endocrinology, Affiliated Hospital 2 of Nantong University and First People's Hospital of Nantong City, Nantong, China, <sup>2</sup>Department of General Surgery, Affiliated Hospital 2 of Nantong University and First People's Hospital of Nantong City, Nantong, China, <sup>3</sup>Department of Ophthalmology, Affiliated Hospital 2 of Nantong University and First People's Hospital of Nantong City, Nantong, China

## KEYWORDS

type 2 diabetes, bone turnover, bone formation, bone resorption, sodium, bone mineral density

## A corrigendum on

## The association between normal serum sodium levels and bone turnover in patients with type 2 diabetes


by Huang H-y, Huang Z-q, Hua L-y, Liu W-s, Xu F, Ge X-q, Lu C-f, Su J-b and Wang X-q (2022). *Front. Endocrinol.* 13:927223. doi: 10.3389/fendo.2022.927223

In the published article, there was an error in **Table 3** as published. We made a mistake in the title of the table when revising the manuscript. It was an oversight on our part that we typed the same “ $\beta$ ” and “p” values for CTx model 0 when we made this table. The corrected **Table 3** and its amended caption “Independent associations of serum sodium level with BTMs levels via multivariate linear regression analysis” appear below.

TABLE 3 Independent associations of serum sodium level with BTMs levels via multivariate linear regression analysis.

Models	B (95% CI)	$\beta$	t	p	R <sup>2</sup> for model
OC					
Model 0	0.038 (0.020-0.056)	0.210	4.141	<0.001	0.044
Model 1	0.040 (0.022-0.059)	0.222	4.249	<0.001	0.086
Model 2	0.037 (0.019-0.055)	0.204	3.957	<0.001	0.169
Model 3	0.025 (0.003-0.047)	0.134	2.281	0.023	0.227
CTx					
Model 0	0.023 (-0.002-0.058)	0.092	1.777	0.076	0.008
Model 1	0.031 (0.006-0.057)	0.124	2.391	0.017	0.096
Model 2	0.031 (0.005-0.058)	0.125	2.362	0.019	0.128
Model 3	0.019 (-0.011-0.050)	0.074	1.247	0.213	0.212
PINP					
Model 0	0.038 (0.019-0.058)	0.196	3.854	<0.001	0.039
Model 1	0.043 (0.024-0.063)	0.224	4.409	<0.001	0.133
Model 2	0.041 (0.022-0.061)	0.214	4.150	<0.001	0.169
Model 3	0.036 (0.013-0.059)	0.179	3.023	0.003	0.215

In the published article, there also was an error in **Figure 4** as published. We made the modifications on the same template when drawing the figures, but we missed the need to modify the  $p$  value. The corrected **Figure 4** and its caption “The relationship between serum sodium and lnPINP in patients with T2D (A) unadjusted; (B) partially adjusted for age, sex, BMI and HbA1c.” appear below.



The authors apologize for these errors and state that they do not change the scientific conclusions of the article in any way. The original article has been updated.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.