



OPEN ACCESS

EDITED BY

Surasak Saokaew,
University of Phayao, Thailand

REVIEWED BY

Olarik Musigavong,
Ministry of Public Health, Thailand
William J. Massey,
Cleveland Clinic, United States

*CORRESPONDENCE

Anna Bendarska-Czerwińska
[✉czerwinskaa002@gmail.com](mailto:czerwinskaa002@gmail.com)
Nikola Zmarzły
[✉nikola.zmarzly@gmail.com](mailto:nikola.zmarzly@gmail.com)
Beniamin Oskar Grabarek
[✉Bgrabarek7@gmail.com](mailto:Bgrabarek7@gmail.com)

SPECIALTY SECTION

This article was submitted to
Reproduction,
a section of the journal
Frontiers in Endocrinology

RECEIVED 15 June 2022

ACCEPTED 28 December 2022

PUBLISHED 17 January 2023

CITATION

Bendarska-Czerwińska A, Zmarzły N,
Morawiec E, Panfil A, Bryś K, Czarniecka J,
Ostenda A, Dziobek K, Sagan D, Boroń D,
Michalski P, Pallazo-Michalska V and
Grabarek BO (2023) Endocrine disorders
and fertility and pregnancy: An update.
Front. Endocrinol. 13:970439.
doi: 10.3389/fendo.2022.970439

COPYRIGHT

© 2023 Bendarska-Czerwińska, Zmarzły,
Morawiec, Panfil, Bryś, Czarniecka,
Ostenda, Sagan, Boroń, Michalski,
Pallazo-Michalska and Grabarek. This is an open-
access article distributed under the terms of
the [Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Endocrine disorders and fertility and pregnancy: An update

Anna Bendarska-Czerwińska^{1,2,3*}, Nikola Zmarzły^{4*},
Emilia Morawiec^{1,4,5}, Agata Panfil⁴, Kamil Bryś⁴,
Justyna Czarniecka⁴, Aleksander Ostenda², Konrad Dziobek²,
Dorota Sagan⁶, Dariusz Boroń^{4,7,8,9}, Piotr Michalski²,
Veronica Pallazo-Michalska² and Beniamin Oskar Grabarek^{1,4,7,8,9*}

¹Department of Molecular, Biology Gyncentrum Fertility Clinic, Katowice, Poland, ²Faculty of Medicine, Academy of Silesia, Zabrze, Poland, ³American Medical Clinic, Katowice, Poland, ⁴Department of Histology, Cytophysiology and Embryology, Faculty of Medicine, University of Technology, Academy of Silesia in Katowice, Zabrze, Poland, ⁵Department of Microbiology, Faculty of Medicine, University of Technology, Academy of Silesia in Katowice, Zabrze, Poland, ⁶Medical Center Dormed Medical SPA, Busko-Zdroj, Poland, ⁷Department of Gynaecology and Obstetrics, Faculty of Medicine, Academy of Silesia, Zabrze, Poland, ⁸Department of Gynecology and Obstetrics with Gynecologic Oncology, Ludwik Rydygier Memorial Specialized Hospital, Kraków, Poland, ⁹Department of Gynecology and Obstetrics, TOMMED Specjalisci od Zdrowia, Katowice, Poland

It is estimated that more and more couples suffer from fertility and pregnancy maintenance disorders. It is associated with impaired androgen secretion, which is influenced by many factors, ranging from genetic to environmental. It is also important to remember that fertility disorders can also result from abnormal anatomy of the reproductive male and female organ (congenital uterine anomalies – septate, unicornuate, bicornuate uterus; acquired defects of the uterus structure – fibroids, polyps, hypertrophy), disturbed hormonal cycle and obstruction of the fallopian tubes resulting from the presence of adhesions due to inflammation, endometriosis, and surgery, abnormal rhythm of menstrual bleeding, the abnormal concentration of hormones. There are many relationships between the endocrine organs, leading to a chain reaction when one of them fails to function properly. Conditions in which the immune system is involved, including infections and autoimmune diseases, also affect fertility. The form of treatment depends on infertility duration and the patient's age. It includes ovulation stimulation with clomiphene citrate or gonadotropins, metformin use, and weight loss interventions. Since so many different factors affect fertility, it is important to correctly diagnose what is causing the problem and to modify the treatment regimen if necessary. This review describes disturbances in the hormone secretion of individual endocrine organs in the context of fertility and the maintenance of pregnancy.

KEYWORDS

endocrine disorders, fertility, infertility, pregnancy, reproductive health, hormones

1 Introduction

Infertility is the inability to conceive within 12 months of regular intercourse (2-4 times a week) without using any contraceptive methods. Involuntary childlessness is a significant social problem faced by 20% of couples worldwide, and only in Poland the problem of infertility affects approximately 1.5 million couples each year (1).

Factors that influence fertility in both sexes include hyperprolactinemia, hypogonadotropic hypogonadism, infections, systemic diseases, and even lifestyle (2). More and more attention is also paid to the problem of obesity. Adipocytes act as an endocrine organ, and their excess promotes disorders of the hypothalamic-pituitary-ovarian axis. Secreted adipokines include leptin, adiponectin, resistin, interleukin 6, interleukin 1 β , and tumor necrosis factor α (TNF α), involved in inflammatory processes and the regulation of metabolism (3).

According to the recommendations of the Fertility and Infertility Section at the Polish Society of Gynecologists and Obstetricians and the Polish Society of Reproductive Medicine and Embryology, there are several causes of male infertility (4): 1) pre-testicular, i.e. related to the malfunctioning of the endocrine system in terms of impaired secretion of luteinizing hormone, sex steroids – testosterone, inhibin, follicle-stimulating hormone or resulting from mutations which cause impaired sperm movement or chromosomal aberrations observed, e.g. in Klinefelter's syndrome, 2) testicular, e.g. cryptorchidism, varicocele, infectious diseases; 3) extra-testicular, which include congenital absence of vas deferens, underdevelopment of seminal vesicles, epididymis defects, polycystic kidney disease, cystic fibrosis, diabetes; 4) sexual disorders – lack of erection or ejaculation, penis structure abnormalities (5). During the diagnosis of infertility, the assessment of male fertility should be obligatory, and the examination should be performed after maintaining a 2-7 day period of sexual abstinence (6).

On the other hand, infertility in women may be mainly due to disturbances in the occurrence of ovulatory cycles, caused, inter alia, by woman's age, hormonal disorders in the course of polycystic ovary syndrome (PCOS), hyperprolactinemia, abnormal anatomy of the reproductive organ (congenital uterine anomalies – septate, unicornuate, bicornuate uterus; acquired defects of the uterus structure – fibroids, polyps, hypertrophy), disturbed hormonal cycle and obstruction of the fallopian tubes resulting from the presence of adhesions due to inflammation, endometriosis, and surgery (7).

Diagnostics of female infertility should include: medical history regarding the regularity of menstruation, gynecological examination, determination of the concentration of selected sex hormones, including a single measurement of progesterone level 7 days before the planned menstruation to assess ovulation, ultrasound (USG) at the end of the follicular phase of the cycle (8, 9). However, in the case of ovulation disorders, manifested by an abnormal rhythm of menstrual bleeding, progesterone level lower than 2 ng/mL in the middle of the luteal phase, the diagnosis should be supplemented with the determination of the concentration of gonadotropins, androgens, thyroid-stimulating hormone (TSH), prolactin (PRL), anti-müllerian hormone (AMH) and assessment of the reproductive potential of the ovaries (10). The form of infertility treatment depends on its duration and the patient's age. It includes ovulation stimulation with clomiphene citrate or gonadotropins, intrauterine insemination, *in vitro* fertilization (11).

The aim of this review was to present disturbances in the hormone secretion of individual endocrine organs in the context of fertility of both sexes and the maintenance of pregnancy (Figure 1).

2 Differences between male and female embryogenesis

Sex differentiation is a complex process that depends on the activity of many genes. The key to sex dimorphism is the SRY (sex-determining region on Y) gene, whose protein product initiates the cascade of expression of genes determining the formation of testicles. The lack of this protein allows development towards a female individual (12, 13).

2.1 Undifferentiated stage

The sex of the embryo is determined already during fertilization, but until the 7th week of development, the gonads do not show gender-related differences. Initially, the gonads are in the form of longitudinal folds, which are formed due to the proliferation of the body cavity epithelium and the thickening of the stromal mesenchyme. Primordial germ cells appear no earlier than the 6th week of development and it is important for the gonads development that they populate the genital crests. Genital crest epithelial cells proliferate and penetrate into the stromal mesenchyme, where they form irregularly shaped primary sex cords. In both female and male embryos, these cords maintain a connection to the surface epithelium. This makes it impossible to distinguish between male and female gonads, which is why these structures are referred to as undifferentiated gonads (14).


Female and male embryos have two pairs of ducts: mesonephric (Wolffian) and paramesonephric (Müllerian). The caudal part of the connected ducts is incorporated into the posterior wall of the urogenital sinus, forming a sinus tubercle.

At 3 weeks of development, the mesenchyme cells migrate around the cloacal membrane, forming a pair of slightly elevated cloacal folds. They join cranially to form the genital tubercle, and caudally to the urethral and the anal folds. At the same time, another pair of genital eminences is visible on either side of the urethral folds. Later, they form the scrotal eminences in men and the labia majora in women (15).

2.2 Embryogenesis of the female reproductive system

2.2.1 Ovaries

In female development, the primary sex cords break down into irregular clusters of cells. During further development, they disappear and are replaced by vascularized stroma tissue, forming the ovarian core. The surface epithelial cells of the female gonad (as opposed to the epithelial cells of the male gonad) continue to divide, resulting in the formation of cortical cords around the 7th week of development. In the third month, they disintegrate into separate cell foci forming



follicular cells, creating, together with the oogonium, the primary (resting) follicle (16).

2.2.2 Genital ducts

In the presence of estrogen, the main female genital ducts are formed from the paramesonephric duct. With the descent of the ovary, part of the ducts transform into the fallopian tube and uterine canal. The connected paramesonephric ducts become the corpus and cervix and the upper part of the vagina. In female embryos, the mesonephric ducts regress due to a lack of testosterone (16).

2.2.3 External genitalia

The development of female external genitalia is stimulated by estrogens. The genital tubercle lengthens to form the clitoris, and the folds of the urethra give rise to the labia minora. In turn, the vaginal prominences enlarge and produce the labia majora. The urethral groove remains open and forms the vestibule (17).

2.3 Embryogenesis of the male reproductive system

2.3.1 Testes

The genetically male embryo, under the influence of the SRY gene product, develops sex cords that penetrate deeper towards the core part

of the gonad, forming medullary cords (13). These, in turn, continue to disintegrate, becoming the tubules of the testicle network. In further development, the testicular cords are separated from the surface epithelium by the tunica albuginea. In the fourth month, the testicular cords join the nucleus network. It is worth noting that the testicular cords retain the form of solid structures until puberty, and then transform into convoluted seminiferous tubules (14, 18).

2.3.2 Genital ducts

They develop under the influence of testosterone and originate from part of the mesonephric organ. With the exception of the appendix of the epididymis, the mesonephric duct does not regress and forms the main genital ducts. Below the orifice of the efferent ducts, the epididymal duct is elongated and formed. The mesonephric duct from the tail of the epididymis to the protrusion of the seminal vesicle is called the vas deferens.

2.3.3 External genitalia

The development of male external genitalia is stimulated by androgens and is manifested by the rapid lengthening of the genital tubercle, the penis. In the process of elongation, the penis pulls on the folds of the urethra, which form the urethral grooves. The furthest part is the penis glans. The epithelial lining of the groove forms the urethral plate. At the end of the 3rd month, the folds of the urethra close above the plate, forming the spongy part of the urethra, then the urethral lumen is

formed. In turn, the genital (scrotal) eminences move in the caudal direction during development and form the scrotum (17).

3 Hypothalamus and pituitary gland

The hypothalamus and pituitary gland control and regulate the proper functioning of the endocrine glands. The gonadotropic pathway is especially important in the case of fertility and the maintenance of pregnancy (Tables 1, 2). The hypothalamus produces gonadotropin-releasing hormone (GnRH), the appropriate concentration of which is necessary for the secretion of gonadotropins, i.e. luteinizing hormone (LH) and follicle-stimulating hormone (FSH) by the anterior (glandular) lobe of the pituitary gland (22). Gonadotropins, in turn, have a direct influence on the production of sex hormones, sperm and ova, as well as the course of pregnancy. Thus, the malfunction of the hypothalamus or pituitary gland will be associated with low production of GnRH and gonadotropins, which in turn will lead to the gonadal failure, known as hypogonadotropic hypogonadism (23).

Other hormones produced by the anterior pituitary gland include adrenocorticotropic hormone (ACTH) and growth hormone (GH). ACTH is produced from the precursor hormone proopiomelanocortin (POMC), which stimulates the production of cortisol in the adrenal glands *via* melanocortin receptors. Corticotropin-releasing hormone (CRH), along with vasopressin (ADH), are the main hormones that control ACTH secretion. GH and ACTH secretion is pulsatile and subject to circadian rhythms. Factors that influence the increase in growth hormone secretion are sleep and physical activity, as well as fasting, hypoglycemia, hypovolemia, and surgery. Hyperglycemia initially reduces and then increases GH levels (rebound effect). GH secretion shows gender differences, in males it is pulsatile and in females secretion is continuous. GH levels decline with age and somatopause occurs in the elderly (24).

3.1 Female

3.1.1 *In vivo* human studies

Growth hormone, by activating insulin-like growth factor 1 (IGF-1) synthesis, directly improves the quality of oocytes, and also increases

FSH-induced ovarian steroidogenesis. Administration of GH during ovulation stimulation enhances the effectiveness of the *in-vitro* procedure by increasing the percentage of mature oocytes and embryos viable to the day of transfer. Daily administration of a low dose of growth hormone (0.5 IU) started on the day of GnRH agonist administration increases the pregnancy rate (34.4% vs. 0%), the percentage of good-quality embryos and cryopreserved embryos. In addition, the use of a low dose of GH (4IU/d) in patients with reduced response to GnRH antagonists preparing for the IVF/ICSI procedure decreases the effective dose of gonadotropins and the duration of stimulation, increasing the total number of oocytes and oocytes in metaphase II of meiosis (48).

In obese women, reduced fertility results, among others, from decreased LH levels. They experience accelerated sexual maturation, menstrual disorders, in particular prolongation of the follicular phase, indicating ovulation disorders, and a higher frequency of obstetric complications, including spontaneous miscarriages (49).

Interestingly, the serine protease inhibitor (SERPINA12, also known as Vaspin) plays a significant role in the pathogenesis of type 2 diabetes, inflammation and infertility. Its concentration increases with weight gain and the worsening of insulin resistance, and decreases with weight loss. Metformin reduces the serum concentration of Vaspin in patients with PCOS (50).

Too low body weight, intense exercise, chronic diseases, malabsorption disorders such as celiac disease, hyperthyroidism, and thus energy deficiencies lead to impaired pulsatile GnRH secretion, and as a result, the concentration of LH and FSH necessary for effective stimulation of the gonads is too low (51). Studies have also shown that women with a BMI below 18.5 have a 72% higher risk of miscarriage in the first trimester of pregnancy than women with a normal body weight (52).

Disturbance of the circadian rhythm including sleep quality, wakefulness and the above-related pulsatile hormone secretion plays a significant role in infertility. It is known that insomnia in female shift workers results in decreased melatonin production. Low melatonin concentration in Graaf vesicles is associated with higher generation of free oxygen radicals (ROS), and thus poorer quality of oocytes. Melatonin supplementation in patients undergoing the *in vitro* procedure increases its effectiveness by improving the quality of oocytes, increasing the percentage of fertilization and the quality of the emerging embryos (24).

TABLE 1 Fertility studies related to the hypothalamus and pituitary gland.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Abbara et al. (19)	<i>in vivo</i> study (human) female	Comparative study	Evaluation of gonadotropin levels in a group of 243 patients with hyperprolactinemia	In patients with PCOS and hyperprolactinemia, there is an increase mainly in LH	Depending on the severity of hyperprolactinemia, a variable pattern of gonadotropin secretion is noted
Feng et al. (20)	<i>in vivo</i> study (human) female	Comparative study	Evaluation of the effectiveness of the use of Bushen-zhu-yun decoction (BSZY-D) and dopamine agonists in infertile patients with hyperprolactinemia	Adding decoction to traditional treatment results in a reduction in the number of miscarriages, fewer side effects, lower prolactin levels	Bushen-zhu-yun decoction (BSZY-D) and dopamine agonists are characterized by synergism of action, which is beneficial
Sermondade et al. (21)	<i>in vivo</i> study (human) male	Meta-analysis	Confirmation of the relationship between BMI and sperm count	Overweight and obesity were associated with an increased prevalence of azoospermia or oligozoospermia	BMI is associated with sperm count

TABLE 2 Fertility studies related to the pineal gland activity.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Espino et al. (25)	<i>in vivo</i> human female	Clinical experiment	Assessment of the effect of melatonin administration on reproduction during the use of an assisted reproductive protocol	The study group showed a higher percentage of pregnancies and live births	Melatonin may be a fertility-enhancing drug in women
Mokhtari et al. (26)	<i>in vivo</i> human female	Double-blinded randomized clinical trial	Assessment of the effect of melatonin administration on the fertility of women suffering from PCOS	The study group showed a higher percentage of pregnancies	Melatonin can potentially increase a chance of pregnancy
Celik et al. (27)	<i>in vivo</i> human female	Clinical experiment	Determining the relationship between melatonin concentration and the occurrence of cholestasis in pregnant women	Lower melatonin concentration was observed in the group of women with cholestasis	Melatonin concentration may be a predictor of the occurrence of maternal cholestasis
Hobson et al. (28)	<i>in vivo</i> human female	Clinical experiment	Assessment of melatonin use in pre-eclampsia	Blood pressure was lower in the study group	Melatonin can be used in pre-eclampsia to lower blood pressure
Zhang et al. (29)	<i>in vivo</i> animal (mouse) female	Animal medical experiment	Evaluation of the effect of melatonin concentration on female reproductive performance	Melatonin deficiency reduces the gonads and reproductive potential of females	The decreased melatonin concentration reduces the reproductive potential
Lv et al. (30)	<i>in vivo</i> animal (mouse) female	Animal medical experiment	Assessment of the effect of melatonin and leptin on female reproductive potential	Blocking the melatonin mt1 receptor causes a decrease in the concentration of leptin and steroid hormones	Melatonin is involved in leptin-mediated regulation of steroid hormone level
Lombardo et al. (31)	<i>in vivo</i> animal (Fundulus heteroclitus) female	Animal medical experiment	Evaluation of the influence of melatonin concentration on the survival of the embryo	There is a positive correlation between melatonin concentration and embryo survival	The higher the melatonin concentration, the higher the probability of embryo survival
Zhang et al. (32)	<i>in vivo</i> animal (mouse) female	Animal medical experiment	Evaluation of the influence of the MTNR1A polymorphism on the reproductive potential of females	Melatonin has a positive effect on reproduction	Higher melatonin levels may positively affect fertility
Cosso et al. (33)	<i>in vivo</i> animal (ewe lamb) female	Animal medical experiment	Evaluation of the effect of melatonin administration on female reproduction	Higher fertility and an earlier first estrus were shown in the study group	Melatonin may be a fertility enhancer for llamas
Song et al. (34)	<i>in vivo</i> animal (mice) female	Animal medical experiment	Evaluation of the effect of melatonin administration on female reproduction	The study group showed a decrease in ovarian aging, higher fertility and oocyte quality	Melatonin may increase fertility in mice
Dholpuria et al. (35)	<i>in vivo</i> animal (camel) female	Animal medical experiment	Evaluation of the effect of melatonin-releasing implants on female fertility	The study group showed a higher percentage of pregnancies	Melatonin may be a fertility drug for camels
Peng et al. (36)	<i>in vivo</i> animal (pig) female	Animal medical experiment	Assessment of the effect of melatonin on the course of pregnancy	Melatonin increases the expression of antioxidant factors	Melatonin supplementation may have a positive effect on the maintenance of pregnancy
Arend et al. (37)	<i>in vivo</i> animal (swine) female	Animal medical experiment	Evaluation of the effect of melatonin administration on female reproduction	The study group showed a higher percentage of surviving pregnancies	Melatonin may be a fertility enhancer in pigs
Bai et al. (38)	<i>in vivo</i> animal (sheep) female	Animal medical experiment	Assessment of the effect of melatonin on the immune system functioning in pregnant females	Cd4 and mtnr1a expression was higher in samples taken from the immune system in early pregnancy	Melatonin has been linked to changes in the immune system functioning during pregnancy

(Continued)

TABLE 2 Continued

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Gunwant et al. (39)	<i>in vivo</i> animal (water buffalo) female	Animal medical experiment	Determination of the influence of MTNR1A receptor polymorphism on reproductive potential	The reproductive potential varies depending on the MTNR1A gene variant	Melatonin receptor polymorphism affects the occurrence of fertility periods
Fathy et al. (40)	<i>in vivo</i> animal (sheep breeds) female	Animal medical experiment	Assessment of the influence of MTNR1A and AA-NAT polymorphism on reproductive potential	The reproductive potential varies with the polymorphism	The MTNR1A and AA-NAT polymorphism affects the age of the first estrus
Mura et al. (41)	<i>in vivo</i> animal (Sarda ewes) female	Animal medical experiment	Assessment of the influence of MTNR1A polymorphism on reproductive potential	The reproductive potential varies depending on the MTNR1A gene variant	MTNR1A polymorphism affects fertility
do Nascimento Marinho et al. (42)	<i>in vivo</i> animal (rats) female	Animal medical experiment	Determination of the protective effect of melatonin when exposed to pesticides during pregnancy	The genetic changes caused by exposure to cypermethrin were reduced in the study group	Melatonin can be used as a drug to protect against the negative effects of cypermethrin, but it is not effective in reversing the effect of methomyl
Huang et al. (43)	<i>in vivo</i> animal (mouse) female	Animal medical experiment	Determination of the melatonin protective effect on the ovaries during the use of cisplatin	Ovarian toxicity of cisplatin is decreased when melatonin is used during chemotherapy	Melatonin can be used in a protective way during cisplatin administration to preserve fertility
de Almeida (44)	<i>in vivo</i> animal (rats) female	Animal medical experiment	Evaluation of the effect of melatonin on fertility after exposure to herbicides in female rats	The study group showed a higher percentage of pregnancies and fetal survival	Melatonin can be used as protection against herbicide exposure
de Sousa Coelho et al. (45)	<i>in vivo</i> animal (rats) female	Animal medical experiment	Determination of the protective effect of melatonin on the fetus exposed to ethanol	Melatonin reduces the defects caused by exposure of the fetus to ethanol	Melatonin can be used to prevent fas
Wang et al. (46)	<i>in vivo</i> animal (mouse) male	Animal medical experiment	Determination of the melatonin protective effect on the testes during the use of paclitaxel	The sperm quality increases after the use of melatonin during treatment with paclitaxel	Melatonin can be used in a protective way during paclitaxel administration to preserve fertility
El Gheit et al. (47)	<i>in vivo</i> animal (rat) male	Animal medical experiment	Evaluation of the effectiveness of melatonin in the treatment of varicocele	Less histopathological changes in the testicle were found in the melatonin group	Melatonin can be used as an adjunct therapy for varicocele to reduce damage to the testes

Prolactin is a peptide hormone synthesized by lactotrophic cells of the anterior pituitary gland. Its secretion is mainly influenced by dopamine, which negatively influences its secretion through negative feedback. The main function of prolactin is to influence the development of the mammary glands during pregnancy and to support milk production after delivery. Plasma prolactin levels increase rapidly during pregnancy with an increase in the size and number of the anterior pituitary lactotrophic cells. During breastfeeding, the sucking of the nipple by the baby causes a rapid secretion of prolactin through a mechanism of rapid neuroendocrine reflex. In pathological situations, the symptoms of hyperprolactinemia include hypogonadotropic hypogonadism (menstrual disorders and infertility) or visual field disturbances due to the mass effect caused by prolactinoma, the most common pituitary tumor. Hyperprolactinemia is diagnosed by measuring plasma prolactin. A false-positive result may be due to the presence of macroprolactin in the plasma. Its verification is possible with the use of the polyethylene glycol (PEG) precipitation method, which allows to avoid unnecessary treatment (53).

Hypothyroidism leads to a compensatory increase in the production of thyrotropin-releasing hormone (TRH) by the hypothalamus, which in turn results in suppression of dopamine production, resulting in excessive prolactin secretion, i.e. hyperprolactinemia (54). Central nervous system (CNS) tumors that produce prolactin, called prolactinomas, also cause hyperprolactinemia. Hyperprolactinemia is of key importance in infertility as it inhibits the production of GnRH and, consequently, also of gonadotropins (LH, FSH) by the pituitary gland. The therapy most often uses bromocriptine, while the basic diagnostic imaging test is magnetic resonance imaging (MR) in the T1 sequence (19, 20, 55). Prolactinomas are the most common hormonally active tumors in pregnant women. A case of a 30-year-old woman in the 36th week of pregnancy with headaches and left-sided vision loss was described. MRI of the pituitary gland confirmed a 10x11mm suprasellar lesion. After delivery, the lesion was removed by endoscopic transnasal resection. Histopathological examination revealed a tissue of prolactinoma. Therefore, a number of potential causes, including

prolactinoma, should be considered in the differential diagnosis of visual loss during pregnancy (56).

Studies on the neuropeptide kisspeptin encoded by the Kiss1 gene prove that it is a potent inducer of GnRH production. In animal models, mice with hyperprolactinemia were administered kisspeptin once a day for 20 days, which led to the restoration of estrus cycles and an increase in FSH and LH levels. Thus, kisspeptin may be used in the future to treat infertility associated with hyperprolactinemia (54). In studies with mice and sheep, paracrine substances secreted by the uterine glands have been shown to play a key role in implantation of the embryo and maintenance of pregnancy in the early stages. Therefore, problems with getting pregnant and with its maintenance may result from impaired functioning of the uterine glands. A thorough understanding and study of this issue may in the future contribute to improving the outcomes of infertility treatment as well as the maintenance of health by the mother and the fetus (57).

Multi-hormonal pituitary insufficiency is most often idiopathic and is associated with a mutation in one or several genes, among which the most important are transcription factors such as GLI2, LHX3, LHX4, HESX1, PROP1, POU1F1, SOX2, PITX2, OTX2, SOX3. Clinically, it is possible to encounter an isolated deficiency of one of the pituitary gland hormones or with a disturbance in the production of some or even all pituitary hormones, including gonadotrophins. It is therefore another potential cause of infertility (58). Postpartum pituitary hemorrhage, known as Sheehan's syndrome, is a common cause of multi-hormonal pituitary insufficiency in tropical countries. Hypopituitarism also occurs in 10% of survivors of a poisonous snake. In turn, post-traumatic pituitary insufficiency accounts for approximately 7% of all cases (59). Empty sella syndrome can be another cause of hypopituitarism. It may be idiopathic or associated with postpartum hemorrhage, head trauma, CNS stroke, hormonally active pituitary microadenoma, and after radiation therapy or surgery. It manifests itself clinically with headaches, visual disturbances and hormonal insufficiency of the pituitary gland (60).

Autoimmune diseases, especially systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA), as well as the toxicity of disease-modifying drugs, in particular non-steroidal anti-inflammatory drugs (NSAIDs) and glucocorticoids (GCs), also affect infertility. Although recently there has been an improvement, especially thanks to the use of biological drugs that reduce disease activity to a greater extent (61).

Infections in the hypothalamic-pituitary region account for less than 1% of the causes of its damage. The etiological factors can be bacteria, viruses, fungi and parasites. In imaging studies, infection may appear as a tumor in the area of the sella turcica, which raises the suspicion of a proliferative process. Risk factors for infection include meningitis, sinusitis, neurosurgery, and immune system disorders. Infection may develop in glands unchanged or damaged by previous diseases (adenomas, Rathke's cyst, craniopharyngioma). Diagnostics is difficult due to the lack of specific symptoms. Patients may experience blurred vision or headache, fever and leukocytosis. During the acute phase of the disease, but also several months or years after successful antibiotic therapy, a significant proportion of patients develop symptoms of hypothalamic-pituitary insufficiency (62). Hypopituitarism can be caused by inflammation of the pituitary gland and can be primary or secondary. The most common form of primary inflammation is lymphocytic pituitary inflammation, which

is most frequent in pregnant or postpartum women. In turn, immunotherapy, used in the treatment of neoplastic diseases, is important in secondary pituitary inflammation. Autoimmune complications, which can also affect the pituitary gland, are a side effect of such treatment. The most common clinical symptoms are headache, excessive thirst and visual disturbances, while laboratory deficiencies include ACTH, TSH, FSH, LH, GH and hyperprolactinemia. Although biopsy remains the gold standard in the diagnosis of pituitary inflammation, the most important in clinical practice is MRI, which also helps to differentiate pituitary inflammation from pituitary adenomas (63).

3.1.2 *In vivo* animal studies

It is known that stress negatively influences the homeostasis of the circadian rhythm, which results in disturbance of the hypothalamic-pituitary-gonadal axis. In animal models, knock-out of clock genes such as period circadian regulator 1 and 2 (Per1, Per2), brain and muscle Arnt-like protein-1 (Bmal1) led to infertility in mice (64).

The female reproductive system is also affected by toxic substances such as pesticides, heavy metals, diethylstilbestrol, phenols, bisphenols, parabens. They can interfere with receptor binding, steroidogenesis and hormone metabolism. They also have a proven effect on the increased incidence of preterm labor, fetal growth disorders, miscarriages and difficulties in getting pregnant. Animal studies have shown that exposure to pesticides caused damage to oocytes, decreased production of steroid hormones by the ovaries, and lower fertility. Exposure to diethylstilbestrol suppressed gonadotropin production and caused damage to the pituitary gland and uterine tissue (65).

3.1.3 *In vitro* human studies

In vitro studies have shown that cannabinoids influence the functioning of the hypothalamus and pituitary gland through CB-1 receptors. Blocking these receptors results in an increase in the concentration of ACTH, GH, LH, and TSH, which proves that cannabinoids have an inhibitory effect on the activity of the hypothalamic-pituitary axis (66).

3.2 Male

3.2.1 *In vivo* human studies

Gonadotropin-releasing hormone produced by the hypothalamus regulates the release and secretion of gonadotrophins that control the testes functioning, i.e. luteinizing hormone and follicle-stimulating hormone from the anterior pituitary. FSH receptors are located on Sertoli cells and LH receptors on Leydig cells. The effect of gonadotropins on testicular cells determines the proper course of testosterone synthesis, spermatogenesis, and the quality of sperm. The functioning of the male reproductive system is also influenced by hormones such as estradiol (E2) and prolactin. Estradiol, produced both by the testes and by the peripheral conversion of testosterone, is a potent inhibitor of LH and FSH. On the other hand, prolactin, by inhibiting the production of GnRH, reduces the concentration of LH and testosterone, which results in hypogonadism. In response to the generation of free oxygen radicals (ROS) by the male reproductive system cells, the hypothalamic-pituitary-adrenal axis is activated and

cortisol is produced in response to stress. Cortisol, in turn, inhibits the production of gonadotropins, which causes a decrease in the testosterone production by Leydig cells (due to the lowered LH levels). Moreover, low FSH levels reduce the release of androgen-binding protein (ABP) by Sertoli cells, which further decreases circulating testosterone levels. Interestingly, obesity affects not only the hypothalamic-pituitary-adrenal and hypothalamic-pituitary-thyroid axes, but by generating ROS, it affects the increased production of leptin by adipocytes, which, together with insulin, reduces the concentration of triiodothyronine (T3), and thus adversely affects the testes functioning. Leptin is also an inhibitor of GnRH production by the hypothalamus (67). The pre-pubertal rise in leptin levels is responsible for the proper testes development. In turn, an increase in androgen concentration in adolescent boys causes a decrease in leptin concentration (68).

Obesity is not only associated with a higher risk of cardiovascular disease, stroke, type 2 diabetes, but also with disorders in the reproductive system in both women and men. It also leads to hyperinsulinemia, hyperlipidemia, hyperleptinemia and chronic inflammation. In obese men, decreased levels of testosterone, luteinizing hormone and sex hormone-binding globulin (SHBG) are observed. Increased amount of adipocytes negatively affects the quality and quantity of sperm, and as BMI increases, total sperm count decreases. A meta-analysis of 21 studies involving 13,000 men found a higher incidence of oligozoospermia and azoospermia in obese men. Moreover, the effectiveness of the IVF procedure decreases when the partner is obese (21).

In addition, studies in rats have shown that sleep disorders reduce testosterone levels, decrease sperm motility and apoptosis of Leydig cells in the testes. Moreover, it had a negative effect on the initiation of sexual behavior in males and on the percentage of ejaculation (69).

Infection with the severe acute respiratory syndrome coronavirus 2 (SARS-CoV2) is the cause of a cytokine storm that damages many organs. In addition to the well-known effects of SARS-CoV2 infection on the lungs, heart, and CNS, testicular damage is also under discussion. Evidence for the presence of SARS-CoV2 in sperm samples and in testicular tissue is limited, while an impaired immune system response, hyperthermia and oxidative stress caused by infection, damage testicular tissue. Testicular steroidogenesis defects related to SARS-CoV2 infection are the cause of decreased testosterone concentration, which leads to impaired spermatogenesis, impotence and, consequently, infertility in a significant percentage of convalescents. Increased levels of FSH and LH are biomarkers of testicular damage. Hypothalamic-pituitary-gonadal dysregulation due to SARS-CoV2 infection is also associated with the risk of acute renal failure and cirrhosis (70). It should also be noted that HIV infection is associated with an increased risk of pituitary gland stroke as well as pituitary lymphoma and may therefore lead to hypopituitarism, one of the effects of which is infertility (71).

4 Pineal gland

The pineal gland is a relatively small gland located within the cranial cavity behind the third ventricle that connects to the cerebrum by a peduncle. It is a highly vascularized part of the brain not covered by the blood-brain barrier. 90-95% of the cells of this gland are

pinacocytes, whose main function is the production of melatonin (N-acetyl-5-methoxytryptamine) (72). Its synthesis is regulated by the superior cervical ganglion, which receives information about the intensity of light through the suprachiasmatic nuclei of the hypothalamus, considered to be the anatomical center of the biological clock (73). The pineal gland, through melatonin, is responsible for controlling the regulation of the circadian rhythm, mainly sleep and wakefulness. Its synthesis and secretion significantly increases in the dark (74), and exposure to a light source causes a significant reduction in its synthesis (75). Melatonin acts through the MT1 and MT2 receptors located in the suprachiasmatic nucleus (SCN), reducing the activity of its neurons, which ultimately leads to the feeling of drowsiness and falling asleep (76). Another important function of melatonin is the inhibition of sexual maturation. In humans, the decline in the average daily melatonin production progresses with age and is associated with the progression on the Tanner scale (77, 78). This is due to the inhibition of kisspeptin expression by melatonin, which stimulates the hypothalamus cells to synthesize GnRH (79, 80). It has been proven that the administration of exogenous melatonin causes delayed puberty in children of both sexes (81). Melatonin is also produced in other organs, including the reproductive system (82–87), the retina and lens (88), the gastrointestinal tract (89, 90) and blood cells (91), acting as an autocrine or paracrine hormone (90). However, it most likely does not significantly affect the plasma concentration of melatonin as it was found to be undetectable in rats after pituitary removal (92). Melatonin is also considered to be a very effective antioxidant (73). By activating its MT1 and MT2 receptors, it stimulates the expression of superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and glutathione reductase (GRd) (93). Another mechanism of antioxidant activity is the stimulation of the Nrf2 expression, which is a transcription factor for many genes encoding antioxidant enzymes (94). There are also many reports on the direct antioxidant activity of its metabolites: N1-acetyl-N2-formyl-5-methoxykynuramine (AFMK), N1-acetyl-5-methoxykynuramine (AMK) and cyclic 3-hydroxymelatonin (3-OHM) (95).

4.1 Female

4.1.1 *In vivo* human studies

Melatonin appears to be an important hormone related to female fertility. Its higher plasma levels have been found in women with infertility caused by hypothalamus dysfunction (96, 97). At the same time, it has been shown that melatonin supplementation can positively affect fertility and improve the effectiveness of assisted reproductive protocols (25), which may be related to its influence on the increase of melatonin concentration in the follicular fluid (25, 98). Due to its antioxidant effect, melatonin has a positive effect on the quality of oocytes. Negative effects on fertility may be related to the inhibition of GnRH expression, which promotes ovulation by stimulating LH expression. Melatonin supplementation has also been shown to be beneficial in women suffering from polycystic ovary syndrome (26). This is due to a decrease in plasma testosterone concentration and elevated probability of ovulation (99). Studies carried out in a group of transgender men showed that testosterone inhibited the expression of gonadotropic hormones, which

contributed to a strong suppression of ovulation (100). The reasons for the positive effect on the fertility of melatonin supplementation in PCOS patients are likely to be found in the antioxidant potential decrease (101).

Melatonin is associated not only with fertility, but also with the course of pregnancy. An increase in melatonin concentration during pregnancy and its sudden decrease after delivery have been shown (102). Furthermore, an increased risk of cholestasis has been demonstrated in women with a lower melatonin concentration (27) and the effectiveness of melatonin administration in lowering blood pressure and extending the duration of pregnancy in the course of pre-eclampsia has been confirmed (28). Studies are still needed to establish the relationship between melatonin levels and the course of pregnancy in more detail.

4.1.2 *In vivo* animal studies

In animals, a strong relationship between melatonin and fertility has also been demonstrated. Melatonin deficiency is correlated with decreased fertility (29, 30), lower levels of sex steroid hormones (30), and decreased survival of embryos after fertilization (31). At the same time, a positive correlation was observed between serum melatonin concentration and fertility (32–36). Administration of exogenous melatonin is associated with a higher percentage of maintained pregnancies (37). Melatonin also participates in the activity changes of the female immune system during pregnancy in sheep, influencing the immune activity of the thymus, lymph nodes and spleen, which are involved in the process of fetal immune tolerance. The effect of melatonin on the immune system is complex and pleiotropic. At the beginning of the inflammatory reaction, it is immunostimulating, later it changes to immunosuppressive. The changes induced by melatonin in pregnancy are mainly related to the stimulation of the helper lymphocyte population by directly affecting the MT1 and MT2 receptors (38).

The relationship between melatonin receptors polymorphism and female fertility has been proven (32, 39–41). There are also numerous reports suggesting a protective effect of melatonin administration on fertility, maintenance of pregnancy and prevention of birth defects in females exposed to cytostatics, insecticides, herbicides and alcohol (42–46), which is directly related to its antioxidant nature.

4.1.3 *In vitro* human studies

In vitro studies focus mainly on the antioxidant activity of melatonin and its relationship with the production of gonads. In females, a positive correlation was demonstrated between the melatonin concentration in the follicular fluid and the markers of the ovarian reserve and the quality of oocytes (103–105).

4.1.4 *In vivo* animal studies

Similarly in animal studies, the melatonin level in the follicular fluid correlated with the quality of oocytes and ovarian reserve (106, 107). In addition, melatonin seems to be a good component of cryopreserves used in the storage of the oocytes (108–110). The induction of estradiol production by granular cells was also confirmed (106).

4.2 Male

4.2.1 *In vivo* human studies

In men, melatonin is also very important in terms of fertility. Similarly to women, melatonin causes a decrease in LH secretion, leading to lower testosterone secretion by Leydig cells in the testes, which has a negative impact on the intensity of spermatogenesis (107). A positive correlation was demonstrated between the concentration of melatonin and male fertility, both in the serum and in semen. Interestingly, an increased concentration of melatonin in the plasma and its decreased level in semen was demonstrated in infertile men (111). Studies have shown that the concentration of melatonin in semen inversely correlates with the frequency of sperm DNA damage, which is probably due to its antioxidant activity (112). It has also been reported that melatonin supplementation improves sperm quality (113, 114) and increases the chance of preserving fertility in patients with varicocele (115).

4.2.2 *In vivo* animal studies

There are also studies linking melatonin to male fertility in animals (46). Protective effect on gonads has been demonstrated in the course of varicocele, radiotherapy and metabolic syndrome, exposure to cytostatics and heavy metals (47, 116–119). In addition, the synthesis of melatonin within the testes has been confirmed (120) and the positive effect of its concentration on semen quality was demonstrated (121).

4.2.3 *In vitro* animal studies

The importance of the antioxidant activity of melatonin during spermatogenesis has also been shown in animal studies (122). Furthermore, melatonin seems to be a good component of cryopreserves used in the storage of sperm samples (123).

5 Thyroid

5.1 Female

5.1.1 *In vivo* human studies

Inhibition of the secretion of thyroid hormones is associated with its hypothyroidism. In milder hypothyroidism, infertility is usually not achieved, but the risk of spontaneous miscarriage, premature births and stillbirths increases (124–126) (Table 3). Severe hypothyroidism may lead to infertility as a consequence of a direct inhibitory effect on the ovarian ovulatory activity as well as through an effect on the pituitary-ovarian axis. Decreased activity of sex hormone-binding globulin causes an increase in serum free testosterone and estradiol, moreover, metabolic clearance of androstenedione and estrone is decreased. The elevated level of thyrotropin-releasing hormone in primary hypothyroidism is responsible for the increase in serum prolactin levels and the delayed luteinizing hormone response to the stimulating effect of gonadotropin-releasing hormone, which in turn leads to luteal phase failure in women (124–126, 131, 132).

TABLE 3 Thyroid-related fertility studies.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Vaquero et al. (126)	<i>in vivo</i> human female	Clinical trial	Evaluation of the role of benign thyroid abnormalities in recurrent spontaneous abortion and evaluation of the effects of two different therapeutic protocols	Treatment of mild thyroid disorders with immunoglobulins resulted in termination of pregnancy in 54.5% of cases, while the use of replacement therapy in 81.2% of cases	Thyroid replacement therapy is more effective in the obstetric context than intravenous immunoglobulin
Kakita-Kobayashi et al. (127)	<i>in vitro</i> human female	Clinical experiment	Evaluation of the effect of thyroid hormone on decidualization in human endometrial stromal cells (hescs) and determination of its physiological roles <i>in vitro</i> through gene targeting	A significant increase in decidual response was observed after combined treatment with ovarian steroid hormones and thyroid hormone	Deciduality impairment is a possible cause of infertility in patients with subclinical hypothyroidism (SCH)
Barber et al. (128)	<i>in vitro</i> human female	Clinical experiment	Determination of the expression of thyroid hormone receptors in the extranodal trophoblast, elucidation of the effect of T3 on both the proliferation and differentiation of human trophoblast cells of various origins, and the determination of a potential interaction between EGF and T3	T3 and EGF have an anti-proliferative effect on cells of the extranodal-like lineage (SGHPL-4), with a concomitant proliferation-promoting effect on JEG-3 choriocarcinoma cells	EGF and T3 have a synergistic effect in regulating human trophoblast proliferation and differentiation.
Giuliani et al. (129)	<i>in vitro</i> animal (rat) female	Clinical experiment	Evaluation of the effect of quercetin on the expression of genes encoding thyroid hormones and radioiodine uptake	Quercetin reduces the expression of the thyrotropin receptor, thyroid peroxidase and thyroglobulin genes, and also reduces the uptake of radioiodine by thyroid cells	Quercetin may act as a thyroid disruptor, supplementation should be used with caution
Condorelli et al. (130)	<i>in vitro</i> human male	Clinical experiment	Assessment the <i>in vitro</i> effects of levothyroxine (LT4) on conventional and biofunctional sperm parameters and its implications on fertility	Thyroid hormones have a beneficial effect on sperm mitochondrial function, oxidative stress and DNA integrity	Restoring the normal secretion of thyroid hormones is important in idiopathic male infertility

5.1.2 *In vitro* human studies

Due to the small number of *in vitro* studies and their contradiction regarding the influence of the role of thyroid hormones on reproductive function and semen parameters in humans, it is necessary to exercise caution when interpreting them. Kakita-Kobayashi et al. determined the effect of levothyroxine on decidualization in human endometrial stromal cells (hESCs) and the selection of key genes for this process *in vitro*. After exposure of cells to the drug, a significant increase in the temporal response was observed. Moreover, treatment with LT4 also influenced the regulation of many transcription factors important for decidualization. There was also an increase in type 3 deiodinase in the presence of thyroid hormones, which is an important element in the tissues of the fetus and placenta. In addition, it has been noted that the progesterone receptor and the ovarian steroid hormone receptor are involved in thyroid hormone-induced decidualization. Based on the conducted study, Kakita-Kobayashi et al. claim that impaired decidualization is a possible cause of infertility in patients with subliminal hypothyroidism (127).

It was shown that the trophoblast contains triiodothyronine receptors. *In vitro* studies indicate that thyroid hormones have a direct impact on the early development of the placenta by stimulating angiogenesis and by promoting the invasion and differentiation of embryonic cells (133). Thyroid receptors are found in the endometrium, and their highest levels are observed in the receptive endometrium (134). Barber et al., using immunohistochemical staining, localized specific isoforms of thyroid hormone receptors in

extracellular trophoblasts in placenta biopsies in the first and second trimesters of pregnancy, which indicates a potential sensitivity of cells to T3. Epidermal growth factor (EGF) and T3 exhibited antiproliferative effects on the extravillous trophoblast cell line (SGHPL-4) while promoting proliferation in JEG-3 chorionic cancer cells. The study suggests that T3 and EGF may act synergistically, regulating both the proliferation and the differential function of human trophoblast (128).

5.1.3 *In vitro* animal studies

In 2008, quercetin was shown to inhibit the dose- and time-dependent spread of Fischer rat thyroid cell line (FRTL-5) by inhibiting the insulin-regulated action of Akt kinase. Quercetin interferes with TSH-dependent NIS gene expression and transport in FRTL-5 cells. These observations may help to understand the molecular mechanism of quercetin's anti-thyroid effects on cell growth and function. Even when taken from an *in vitro* thyroid cell line that does not have the characteristics of a transformed cell, these results led to the evaluation of quercetin as an anti-thyroid drug in hyperthyroidism (135, 136). In recent studies, quercetin appears to reduce the expression of the thyroid-stimulating hormone receptor, thyroid peroxidase (TPO) and thyroglobulin (Tg) genes. The anti-thyroid effect of quercetin was further assessed *in vivo*. It was administered (50 mg/kg) to a Sprague-Dawley rat, and after 14 days of treatment, the radioiodine uptake decreased significantly, indicating that quercetin may act as a thyroid disruptor (129, 135). Soybean extracts inhibit iodine uptake and increase the protein

content of a known autoimmune Tg fragment in rat Fischer thyroid cells. These effects may be responsible for the link between more frequent soy consumption and thyroid disorders such as hypothyroidism, goiter, and autoimmune thyroid disease (135, 137). Among the flavonoids, epigallocatechin-3 gallate (EGCG), a catechin rich in green tea, when administered to male rats at doses of 25, 50 and 100 mg/kg of body weight, showed an antithyroid activity, which was manifested by reduced activity of thyroid peroxidase and 5'-deiodinase I and increased activity of the thyroid sodium-potassium pump. In addition, serum T3 and T4 levels were lowered while serum TSH was elevated in rats, demonstrating the possibility of goiter *in vivo* (135, 138).

5.2 Male

5.2.1 *In vitro* human studies

Condorelli et al. assessed the effects of levothyroxine (LT4) on conventional and biofunctional sperm parameters and their impact on fertility. The authors obtained sperm from men with confirmed infertility and exposed them to LT4 *in vitro*. The results confirmed the effect of LT4 on an increase in the number of sperm with a high level of mitochondrial membrane potential (MMP), a reduction in the percentage of sperm with a low MMP and an increase in sperm motility. LT4 was found to induce sperm proliferation and lipid peroxidation, thereby improving the chromatin tightness in the cell nucleus. According to the researchers, the results of the above-mentioned *in vitro* study may have clinical application in patients with idiopathic infertility due to the explanation of the influence of thyroid function on fertility in men (130).

So far, a key role of vitamin D in the male reproductive system has been suggested as its receptors and metabolizing enzymes have been shown to be expressed in the testes and sperm. Vitamin D metabolism is mainly regulated by parathyroid hormone (PTH), produced by the parathyroid glands, and fibroblast growth factor 23 (FGF23), synthesized by osteoblasts and osteoclasts. Decreased levels of circulating calcium and 25-hydroxyvitamin D3 increase PTH secretion, which stimulates 1- α -hydroxylase and inhibits the expression of 24-hydroxylase in the kidney, leading to higher levels of vitamin D and calcium (139, 140). In addition, elevated levels of phosphorus and 25-hydroxyvitamin D3 inhibit 1- α -hydroxylase and stimulate 24-hydroxylase, resulting in a reduction in vitamin D. To close the feedback loop when vitamin D and phosphorus levels decline, FGF23 is inhibited, leading to an increase in vitamin D levels (139, 141). The vitamin D receptor (VDR) and the enzymes that metabolize vitamin D are simultaneously expressed in Sertoli cells, germ cells, Leydig cells, sperm, and cells in the epithelial lining of the male reproductive system. The presence of vitamin D metabolizing enzymes suggests that the reproductive organs may modulate the local response to vitamin D in animals and humans. Nuclear somatic or embryonic cells appear to be able to synthesize and degrade vitamin D locally, independent of systemic vitamin D metabolism. Moreover, expression of VDR in the testes suggests that vitamin D may exert autocrine and paracrine effects, possibly playing a role in regulating testicular function, thus contributing to male infertility. The expression of VDR and vitamin D metabolizing enzymes in the

male reproductive system has been extensively analyzed in animal and human studies. The VDR protein has been found in the prostate, seminal vesicles, epididymis, and also in germ cells, especially spermatogonia, spermatocytes, and Sertoli cells (139, 142). VDR protein expression has been found in animal sperm but has been suppressed in the tail of the epididymis (139, 143). In the same context, testosterone synthesis enzymes in the testes appeared to be reduced in mice fed the vitamin D deficiency (VDD) diet (144).

6 Thymus

Changes in the immune system activity are necessary in the course of a healthy pregnancy as they help to avoid an immune response against an allogeneic fetus (145) (Table 4). One of the organs whose activity change is required for the development of immune tolerance during pregnancy is the thymus, in which the maturation and differentiation of helper T cells take place.

6.1 Female

6.1.1 *In vivo* human studies

Maternal Th, Tc and NK cells express activation markers on their surface after blastocyst implantation in the uterine endometrium, which suggests rapid recognition of trophoblast by these cell populations (146). Many studies have shown changes in the population of T lymphocytes and their activity in normal pregnancy (147, 155–160). In addition, human placenta produces IL-35 with an immunosuppressive effect, stimulating the proliferation of regulatory T cells in the thymus (161, 162).

The activity of the thymus is also important in terms of the etiology of primary ovarian insufficiency (POI), where one of the causes may be an autoimmune reaction. An increased risk of POI in patients with autoimmune diseases and an increased risk of autoimmune diseases in patients with POI have been observed (163).

6.1.2 *In vivo* animal studies

In animal studies, changes in the population of T lymphocytes mainly include an increase in the Th cell population during early pregnancy (149), a greater percentage of type 2 helper cells in relation to type 1 helper cells (148, 164), changes in the toll-like receptor (TLR) expression within thymic epithelial cells, and the synthesis of prostaglandins and proteins stimulated by interferon (150, 151). These changes are directly responsible for the phenomenon of fetal immunotolerance, inhibiting the development and activation of a subpopulation of T lymphocytes that can stimulate inflammatory reactions within the uterine endometrium, while maintaining the endometrial and trophoblast defense against pathogens (165). This is possible due to the activity of the placenta, which produces Th2-specific cytokines such as IL-4, IL-5 and IL-10 throughout the duration of pregnancy (164).

During pregnancy, there is both local (within the uterus) and general increase in the population of regulatory T cells specific for paternal antigens (166, 167), which is responsible for the inhibition of the immune response to them.

TABLE 4 Thymus-related fertility studies.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Saito et al. (146)	<i>in vivo</i> human female	Comparative study	Assessment of activation antigens on T cells of human decidua at an early stage of pregnancy	The decidua contained a small number of T cells and both CD4+ and CD8+ subsets expressed CD69, HLA-DR, IL-2R alpha and IL-2R beta antigens	T cells in the decidua in the first trimester of pregnancy are regionally activated
Watanabe et al. (147)	<i>in vivo</i> human female	Comparative study	Assessment of changes in T, B, and NK lymphocyte subsets during and after pregnancy	Suppressor T and NK+3 cells increase in early pregnancy and decrease in late pregnancy	The number of T, B, and NK lymphocyte subsets changes throughout pregnancy
Li et al. (148)	<i>in vivo</i> animal (sheep) female	Animal medical experiment	Assessment of TLR expression change in early pregnancy in females	Changes in TLR expression have been demonstrated in early pregnancy	Alteration of TLR expression may be involved in the generation of immune tolerance
Zhang et al. (149)	<i>in vivo</i> animal (sheep) female	Animal medical experiment (sheep)	Assessment of the influence of early pregnancy hormones on thymic cytokine expression in females	Increased expression of cytokines stimulating the production of th1 and th2 lymphocytes	In early pregnancy, the production of helper t cells that may be involved in the production of immune tolerance is stimulated
Yang et al. (150)	<i>in vivo</i> animal (sheep) female	Animal medical experiment	Assessment of the influence of early pregnancy hormones on prostaglandin synthesis in the female thymus	A large increase in thymic prostaglandin synthesis has been demonstrated	In early pregnancy, the expression of prostaglandin synthases in the thymus changes, which may be associated with the development of immune tolerance
Zhang et al. (151)	<i>in vivo</i> animal (sheep) female	Animal medical experiment	Assessment of the interferon stimulated change in protein expression during early pregnancy in females	Changes in the expression of interferon-stimulated genes have been demonstrated in early pregnancy	The altered expression of certain proteins may be associated with the development of immune tolerance during pregnancy
Wise (152)	<i>in vivo</i> animal (rat) male	Animal medical experiment	Assessment of the effects of thymulin on rat testicular steroid synthesis	Increased doses of thymulin (100 ng/ml) resulted in decreased testicular steroid synthesis	Increased concentrations of thymulin had inhibitory effects on testicular steroidogenesis
Wise and Ford (153)	<i>in vivo</i> animal male	Animal medical experiment (boar)	Assessment of the effects of thymic peptide thymulin on testicular steroid synthesis	Injection of thymulin increased circulating testosterone concentrations	Thymulin increases androgen stimulation in boar testes
Jacobo et al. (154)	<i>in vivo</i> animal (rat) male	Animal medical experiment	Analysis the phenotype and number of T lymphocytes in the testicular interstitium of rats during EAO development	CD4+Foxp3+ T(reg) cells were more abundant than CD8+Foxp3+ T(reg) cells	The numbers of T(reg) cell subsets increased in the testis of rats with orchitis

The activity of thymus is related not only to pregnancy, but also to the maturation of the ovaries and their activity. Studies in mice have shown that early removal of the thymus causes disorders in the ovarian development, failure of maturation and sterility. This effect was reversible by administering thymulin, although a decrease in fertility was observed due to the presence of autoreactive T cells (168, 169). It is related to the influence of thymulin on the expression of LH and FSH by the pituitary gland. It has been demonstrated that GnRH and thymulin act synergistically with the release of LH and additively with the FSH secretion (170). The variable effect of thymulin on LH and FSH expression depending on the time of the menstrual cycle has also been reported (171).

6.2 Male

6.2.1 *In vivo* animal studies

A positive correlation was demonstrated between the concentration of thymulin and LH, as well as a reduction in

testosterone production with a decrease in thymulin concentration (152, 153). The presence of regulatory T cells within the testes has also been demonstrated (154, 172), which, through the secretion of IL-10, affect the ratio of the Th1 to Th2 subpopulation. Their presence is essential for the immunotolerance of spermatocyte antigens. In the case of disorders leading to a reduction in their population size, autoimmune orchitis may occur (154).

7 Pancreas

Disorders of lipid and glucose metabolism, recently associated with the increasingly frequent obesity, are a current medical problem. They not only pose a health risk, but also have an adverse effect on reproductive function (Table 5). Insulin resistance, i.e. reduced cell sensitivity to insulin and compensatory hyperinsulinemia, induce an early response to luteinizing hormones and cause premature differentiation of small follicles, resulting in anovulation. On the other hand, they adversely affects the functions and environment of

the endometrium and are responsible for disturbances in embryo implantation (180). In addition, hyperinsulinemia disturbs the intra-follicular microenvironment during folliculogenesis, decreases the fertilization rate and the potential for embryonic development during natural ovarian stimulation cycles (173). Insulin primarily acts on its own receptors located on the theca cells surrounding the stromal and granulosa cells, thereby stimulating ovarian steroidogenesis (174). Research shows that inflammation (elevated levels of IL-6 and IL-17) accompanying insulin resistance and obesity affect ovulation and fertilization and increase the risk of early miscarriage. Moreover, obese people show increased levels of estrogens due to the overexpression of aromatase in adipose tissue, which in turn disturbs ovulation (173, 175).

7.1 Female

7.1.1 *In vivo* human studies

Glucagon is a central mediator of glycemic control, released by pancreatic alpha cells in response to hypoglycemia. It is involved in insulin signaling and action, but also predisposes to adverse pregnancy outcomes. Interestingly, more and more reports confirm

the modulating activity of glucagon-like peptide-1 (GLP-1) in reproduction. GLP-1 receptor antagonists play an important role in the treatment of type 2 diabetes and obesity by lowering glucose levels, reducing body weight and improving reproductive health. Obesity inhibits the hypothalamic-pituitary-gonadal axis, disrupting ovarian function, ovulation index and endometrial receptivity, and affects the molecular mechanisms that regulate the biological activity of the reproductive system (3, 181, 182).

Pancreatitis accompanying pregnancy is not common, but it carries a burden in terms of maternal mortality or perinatal mortality (183). Commonly used in hormone contraception, hormone replacement therapy (after menopause) or hormone therapy in *in vitro* fertilization (development of the endometrium in preparation for embryo transfer), estrogens may be one of the causes of acute drug-induced pancreatitis (184). These cases are rare and the clinical course is usually mild to moderate (185). The mechanism of acute estrogen-induced pancreatitis is not fully understood, but it is presumed to be due to hypertriglyceridemia. Mechanisms related to this phenomenon include an increase in the synthesis of TG in the liver, secretion of VLDL into the circulation, an increase in newly synthesized TG in the liver, excessive secretion of TG and apolipoprotein, and the effect of estrogens on the inhibition of lipoprotein lipase (LpL) promoter activity

TABLE 5 Pancreatic-related fertility studies.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Wang et al. (173)	<i>in vivo</i> human female	Prospective cohort study	Evaluation of the impact of insulin resistance on IVF outcomes in women without PCOS	The percentage of mature oocytes and the rate of blastocyst formation were significantly lower in the IR group compared to the group without IR	In lean infertile women without PCOS, insulin resistance is associated with a reduced percentage of mature eggs and poor embryo quality, in which pancreatic B-cell immunity may play a role
Mekaru et al. (174)	<i>in vivo</i> human female	Retrospective study	Evaluation of whether insulin resistance in patients without PCOS affects the results of <i>in vitro</i> fertilization and pregnancy	There were no differences in response to controlled ovarian hyperstimulation, number of oocytes retrieved, conception rate, pregnancy rate, live birth rate, and incidence of gestational diabetes in women with and without insulin resistance	Insulin resistance in patients without PCOS has no impact on IVF outcomes or perinatal prognosis
Li et al. (175)	<i>in vivo</i> human female	Retrospective study	Assessment of the importance of central obesity on the effectiveness of IVF	Women with central obesity had significantly more endocrine and metabolic disorders and required significantly higher doses of gonadotropins, longer duration of ovarian stimulation	Central obesity negatively affects the effectiveness of IVF
Issa et al. (176)	<i>in vivo</i> human female	Case study	Case presentation	Severe hypertriglyceridemia and secondary acute pancreatitis and diabetic ketoacidosis have occurred following <i>in vitro</i> fertilization	IVF may be associated with hypertriglyceridemia with secondary acute pancreatitis
Vuguin et al. (177)	<i>in vivo</i> animal (mouse) female	Animal medical experiment	Assessment of the importance of glucagon in the regulation of fetal growth and maturation	Deletion of the GLU receptor negatively affected fetal survival and in adult animals resulted in a change in the β/α cell ratio	Glucagon plays an important role in embryogenesis
Saleh et al. (178)	<i>in vivo</i> human male	Comparative study	Assessment of the importance of insulin resistance and irisin concentration in the etiology of male idiopathic infertility	Infertile patients had higher HOMA-IR and lower irisin compared to controls	Insulin resistance is one of the potential factors of idiopathic male infertility
Izzi-Engbeaya et al. (179)	<i>in vivo</i> human male	Randomized controlled trial	Evaluation of the effect of glucagon administration on the secretion of sex hormones in healthy young men	There were no significant differences in circulating LH, FSH or testosterone levels, although glucagon administration had a metabolic effect	Intravenous administration of glucagon does not affect the secretion of sex hormones in healthy men

(176). Transported by chylomicrons and very low-density lipoproteins, triglycerides are hydrolyzed by high lipase concentrations in the pancreatic capillaries, where they form a large amount of toxic free fatty acids that cause lipotoxicity in acute pancreatitis (186). Hence, in the case of patients qualified for IVF, especially those from the high-risk group (such as diabetes, polycystic ovary syndrome, obesity, dyslipidemia), it is reasonable to screen for lipid abnormalities before starting the *in vitro* procedure and constantly monitor key biochemical parameters (176).

7.1.2 *In vivo* animal studies

Studies in a mouse model (glucagon receptor knockout mice) showed that the lack of glucagon signaling did not alter the hypothalamic-pituitary-ovarian axis. Pregnant knockout female mice exhibited hypoglycemia and hyperglucagonemia accompanied by decreased fetal weight, increased late-stage fetal mortality, and placental abnormalities. Moreover, the lack of glucagon signaling significantly reduced the level of expression of genes controlling growth, adrenergic signaling, vascularization, oxidative stress and the activity of G-protein-coupled receptors (187). Studies on an animal model allow the conclusion that, similar to insulin, glucagon contributes to the proper reproductive function of women. It has also been shown that the lack of glucagon signaling creates a poor quality uterine environment for fetal growth. Animal studies have demonstrated that the lack of glucagon signaling during pregnancy was associated with reduced litter size, limited intrauterine growth, and increased neonatal mortality (177).

7.2 Male

7.2.1 *In vivo* human studies

In a study of male patients with idiopathic infertility, the relationship of metabolic syndrome, obesity and diabetes mellitus was shown as factors contributing to the pathogenesis of male infertility. A strong association of insulin resistance has also been shown in patients with unexplained fertility problems, presuming that elevated blood insulin levels may impair spermatogenesis, and in patients with hyperinsulinemia and type 2 diabetes, also generate nuclear and mitochondrial DNA damage in sperm (178, 188). In the aspect of metabolic diseases, the relationship between sperm count and BMI was also investigated and it was shown that obese men had a twofold increase in the risk of oligozoospermia and increased scrotal temperature, leading to sperm dysfunction, reduced sperm count and mobility, and DNA damage (188).

Moreover, it has been confirmed that glucagon can directly stimulate the reproductive axis, which has not been observed in studies on men where there was no effect of glucagon administration on reproductive hormone levels (179). On the other hand, administration of glucagon receptor antagonists can cause metabolic effects (weight loss, i.e. alleviation of hypogonadism of the hypothalamus in obese men) without direct negative effects on the reproductive system (179).

8 Adrenal glands

Changes in the hypothalamic-pituitary-adrenal axis and the subsequent alterations in the concentration of circulating hormones

constitute the body's response to stressful challenges (Table 6). Mobilization of resources during the stress response suppresses the reproductive axis, which gives the survival of an individual higher priority than the preservation of the species (195). Stressors also affect the adrenal medulla, which secretes catecholamines, i.e. adrenaline and noradrenaline, and the adrenal cortex that secretes aldosterone, which is, next to cortisol, a key regulator of blood pressure (196).

8.1 Female

8.1.1 *In vivo* human studies

Past studies have shown that patients with lower levels of adrenaline had a higher rate of implantation (the group of patients undergoing IVF), which suggests that the less stressful life they lead, the greater their reproductive success (189, 190). It has been found that stressors activate the hypothalamic-pituitary-adrenal axis, causing changes that have a significant impact on female fertility (197).

The IVF procedure itself may be a stressful condition. In a study by Smeenk et al. there was no difference in the stress response between women who became pregnant after treatment and those who did not. Little is known about the effects of catecholamines and cortisol on the physiological processes related to reproduction. It is concluded that catecholamines may affect fertility by altering uterine blood flow, while cortisol with immunosuppressive properties may additionally affect the immune states necessary for implantation. Studies have shown that a higher ratio of serum cortisol to follicular cortisol was associated with pregnancy, and that infertile women had higher levels of stress in terms of circulating prolactin and cortisol compared to fertile women in the control group (190).

Aldosterone, as a mineralocorticoid hormone secreted by the zona glomerulosa of the adrenal cortex, is responsible for the regulation of water and electrolyte balance and blood pressure. During physiological pregnancy, aldosterone levels increase, inducing elevated plasma volume, which is essential for the maintenance of circulating blood volume, blood pressure, and uteroplacental perfusion. Aldosterone levels remain high throughout pregnancy, suggesting a possible role in the regulation of placental and fetal development (198, 199). Recent studies have shown the possible involvement of aldosterone in some gynecological conditions and diseases, including endometriosis combined with infertility. Aldosterone may exacerbate systemic and local inflammatory states underlying endometriosis by activating mineralocorticoid receptors present in inflammatory cells (198).

8.1.2 *In vivo* animal studies

High levels of the norepinephrine transporter (NET) have been documented in mammalian placenta tissue. Placental changes in NET were involved in the modification of reproductive function and fertility of the offspring caused by pregnancy stress (191). Studies in rats showed that exposure of pregnant rats to sympathetic stress affected the placental transport of norepinephrine, leading to a reduction in the ability of the placenta to remove norepinephrine from the fetus into the maternal circulation. It also resulted in impaired fertility of the offspring in adulthood (191). In addition, it

TABLE 6 Adrenal-related fertility studies.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Csemiczky et al. (189)	<i>in vivo</i> human female	Comparative study	Assessment of prolactin and cortisol levels as well as assessment of the personality profile of infertile women	Infertile women also had significantly higher levels of prolactin and cortisol throughout their menstrual cycle	Infertile women have higher serum cortisol and anxiety levels
Smeenk et al. (190)	<i>in vivo</i> human female	Multicenter study	Evaluation of the relationships between the concentrations of stress hormones in the urine: adrenaline, noradrenaline and cortisol in women entering IVF for the first time	A significantly higher concentration of the assessed hormones was found before IVF, which decreased in the case of IVF success	Cortisol is the missing link in the relationship between psychosocial stress and outcome after IVF/ICSI
Piquer et al. (191)	<i>in vivo</i> animal (rat) female	Animal medical experiment	Determination of changes in the expression of the placental norepinephrine transporter during pregnancy and their relationship with its ability to transport norepinephrine under stress conditions	Exposure of pregnant rats to sympathetic stress resulted in increased levels of norepinephrine and corticosterone throughout pregnancy, decreased placental capacity to clear fetal norepinephrine into the maternal circulation, altered levels of placental epinephrine transporter protein depending on fetal sex, and increased placental and offspring body weight	Increased placental adrenaline transporter levels in pregnancy have been associated with decreased adult fertility of offspring
Kapoor et al. (192)	<i>in vivo</i> animal (guinea pigs) female	Animal medical experiment	Assessment of the impact of prenatal stress on the function of the hypothalamic-pituitary-adrenal (HPA) axis in adult offspring	The concentration of cortisol in the offspring of mothers exposed to stress was significantly higher than in the control	Stress exerted on the pregnant female during neuroendocrine development programs growth, HPA axis function and stress-related behavior in adult male guinea pigs
Mayerhofer et al. (193)	<i>in vivo</i> animal (hamster) male	Animal medical experiment	Evaluation of the effect of catecholamines on androgen production during periods of gonadal activity and rest in a seasonally reproducing species	Catecholamines have been shown to modulate the Leydig cell response to gonadotropins in this species of hamster	Stress hormones have a negative impact on reproductive functions and abilities
Rehman et al. (194)	<i>in vivo</i> human male	Retrospective study	Comparing the concentration of stress markers and antioxidants in fertile and infertile men and examining their impact on reproductive hormones and fertility	Cortisol, epinephrine, follicle-stimulating hormone and luteinizing superoxide dismutase, glutathione levels were significantly higher in the group of patients compared to the control group	Stress together with the reduction of antioxidant concentration plays an important role in reducing the reproductive potential in infertile men

has been observed that increased plasma levels of norepinephrine can result in the narrowing of the uterine arteries leading to failure of trophoblast invasion.

Pregnancy is particularly sensitive to stress factors that may lead to permanent modifications in the postpartum development of newborns, predisposing them to diseases that appear in adulthood (200, 201). In a guinea pig study, maternal access to nutrients was drastically restricted during the period of maximum fetal brain growth (70% of pregnancy), resulting in adult male offspring with elevated basal cortisol levels but normal adrenal stress response. However, the same stress applied to 90% of pregnancy results in male offspring with normal basal cortisol levels but increased responsiveness of the hypothalamic-pituitary-adrenal axis to challenge (192). Interspecies differences in the effects of prenatal stress on adult behavior may also be due to the unique development profiles of fetal bodies and brains. In humans, the rapid phase of fetal brain development occurs from about 27 to 30 weeks of gestation and extends to the postpartum period, while in rats or mice, maximum brain growth does not begin until postnatal life. Hence, the period of maternal stress in a rodent would probably

correspond to a completely different phase in the human fetal brain.

8.2 Male

8.2.1 *In vivo* human studies

Apart from oxidative stress, which has a negative impact on the quality of sperm, long-term stress manifested by an increased level of cortisol is also mentioned. This is related to male infertility due to the decreased conversion of androstenedione to testosterone, thereby reducing the volume and concentration of sperm in semen (202). Studies have shown that cortisol and adrenaline levels were significantly higher in the group of infertile men compared to the control group of men without fertility problems (194). Stress in this sense is both a physical and an emotional factor causing the activation of neurons that secrete the corticotropin-releasing hormone, leading to higher plasma cortisol levels. Excess cortisol reduces testosterone production, reducing sperm parameters. Cortisol may also directly reduce testosterone

production by blocking the transcription of genes encoding enzymes necessary for testosterone synthesis (195).

8.2.2 *In vivo* animal studies

High levels of adrenaline and associated low testosterone have also been observed in infertile golden hamsters (193). This phenomenon, also accompanying short-term stresses, results from the inhibitory effect of adrenaline and dopamine on testosterone production both by the HPA hypothalamic-pituitary axis and the peripheral blockade of testosterone release (203).

9 Ovaries

Estrogens belong to the family of steroid hormones produced mainly by the gonads and the placenta. In addition to the reproductive system, they play an important role in the immune, skeletal and neuroendocrine systems, therefore disturbances in their concentration or functioning are observed in pathological conditions and diseases such as infertility, cancer, obesity or osteoporosis (204). Estrogens have been shown to have negative and positive feedback on the hypothalamic-pituitary axis, and their action is possible by binding to the estrogen receptor alpha (ER α) and beta (ER β). These receptors have different tissue expression patterns in both humans and rodents (205). Estrogen mediation in biological responses is possible through a genomic mechanism, usually occurring within hours in most tissues, and a non-genomic mechanism, occurring very rapidly within minutes of exposure to hormones (206).

Estrogen binds to the ER, which is mainly found in the nucleus of the target cell. The resulting complex can regulate gene activity by binding directly to DNA regulatory elements called estrogen response elements (EREs), leading to the recruitment of additional factors involved in the regulation of transcription (207). An alternative approach, also called indirect, involves interaction with transcription factors, including TF, AP-1, SP1, and NF- κ B, which recruits chromatin-modifying coregulator proteins and enables activation or repression of ER target genes to direct cell proliferation (208). The non-genomic mechanism involves the interaction of the ER located in the plasma membrane or its vicinity with adapter proteins, including Shc and Src, as well as signaling through GPR30, resulting in activation of the MAPK cascades, PI3K, and adenylate cyclase (209).

Progesterone is an endogenous steroid hormone commonly produced by the cortex of the adrenal glands and the gonads. The ovarian follicles are the main source of peripheral progesterone from the late follicular to luteal phase (210). It is also secreted by the corpus luteum of the ovary, generally during the first ten weeks of pregnancy, and then through the placenta (211). The action of progesterone is based on binding to a receptor located in the cell cytoplasm. Then it dimerizes and translocates to the nucleus where it can bind to DNA, which enables the regulation of gene expression. There are three isoforms of progesterone receptors: PR-A, PR-B and PR-C (212).

9.1 Primary ovarian insufficiency

Primary ovarian insufficiency (POI) is defined as the cessation or irregular menstrual cycles under the age of 40 in the presence of

elevated serum FSH levels. This disorder is rare and affects approximately 1% of women. The causes of POIs can be spontaneous, genetic, environmental, infectious, autoimmune, surgical, chemotherapy or radiation related (213). The European Society of Human Reproduction and Embryology (ESHRE) recommends both of the following diagnostic criteria for POIs: hypomenorrhea/amenorrhea for at least four months, elevated follicle-stimulating hormone levels (> 25 mIU/mL) confirmed twice at an interval > 4 weeks (214). Serum FSH determination is the gold standard in the POI diagnosis. The anti-müllerian hormone can only be interpreted in conjunction with the FSH and estrogen levels. In the case of secondary amenorrhea, it is necessary to exclude pregnancy by testing the serum level of beta subunit of human chorionic gonadotropin (beta-HCG) and the concentration of thyroid stimulating hormone and prolactin, as endocrine diseases can lead to menstrual disorders (215). Chromosome analysis and fragile X premutation tests are recommended for all women with POI (216). As some cases of POI are autoimmune in nature, it is important to rule out other autoimmune diseases. According to ESHRE, it is necessary to assess the level of adrenal and thyroid antibodies. If the results are positive, it is imperative to monitor the function of these glands (214).

About 76% of POI patients maintain regular periods during adolescence and adulthood, followed by cycle disruptions (217). Ovarian function can be intermittent and unpredictable, with spontaneous ovulation in up to 20% and conception in approximately 5–10% of women (163). The symptoms these women experience are identical to those during menopause and can include hot flashes, night sweats, dyspareunia, vaginal dryness, sleep disturbances, mood changes, altered urination frequency, low libido, and a lack of energy. They are caused by a decrease in the production of estradiol in the ovaries. Symptoms may be transient or intermittent and may vary in severity due to fluctuations in ovarian activity during the spontaneous occurrence of POI (218). POI significantly reduces patients' quality of life due to increased cardiovascular risk, decreased bone mineral density leading to osteoporosis and atrophic changes in the genitourinary system. The disease has a negative impact on the mental well-being of patients, is associated with pregnancy failures and lower sexual satisfaction (219).

9.2 Polycystic ovary syndrome

Polycystic ovary syndrome (PCOS) is the most common endocrine disorder in women of childbearing age, affecting approximately 6–21% of women (220). PCOS can be diagnosed using the Rotterdam criteria, which require at least two of the three listed criteria: anovulation, clinical signs of hyperandrogenism and/or serological elevation of androgens, polycystic ovaries demonstrated by ultrasound. The National Institutes of Health criteria also require clinical or biochemical hyperandrogenism and oligo- or anovulation. The American Excess PCOS Society requires hyperandrogenism with one of the other two criteria. As this is an exclusion diagnosis, disorders that exhibit PCOS-like features must be ruled out. These include hyperprolactinemia, thyroid disease, non-classical congenital adrenal hyperplasia (221, 222).

In addition to infertility, many diseases are associated with PCOS, including endometrial cancer, type 2 diabetes, impaired glucose

tolerance, metabolic syndrome, cardiovascular risk, non-alcoholic fatty liver disease/non-alcoholic steatohepatitis (NAFLD/NASH) (223). It has been estimated that about 50% of women with PCOS are overweight or obese and have reduced insulin sensitivity (224). Almost all causes of PCOS arise from functional ovarian hyperandrogenism (FOH). Two-thirds of cases have typical FOH, characterized by androgen secretion disruption with an excessive response of 17-hydroxyprogesterone to gonadotrophin stimulation. In other cases of PCOS, an increase in testosterone levels is observed, which can be detected after suppressing the production of androgens in the adrenal glands (225). The causes of dysregulation include excess insulin, which sensitizes the ovaries to luteinizing hormone, as well as an imbalance between the intra-ovarian regulatory systems.

The excess of androgens enhances the growth of primary follicles. At the same time, it initiates premature luteinization, making the selection of the dominant follicle difficult (226). It is estimated that about half of FOH patients have insulin-resistant hyperinsulinism, which prematurely luteinizes granulosa cells, increases steroidogenesis, and stimulates fat accumulation. Hyperandrogenemia causes LH excess, which acts on luteinized granulosa and theca cells (227). Hormonal dysregulation alters the pulsatile release of GnRH, resulting in increased biosynthesis and secretion of LH compared to FSH (228). It has been shown that LH stimulates the production of androgens in the ovaries, while a decrease in FSH levels leads to the inhibition of aromatase activity in granulosa cells, reducing the conversion of androgens to estradiol (229). Moreover, serum androgens are converted peripherally to estrogens. Since it occurs mainly in adipose tissue, estrogen production will be increased in obese PCOS patients. Furthermore, unbalanced estrogen stimulation can lead to endometrial hyperplasia and endometrial cancer (230).

9.3 Female

9.3.1 *In vivo* human studies

Estrogens, in particular estradiol-17 β , are essential fertility regulating hormones. This is related to their participation in the development of ovulatory follicles, induction of pre-ovulatory gonadotropin release in the middle of the cycle, or preparation of the uterine mucosa for implantation. Changes in the production and/or function of estrogen can therefore disrupt these processes, leading to infertility (207) (Table 7). Studies carried out so far in models using rats have shown that disruption of *Esr1* causes infertility in both males and females (231). Rumi et al. also concluded that the proper functioning of *Esr2* is the primary regulator of female fertility, but is not critical to male fertility (232).

9.3.2 *In vivo* animal studies

Physiological estrogen replacement alleviates menopausal symptoms and may improve sexual dysfunction associated with vaginal dryness, dyspareunia, and decreased libido (251). Unfortunately, Hipp et al. showed that more than half of young women with POI either never use hormone replacement therapy (HRT) or start it many years after diagnosis and/or stop using it before the age of 45 (233). The current evidence supports transdermal or vaginal therapy with estradiol as the first line of HRT. They mimic the daily rate of ovarian estradiol production and achieve

mean serum level of 100 pg/mL, which is the average level in women with normal ovarian function throughout the menstrual cycle (252). Oral estradiol may be an alternative, but this is associated with complications related to the hepatic first-pass effect, including increasing the risk of venous thromboembolism (234). The multicenter study of Estrogen and Thromboembolism Risk (ESTHER) confirmed that the odds ratio for this disease was 4.2 in oral estrogen users compared to 0.9 in transdermal estrogen users (235). Interestingly, Renoux et al. also noted that the use of oral HRT was associated with a higher rate of stroke in postmenopausal women (253). In addition, Langrish et al. conducted a study in which they compared the treatment effects of transdermal estradiol and cyclic progestins with treatment with combined oral contraceptives on the circulatory system of young women with POI. They noticed that 12-month transdermal HRT resulted in significantly lower blood pressure, better kidney function and reduced activation of the renin-angiotensin-aldosterone system, suggesting that it was more effective (236).

More and more studies reveal that reproductive disorders can have a negative impact on pregnancy, from implantation to delivery. Many patients with these conditions require assisted reproductive technology (ART), which may influence pregnancy outcomes. As a result, it is difficult to distinguish the contribution of certain factors to poor pregnancy outcomes (254). It has been shown that often multiple risk factors contribute to negative obstetric outcomes in women with reproductive disorders. In addition, many women of childbearing age have more than one reproductive disorder (255).

In the case of POI, the chances of a spontaneous pregnancy are very small. According to the ESHRE guidelines, there are no interventions that increase the chances of natural conception, so oocyte donation or reception is considered a reliable chance to become pregnant (214). Interestingly, most people with POI still have detectable follicles. The mechanism of follicle dysfunction was defined in the NIH study as its inappropriate luteinization due to tonically increased serum LH levels. HRT that suppress high LH levels, can increase the chances of ovulation and improve follicle function. Piedade et al. presented a case report where a patient chose intra-uterine insemination (IUI) in combination with follicle monitoring and HRT and successfully became pregnant (237). In turn, Pinelli et al. conducted a study in which women with reduced ovarian reserve were administered valerate estradiol (2 mg daily) adding dihydrogesterone (10 mg daily) in the luteal phase for 3 months prior to the standard short protocol with a GnRH antagonist for *in vitro* fertilization. They concluded that pretreatment with estrogen appeared to improve IVF outcomes (238).

For overweight and obese women with PCOS, exercise and calorie restriction diets are the best first-line interventions for weight loss and impaired glucose tolerance (IGT) reduction. It is possible to alleviate the clinical symptoms of PCOS, including abnormal ovulation and hormone disturbances. Zhang et al. in their meta-analysis assessed the effectiveness of a low-carbohydrate diet in the treatment of PCOS and showed its beneficial effects (239). Treatments for menstrual disorders, hirsutism and acne include oral contraceptives, transdermal patch or vaginal ring. The progestogen component lowers LH levels, indirectly reducing the production of androgens in the ovaries and increasing the level of sex hormone binding globulin. Metformin is recommended in case of contraindications to the use of hormonal contraceptives. It has been shown to reduce

TABLE 7 Ovarian-related fertility studies.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Rumi et al. (231)	<i>In vivo</i> animal (rat) female	Animal medical experiment	Analysis of <i>Esr1</i> -knockout rats generated by ZFN-mediated genome editing	$\Delta 482$ <i>Esr1</i> mutation created a null allele	Disturbance of ESR1 activity leads to infertility of male and female rats
Rumi et al. (232)	<i>In vivo</i> animal (rat) female	Animal medical experiment	Assessment of the role of ESR2 in fertility regulation	Reproductive functions were impaired in mutant rats	Disturbance of ESR2 activity leads to female infertility
Hipp et al. (233)	<i>in vivo</i> human female	Semi-structured interviews	Analysis of diagnostic and therapeutic regimens in people with fragile X-associated POI	POI diagnosis and hormone treatment occurred later	Women with FXPOI are at risk for delayed POI diagnosis and undertreatment with hormone therapy
Mohammed et al. (234)	<i>in vivo</i> human female	Meta-analysis	Risk comparison of vascular events during the use of oral and transdermal estrogen therapy in postmenopausal women	A higher percentage of vascular events occurred in women using oral estrogens	Oral estrogen therapy may be associated with a higher risk of vascular events compared to transdermal therapy
Canonico et al. (235)	<i>in vivo</i> human female	Clinical experiment	Assessment of the impact of HRT administration on the risk of venous thromboembolism	The odds ratio for venous thromboembolism was higher in women using oral estrogens	Transdermal estrogen therapy is associated with a lower risk of venous thromboembolism
Renoux et al. (235)	<i>in vivo</i> human female	Comparative study	Assessment of the impact of HRT administration on the stroke risk	A higher stroke rate was reported in the oral HRT group	Oral HRT may increase the risk of stroke
Langrish et al. (236)	<i>in vivo</i> human female	Randomized controlled trial	Evaluation of HRT effects in women with POI	Better kidney function and lower blood pressure have been demonstrated for the physiological substitution of sex steroids	Physiological sex steroid substitution is a better option for young women with POI
Piedade et al. (237)	<i>in vivo</i> human female	Observational study (case report)	Analysis of the therapeutic regimen in women with POI	The patient became pregnant and gave birth to a healthy child	Actions should be taken to select the optimal treatment method
Pinelli et al. (238)	<i>in vivo</i> human female	Retrospective analysis	Analysis of estrogen treatment regimen in women with POI	FSH levels were lower and the number of collected and fertilized MII oocytes was higher	Estrogen treatment facilitates conception and improves IVF outcomes
Zhang et al. (239)	<i>in vivo</i> human female	Meta-analysis	Assessment of the impact of a low-carbohydrate diet on the clinical symptoms of PCOS	A low-carbohydrate diet helps to improve the outcomes of PCOS patients	Proper control of carbohydrate intake can alleviate the symptoms of PCOS
Glintborg et al. (240)	<i>in vivo</i> human female	Randomized controlled trial	Evaluation of the effects of metformin use in patients with PCOS	The use of metformin promotes weight loss and improves body composition	Treatment with metformin in combination with oral contraceptive pills is preferable to treatment with the pill alone
Takasaki et al. (241)	<i>in vivo</i> human female	Clinical experiment	Assessment of the effectiveness of modification of standard clomiphene treatment in patients resistant to this drug	About 80% of patients had positive results	Intermittent clomiphene treatment is a useful option in people who are resistant to this drug
Homburg et al. (242)	<i>in vivo</i> human female	Randomized controlled trial	Comparison of the effectiveness of clomiphene and FSH treatment in patients with PCOS	A higher percentage of pregnancies and live births after using FSH instead of clomiphene has been shown	FSH may be an appropriate first-line therapy for some women with PCOS
Waanbah et al. (243)	<i>in vivo</i> human female	Observational study (cohort study)	Comparison of the effectiveness of clomiphene and letrozole treatment in patients with PCOS	A higher percentage of pregnancies and healthy births was observed after treatment with letrozole	Letrozole is a better ovulation inducer than clomiphene in women with PCOS
Ganie et al. (244)	<i>in vivo</i> human female	Randomized controlled trial	Assessment of the effectiveness of spironolactone and metformin treatment in patients with PCOS	Combination treatment gave better results	A low dose of spironolactone and metformin may have beneficial effects in the treatment of PCOS
Glintborg et al. (245)	<i>in vivo</i> human female	Randomized controlled trial	Assessment of the effect of oral contraceptives and metformin on GLP-1 secretion in patients with PCOS	GLP-1 levels increased during treatment	The use of oral contraceptives and metformin affects the secretion of GLP-1

(Continued)

TABLE 7 Continued

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Devin et al. (246)	<i>in vivo</i> human female	Randomized controlled trial	Assessment of the relationship between the levels of growth hormone and sitagliptin and its impact on visceral obesity in patients with PCOS	Sitagliptin lowered visceral fat levels and increased growth hormone half-life and pulse spacing	Sitagliptin may be useful in the treatment of PCOS in obese women
Javed et al. (247)	<i>in vivo</i> human female	Randomized controlled trial	Assessment of the effect of empagliflozin on the metabolic parameters of patients with PCOS	Anthropometric parameters and body composition improved after treatment with empagliflozin	Empagliflozin may be useful in the treatment of PCOS in obese women
Benito et al. (248)	<i>in vivo</i> human female	Observational study (cohort study)	Assessment of the fertility level in women with PCOS after bariatric surgery	Pregnancy and fertility rates were high after surgery and there were few maternal and neonatal complications	Bariatric surgery has a positive effect on pregnancy and fertility indicators
Toulis et al. (249)	<i>in vivo</i> human female	Meta-analysis	Risk assessment of gestational diabetes in women with PCOS	There is a possible risk of gestational diabetes in women with PCOS	PCOS may be a potential risk factor for the development of gestational diabetes
De Frène et al. (250)	<i>in vivo</i> human female	Comparative study	Assessment of the influence of overweight and PCOS on the risk of perinatal complications	Women with PCOS and who were overweight had an increased risk of perinatal complications	Weight loss before pregnancy can reduce the risk of perinatal complications

the progression from impaired glucose tolerance to type 2 diabetes and also to improve menstrual cycles and vascular markers in non-obese women with PCOS (240).

Clomiphene citrate is considered the first-line treatment for infertility in PCOS patients. It is a selective estrogen receptor modulator (SERM) that is a competitive inhibitor of estrogen receptors. Clomiphene improves fertility and ovulation, especially by acting on the hypothalamus, where it binds to estrogen receptors. This causes a pulsatile release of GnRH, promoting the secretion of gonadotrophins from the anterior pituitary. Interestingly, up to 25% of women do not respond to the administration of clomiphene, which is associated with the implementation of gonadotropin therapy. However, Takasaki et al. proposed to modify the clomiphene treatment regimen. They used intermittent dosing, which was effective in approximately 80% of clomiphene-resistant patients (241). On the other hand, Homburg et al. showed that a small dose of FSH could be used instead of clomiphene (242). Another first-line drug for ovulation induction is letrozole. It is an aromatase inhibitor that blocks estrogen synthesis, reducing negative estrogen feedback in the pituitary gland. Waanbah et al. showed that treatment with letrozole, compared to clomiphene, was associated with higher rates of live births and ovulation among infertile women with polycystic ovary syndrome (243). Spironolactone, which is an antagonist of the mineralocorticoid receptor, is also used, as well as its combinations with metformin, which helped to improve the menstrual cycle, glucose and testosterone levels (244).

In PCOS patients, insulin sensitization is also introduced. It is associated with the administration of glucagon-like peptide 1 (GLP-1) agonists which, upon binding to the receptor, stimulate glucose-dependent insulin release from the pancreatic islets. This treatment reduced BMI and testosterone, and improved the ovulation rate in obese women with PCOS (256). Glintborg et al. showed that after treatment with oral contraceptives along with metformin, the level of GLP-1 in women with PCOS was comparable to that in healthy people (245). In addition, Devin et al. conducted studies using DPP4 inhibitors that have beneficial effects on weight loss and lowering

blood glucose levels in obese women with PCOS. They reported that the effect of DPP4 inhibitors on the body weight of women with PCOS is based on the increase in the growth hormone levels, which are lowered in PCOS patients. This, in turn, reduces the mass of visceral adipose tissue (246). On the other hand, Javed et al. noted the promising effect of SGLT2 inhibitors in overweight and obese women with PCOS. After 12 weeks of treatment with empagliflozin, there was an improvement in body composition and anthropometric parameters compared to metformin treatment, however no changes in hormonal and metabolic parameters were observed (247). Interestingly, Benito et al. conducted a study in women with PCOS who had undergone bariatric surgery. They observed a significant improvement in pregnancy and healthy birth rates compared to the control group (248). In turn, Toulis et al. showed that up to 40% of women with PCOS and glucose intolerance can develop gestational diabetes (GDM) (249). There was also an increased risk of pregnancy and neonatal complications, including premature pregnancy loss, gestational hypertension, preterm delivery, low birth weight and the need for cesarean delivery, regardless of obesity (250).

10 Testes

The testes are paired, oval glands located in the scrotum. Their main functions include the production of male gametes (sperm) and male sex hormones. Sperm formation is carried out by the germinal epithelium located in the convoluted tubules of the testes, while the production of testosterone, the main male sex hormone, is the responsibility of Leydig cells. These cells also produce a small amount of androstenedione (A4) (257). The functioning of the testes depends mainly on LH and FSH, as LH stimulates the production of testosterone, and FSH promotes the production of male gametes (258).

The testes are not only the site of testosterone synthesis, but also the main organ regulated by its activity. Testosterone stimulates spermatogenesis by affecting not only spermatogonial stem cells,

but also Sertoli cells, which support spermatogenesis and create a blood-testis barrier within the epithelium (259). In addition, testosterone is the main hormone involved in male sexual maturation (260). Other functions of testosterone are the stimulation of bone and muscle mass growth, and the regulation of sexual drive (261). Unlike LH, FSH acts directly on epithelial cells. It regulates the function of Sertoli cells, stimulating them to paracrine activity, promoting the maturation of spermatogonia into spermatocytes (262). The large increase in testosterone production by the testes in adolescence is one of the main factors in stimulating puberty. During sexual maturation, its synthesis increases 20-30 times (263). Due to this, not only physical but also mental changes are induced, such as increased sex drive, tendencies towards non-physical aggressive behavior and domination (264).

The testes, although being the main androgen-producing organ, are not the only ones. Another source of androgens is the adrenal glands. They produce dehydroepiandrosterone (DHEA) and androstenedione (265). DHEA is an anabolic hormone and strongly influences cognitive functions such as mood and sex drive (266). A4 is a testosterone precursor and has an effect similar to testosterone, but weaker. Both hormones are involved in adrenarche, and the increase in their concentration at the age of 6-8 years is probably of key importance in the development of social and cultural skills (267). Impaired testosterone secretion, regardless of whether it results from primary organ failure (Klinefelter syndrome, Leydig cell aplasia, genetic or drug-induced defects) or a defect associated with the secretion of pituitary hormones (Kallmann's syndrome, hypopituitarism, genetic or drug-induced defects in LH secretion) is a well-known cause of male infertility (268) (Table 8).

10.1 Male

10.1.1 *In vivo* human studies

The percentage of infertile men ranges from 2.5-12% depending on the studied population (289). Infertile males account for 40-50% of couple infertility, and the trend is increasing (290). The main cause of male infertility is poor semen quality, i.e. changes in physical parameters and volume, a decrease in the number of sperm per unit volume, reduced sperm motility and viability, abnormal sperm morphology. There are many reasons for a decline in sperm quality, from primary and secondary testicular failure to numerous genetic, environmental and lifestyle factors (291).

The genetic factors include not only numerical and structural chromosomal aberrations, but also microdeletions of the Y chromosome (292) and mutations of autosomal genes (293). The genes whose mutations have been associated with the reduction of male fertility include methylenetetrahydrofolate reductase (MTHFR), which is important in DNA methylation in stem cells (294), steroid 5-alpha reductase responsible for the conversion of testosterone into dihydrotestosterone (DHT), i.e. a more active metabolite (295), protamines replacing histone proteins during the condensation of the nuclear chromatin of sperm (296), heat shock protein A2 (HSPA2) responsible for the protection of cells from exposure to stress factors (269).

Environmental and lifestyle factors are mainly related to the induction of oxidative stress (297). Reactive oxygen species disrupt

sperm functions and their viability, damaging cell membranes, organelles and genetic material (298). It has been proven that oral administration of antioxidants may be effective in the treatment of some patients with infertility (299).

A positive correlation has also been demonstrated between high plasma testosterone concentration and fertility (300). An important finding is that the administration of exogenous testosterone, which reduces the production of endogenous hormone by suppressing the pituitary gland, correlates with a decrease in male fertility (301). However, it has been proven that early testosterone therapy after spinal cord injury increases the chances of preserving fertility in men (270).

10.1.2 *In vivo* animal studies

Animal studies focus mainly on establishing the relationship of gene expression within the testes with fertility. It has been demonstrated with such genes as ADAD1 and ADAD2 (271), Kindlin-2 (272), Wip 1 (273), and Crybb2 (274). At the same time, the lack of influence on fertility of a dozen or so genes, which are expressed within the testes, has been confirmed (275-279).

Numerous studies have also focused on establishing the effects of various toxins on fertility. It has been shown that exposure to microcystin-leucine-arginine and T-2 toxin suppresses the release of FSH in the pituitary gland, which, by reducing testosterone levels, decreases fertility (280, 281). On the other hand, exposure to mequinox or Mn₃O₄ microparticles reduces fertility due to the induction of oxidative stress within the testes (282). Sleep deprivation has a similar effect, with a reversible decline in fertility due to increased oxidative stress (283).

There are also studies related to the treatment of male infertility. The use of metformin in obese males has been shown to increase fertility (284). The effectiveness of supplementation with apple star fruit in order to improve fertility has also been shown (285). It is probably related to the antioxidant activity of both substances (302).

10.1.3 *In vitro* human studies

Currently, *in vitro* research is focused on determining the relationship between gene expression in testicular cells and fertility. The expression of APP, a protein related to the etiology of Alzheimer's disease, which influences numerous genes directly related to fertility, has been shown to be of great importance (286). Due to bioinformatic analysis, it was possible to determine the potential relationship of 30 genes with fertility and to exclude 54 genes from this group (287, 288).

10.2 The relationship between androgens and female fertility

Recently, increasing attention has been paid to the relationship between female fertility and blood levels of androgens (Table 9). Their sources are the ovaries, adrenal glands and the conversion of estrogen into androgens in other tissues (310). The concentration of this hormone is not subject to lifestyle fluctuations, except for smoking, which increases its level (311). The causes of the increase in androgen levels in women may include, apart from exogenous supplementation, polycystic

TABLE 8 Male fertility studies.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Mohanty et al. (269)	<i>in vivo</i> human male	Bioinformatic analysis of available databases	Assessment of the effect of HSPA2 polymorphism on male fertility	The existence of a variant of the HSPA2 gene negatively affecting fertility was demonstrated	HSPA2 polymorphism may cause male infertility
Choobineh et al. (270)	<i>in vivo</i> human male	Clinical experiment	Assessment of the effect of testosterone administration on the reduction of the negative impact on the testes after spinal cord injury	Increased semen quality in the study group was demonstrated	Early administration of testosterone after a spinal cord injury may positively influence fertility preservation
Snyder et al. (271)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Analysis of the effect of ADAD1 and ADAD2 expression on male fertility	Mice with blocked expression of ADAD1 and ADAD2 were completely sterile	ADAD1 and ADAD2 are key genes related to fertility
Chi et al. (272)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Assessment of the effect of Kindlin-2 expression on male fertility	Testicular hypoplasia and male infertility have been observed after Kindlin-2 expression was blocked	Kindlin-2 is a key gene related to fertility
Wei et al. (273)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Establishing the mechanism of reducing fertility in males with the Wip1 phosphatase mutation	Changes in the expression of numerous proteins related to cell adherence, apoptosis, response to the stimulation of pro-inflammatory cytokines and spermatogenesis were detected	Disabling Wip1 alters the expression of numerous fertility-related genes
Xiang et al. (274)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Assessment of the effect of Crybb2 expression on male fertility	After blocking Crybb2, fertility decreased significantly	Crybb2 is one of the key fertility genes
Sun et al. (275)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Searching for genes important for male reproduction using the CRISP method	A significant role of 13 genes in spermatogenesis and sperm maturation was determined	Genetic mutations can directly lead to infertility
Xia et al. (276)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Analysis of the effect of testis-expressed protein 33 (TEX33) expression on male fertility	Mice with blocked TEX33 expression showed no impairment of fertility	TEX33 is not related to fertility
Shah et al. (277)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Analysis of the influence of c4orf46 expression on male fertility	Mice with blocked c4orf46 expression showed no impairment of fertility	C4orf46 is not related to fertility
Holcomb et al. (278)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Analysis of the effect of expression of testis-specific serine proteases PRSS44, PRSS46, and PRSS54 on male fertility	Mice with blocked expression of testis-specific serine proteases PRSS44, PRSS46, and PRSS54 showed no impairment of fertility	Testis-specific serine proteases PRSS44, PRSS46, and PRSS54 are probably not related to fertility
He et al. (279)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Assessment of the effect of Trim69 expression on male fertility	No changes in fertility were observed after Trim69 blockade	Trim69 is not a key gene associated with fertility
Xiuying et al. (280)	<i>in vivo</i> animal male	Animal medical experiment (Rana nigromaculata)	Assessment of the effect of microcystin-leucine-arginine on reproduction	Decrease in fish synthesis after mclr administration	Exposure to mclrs negatively affects fertility
Yang et al. (281)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Assessment of the effect of T-2 toxin on male fertility	The administration of T-2 toxin caused a significant decrease in the concentration of hormones of the hypothalamic-pituitary-testis axis, which reduced fertility	Exposure to T-2 toxin negatively affects fertility
Zhang et al. (282)	<i>in vivo</i> animal male	Animal medical	Assessment of the effect of exposure to Mn ₃ O ₄ nanoparticles on male fertility	The administration of Mn ₃ O ₄ nanoparticles caused damage to the testes and decreased fertility	Exposure to Mn ₃ O ₄ nanoparticles negatively affects fertility

(Continued)

TABLE 8 Continued

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
		experiment (rat)			
Dominguez-Salazar et al. (283)	<i>in vivo</i> animal male	Animal medical experiment (rat)	Analysis of the effect of sleep deprivation on the blood-testicular barrier in males	In rats subjected to sleep deprivation, a decrease in the expression of proteins that build the blood-testicular barrier was observed, which leads to reduced fertility; fertility functions return to normal after sleep deprivation is reversed	Sleep deprivation has a negative effect on fertility, however it is reversible
Ye et al. (284)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Assessment of the effect of metformin administration in obese males on fertility	Increased fertility in the study group was demonstrated	Metformin can be used to treat male fertility
Akomolafe et al. (285)	<i>in vivo</i> animal male	Animal medical experiment (mouse)	Assessment of the effect of star apple fruit on male fertility	Increased fertility in the study group was demonstrated	Star apple fruit can be used as a nutritional supplement to improve fertility
Silva et al. (286)	<i>in vitro</i> human male	Bioinformatics analysis of gene relationships (in men)	Searching for links between amyloid precursor protein (APP) and genes related to fertility	Numerous relationships of APP expression have been determined not only with genes related to spermatogenesis and sperm maturation, but also with genes regulating the interaction of sperm with oocytes	APP expression may be related to fertility
Miyata et al. (287)	<i>in vivo</i> animal male	Bioinformatic analysis of available databases	Analysis of potential genes related to male fertility	54 genes potentially unrelated to male fertility were identified	The pool of genes suspected of being associated with male fertility was limited
Lu et al. (288)	<i>in vivo</i> animal male	Bioinformatic analysis of available databases	Analysis of potential genes related to male fertility	30 genes potentially unrelated to male fertility were identified	The pool of genes suspected of being associated with male fertility was limited

ovary syndrome, adrenal hyperplasia, Cushing's disease and neoplasms (312).

It has been shown that the increased concentration of androgens increases the frequency of anovulatory cycles (303). The effect of increased testosterone concentration on pregnancy is difficult to clearly establish, as both no effect on the course of pregnancy (304) and an increased risk of preeclampsia (305) have been demonstrated. An increased generation of pro-inflammatory factors within the uterine arteries was also observed, which may have a potentially negative impact on the course of pregnancy (306).

At the same time, it has been shown that the reduced testosterone concentration in women undergoing assisted reproductive protocols may be the cause of a poor response to ovarian stimulation (307). Optimal testosterone concentrations have been demonstrated in terms of the effectiveness of assisted reproductive protocols (308) and oral testosterone supplementation in order to increase the chances of success of such treatments (309).

11 Summary

More and more couples are affected by fertility and pregnancy maintenance disorders. It is estimated that male infertility accounts for 40-50% of couples' infertility. It is associated with impaired testosterone secretion, which may result from primary organ failure or is associated with the disruption of the secretion of pituitary hormones. Normal

ovarian function and placental secretion are essential for female fertility and the maintenance of pregnancy. The most common endocrine disorder in women of reproductive age is polycystic ovary syndrome, where androgen secretion is impaired, and an increase in testosterone levels is also possible. Such dysregulation results in a disruption of the pulsatile secretion of gonadotropin-releasing hormone by the hypothalamus, which in turn leads to abnormal gonadotropin secretion. There is also a rare disorder characterized by inhibition or irregular menstrual cycles known as primary ovarian insufficiency. Fertility is also influenced by autoimmune diseases, the toxicity of disease-modifying drugs, in particular non-steroidal anti-inflammatory drugs and glucocorticoids. Interestingly, HIV infection is associated with an increased risk of pituitary gland stroke and pituitary lymphoma, possibly leading to hypopituitarism, one of the effects of which is infertility. The pituitary-ovarian axis can also be influenced by thyroid hormones. Hypothyroidism leads to its inhibition, which increases the risk of miscarriage, premature births, stillbirths, and even infertility.

The form of infertility treatment depends on its duration and the patient's age. It includes ovulation stimulation with clomiphene citrate or gonadotropins, intrauterine insemination, *in vitro* fertilization. Therapeutic regimens are modified and various methods of drug administration are tried to increase their effectiveness and minimize side effects. However, it is important to correctly diagnose the disorder and take the required medications. Obesity is also getting more and more attention, as it is an important factor contributing to the reduction of fertility and predisposing to

TABLE 9 The relationship of testosterone and female fertility.

Authors (reference)	Type of study	Study design	Aim	Results	Conclusion
Sjaarda et al. (303)	<i>in vivo</i> human female	Clinical experiment	Assessment of the effect of testosterone and anti-müllerian hormone levels on conception and pregnancy	In the group of women with a high concentration of testosterone or amh, non-ovulatory cycles are statistically significantly more frequent; the influence of changes in the concentration of these hormones does not affect the course of pregnancy	High levels of testosterone or amh can negatively affect fertility
Lathi et al. (304)	<i>in vivo</i> human female	Clinical experiment	Assessment of the effect of testosterone levels on pregnancy in women with PCOS	No statistically significant differences in the course of pregnancy were shown	Testosterone in patients with PCOS does not adversely affect pregnant patients
Valdimarsdottir (305)	<i>in vivo</i> human female	Clinical experiment	Assessment of the effect of testosterone levels in the second trimester on pregnancy in women with PCOS and the relationship between testosterone levels and body weight	The higher the body weight, the higher the testosterone concentration; high testosterone levels during the second trimester positively correlate with the risk of preeclampsia	High testosterone levels in pregnant women may result in a higher risk of preeclampsia
Chinnathambi et al. (306)	<i>in vivo</i> animal female	Animal medical experiment (rat)	Assessment of the effect of testosterone on the condition of the uterine arteries during pregnancy in a female	A significant increase in the expression of pro-inflammatory factors and changes in the expression of factors determining the tension in the artery wall were demonstrated	Testosterone can negatively affect the vascularization of the uterus in pregnancy
Sun et al. (307)	<i>in vivo</i> human female	Clinical experiment	Assessment of the influence of testosterone concentration on the effectiveness of ovarian stimulation and the course of pregnancy achieved by IVF	Low testosterone levels correlate with a poor response to ovarian stimulation, no association with the course of pregnancy has been demonstrated	The use of testosterone during ovarian stimulation may increase the effectiveness of these treatments
Chen et al. (308)	<i>in vivo</i> human female	Clinical experiment	Assessment of the influence of testosterone concentration in women on the effectiveness of the ovarian stimulation	Optimal testosterone concentrations have been demonstrated at various stages of the ovarian stimulation	The use of testosterone during the ovarian stimulation process may increase the effectiveness of these treatments
Saharkhiz et al. (309)	<i>in vivo</i> human female	Clinical experiment	Assessment of the influence of testosterone administration on ovarian response in IVF cycles	An improvement in ovarian response in women in the study group and a higher pregnancy rate was demonstrated	Testosterone administration in women with a poor ovarian response to stimulation may be effective in IVF

adverse pregnancy outcomes. Weight loss interventions that include exercise and calorie restriction diets seem to be a good idea. In obese people, metformin is also often used, which improves menstrual cycles, reduces progression from impaired glucose tolerance to type 2 diabetes, and is also recommended in the case of contraindications to the use of hormonal contraceptives. Therefore, infertility as well as disorders of the maintenance of pregnancy and childbirth are a multifactorial issue related to numerous relationships between endocrine organs. Understanding these relationships as well as the importance of the influence of external factors is necessary to develop new and better diagnostic and therapeutic schemes.

Author contributions

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Janicka A, Spaczynski RZ, Kozioł K, Radwan M, Kurzawa R. Assisted reproductive medicine in Poland, 2013–2016: Polish society of reproductive medicine and embryology (PTMRIE) and fertility and sterility special interest group of the Polish society of gynaecologists and obstetricians (SPiN PTGiP) report. *Ginekol Pol* (2021) 92:7–15. doi: 10.5603/GP.a2020.0142
- Vander Borgh T, Wyns C. Fertility and infertility: Definition and epidemiology. *Clin Biochem* (2018) 62:2–10. doi: 10.1016/j.clinbiochem.2018.03.012
- Silvestris E, de Pergola G, Rosania R, Loverro G. Obesity as disruptor of the female fertility. *Reprod Biol Endocrinol* (2018) 16:22. doi: 10.1186/s12958-018-0336-z
- Łukaszuk K, Kozioł K, Jakiel G, Jakimiuk A, Jędrzejczak P, Kuczyński W, et al. Diagnostyka i leczenie niepłodności — rekomendacje polskiego towarzystwa medycyny rozrodu i embriologii (PTMRIE) oraz polskiego towarzystwa ginekologów i położników (PTGP). *Ginekol Perinatol Prakt* (2018) 3:112–40.
- Wdowiak A, Bazylewicz A, Dolzhenko MN, Nosenko NN, Hdyrya O. Technologie medyczne w diagnostyce męskiej niepłodności. *Eur J Med Technol* (2015) 1:7–17.
- Khatun A, Rahman MS, Pang M-G. Clinical assessment of the male fertility. *Obstet Gynecol Sci* (2018) 61:179–91. doi: 10.5468/ogs.2018.61.2.179
- Janicka A, Spaczynski RZ, Kurzawa R, Sp P, Clinics F. Assisted reproductive medicine in Poland – fertility and sterility special interest group of the Polish gynaecological society (SPiN PTG) 2012 report. *Ginekol Pol* (2015) 86:932–39. doi: 10.17772/gp/60549
- Practice committee of the American society for reproductive medicine. Diagnostic evaluation of the infertile female: a committee opinion. *Fertil Steril* (2015) 103:e44–50. doi: 10.1016/j.fertnstert.2015.03.019
- van Rijswijk J, Pham CT, Dreyer K, Verhoeve HR, Hoek A, de Bruin J-P, et al. Oil-based or water-based contrast for hysterosalpingography in infertile women: a cost-effectiveness analysis of a randomized controlled trial. *Fertil Steril* (2018) 110:754–60. doi: 10.1016/j.fertnstert.2018.05.001
- Tomassetti C, D'Hooghe T. Endometriosis and infertility: Insights into the causal link and management strategies. *Best Pract Res Clin Obstet Gynaecol* (2018) 51:25–33. doi: 10.1016/j.bpobgyn.2018.06.002
- Practice committee of the American society for reproductive medicine. Electronic address: ASRM@asrm.org, practice committee of the American society for reproductive medicine. The role of immunotherapy in in vitro fertilization: A guideline. *Fertil Steril* (2018) 110:387–400. doi: 10.1016/j.fertnstert.2018.05.009
- Parma P, Radi O. Molecular mechanisms of sexual development. *Sex Dev Genet Mol Biol Evol Endocrinol Embryol Pathol Sex Determ Differ* (2012) 6:7–17. doi: 10.1159/000332209
- Stévant I, Nef S. Genetic control of gonadal sex determination and development. *Trends Genet TIG* (2019) 35:346–58. doi: 10.1016/j.tig.2019.02.004
- Biason-Laubler A. Control of sex development. *Best Pract Res Clin Endocrinol Metab* (2010) 24:163–86. doi: 10.1016/j.beem.2009.12.002
- Makiyan Z. Studies of gonadal sex differentiation. *Organogenesis* (2016) 12:42–51. doi: 10.1080/15476278.2016.1145318
- Lamothe S, Bernard V, Christin-Maitre S. Gonad differentiation toward ovary. *Ann Endocrinol* (2020) 81:83–8. doi: 10.1016/j.ando.2020.04.004
- Cunha GR, Baskin LS. Development of the external genitalia. *Differ Res Biol Divers* (2020) 112:7–9. doi: 10.1016/j.diff.2019.10.008
- Gustafson ML, Donahoe PK. Male Sex determination: current concepts of male sexual differentiation. *Annu Rev Med* (1994) 45:505–24. doi: 10.1146/annurev.med.45.1.505
- Abbara A, Clarke SA, Nesbitt A, Ali S, Comminos AN, Hatfield E, et al. Interpretation of serum gonadotropin levels in hyperprolactinaemia. *Neuroendocrinology* (2018) 107:105–13. doi: 10.1159/000489264
- Feng H, Zhong Q, Zhou H, Jiang X, Ding Y. The effects of combined bromocriptine and bu-shen-zhu-yun decoction on serum hormones, anxiety, and pregnancy in hyperprolactinemic infertility patients. *Ann Palliat Med* (2021) 10:12230–43. doi: 10.21037/apm-21-3111
- Sermondade N, Faure C, Fezeu L, Shayeb AG, Bonde JP, Jensen TK, et al. BMI in relation to sperm count: an updated systematic review and collaborative meta-analysis. *Hum Reprod Update* (2013) 19:221–31. doi: 10.1093/humupd/dms050
- Stamatiades GA, Kaiser UB. Gonadotropin regulation by pulsatile GnRH: Signaling and gene expression. *Mol Cell Endocrinol* (2018) 463:131–41. doi: 10.1016/j.mce.2017.10.015
- Stamatiades GA, Carroll RS, Kaiser UB. GnRH-a key regulator of FSH. *Endocrinology* (2019) 160:57–67. doi: 10.1210/en.2018-00889
- Lim CT, Khoo B. Normal physiology of ACTH and GH release in the hypothalamus and anterior pituitary in man. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al, editors. *Endotext*. South Dartmouth (MA: MDText.com, Inc (2000). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK279116/>.
- Espino J, Macedo M, Lozano G, Ortiz Á, Rodríguez C, Rodríguez AB, et al. Impact of melatonin supplementation in women with unexplained infertility undergoing fertility treatment. *Antioxidants* (2019) 8:338. doi: 10.3390/antiox8090338
- Mokhtari F, Akbari Asbagh F, Azmoodeh O, Bakhtiyari M, Almasi-Hashiani A. Effects of melatonin administration on chemical pregnancy rates of polycystic ovary syndrome patients undergoing intrauterine insemination: A randomized clinical trial. *Int J Fertil Steril* (2019) 13:225–9. doi: 10.22074/ijfs.2019.5717
- Çelik S, Guve H, Çalıřkan C, Çelik S. The role of melatonin, IL-8 and IL-10 in intrahepatic cholestasis of pregnancy. *Z Geburtshilfe Neonatol* (2021) 225:238–43. doi: 10.1055/a-1233-9084
- Hobson SR, Gurusinghe S, Lim R, Alers NO, Miller SL, Kingdom JC, et al. Melatonin improves endothelial function *in vitro* and prolongs pregnancy in women with early-onset preeclampsia. *J Pineal Res* (2018) 65:e12508. doi: 10.1111/jpi.12508
- Zhang C, Clough SJ, Adamah-Biassi EB, Sveinsson MH, Hutchinson AJ, Miura I, et al. Impact of endogenous melatonin on rhythmic behaviors, reproduction, and survival revealed in melatonin-proficient C57BL/6J congenic mice. *J Pineal Res* (2021) 71:e12748. doi: 10.1111/jpi.12748
- Lv D, Tan T, Zhu T, Wang J, Zhang S, Zhang L, et al. Leptin mediates the effects of melatonin on female reproduction in mammals. *J Pineal Res* (2019) 66:e12559. doi: 10.1111/jpi.12559
- Lombardo F, Giorgini E, Gioacchini G, Maradonna F, Ferraris P, Carnevali O. Melatonin effects on fundulus heteroclitus reproduction. *Reprod Fertil Dev* (2012) 24:794–803. doi: 10.1071/RD11267
- Zhang L, Zhang Z, Wang J, Lv D, Zhu T, Wang F, et al. Melatonin regulates the activities of ovary and delays the fertility decline in female animals via MT1/AMPK pathway. *J Pineal Res* (2019) 66:e12550. doi: 10.1111/jpi.12550
- Cosso G, Luridiana S, Pulinas L, Curone G, Pich G, Carcangiu V, et al. Melatonin treatment in rams and their replacement with novel treated rams advance first lambing and increase fertility in sarda ewe lambs. *Anim Open Access J MDPI* (2021) 11:1227. doi: 10.3390/ani11051227
- Song C, Peng W, Yin S, Zhao J, Fu B, Zhang J, et al. Melatonin improves age-induced fertility decline and attenuates ovarian mitochondrial oxidative stress in mice. *Sci Rep* (2016) 6:35165. doi: 10.1038/srep35165
- Dholpuria S, Vyas S, Purohit GN, Pathak KML. Sonographic monitoring of early follicle growth induced by melatonin implants in camels and the subsequent fertility. *J Ultrasound* (2012) 15:135–41. doi: 10.1016/j.jus.2012.02.008
- Peng X, Cai X, Li J, Huang Y, Liu H, He J, et al. Effects of melatonin supplementation during pregnancy on reproductive performance, maternal-Placental-Fetal redox status, and placental mitochondrial function in a sow model. *Antioxid Basel Switz* (2021) 10:1867. doi: 10.3390/antiox10121867
- Arend LS, Knox RV. Fertility responses of melatonin-treated gilts before and during the follicular and early luteal phases when there are different temperatures and lighting conditions in the housing area. *Anim Reprod Sci* (2021) 230:106769. doi: 10.1016/j.anireprosci.2021.106769
- Bai J, Zhang L, Zhao Z, Li N, Wang B, Yang L. Expression of melatonin receptors and CD4 in the ovine thymus, lymph node, spleen and liver during early pregnancy. *Immunology* (2020) 160:52–63. doi: 10.1111/imm.13180
- Gunwant P, Pandey AK, Kumar A, Singh I, Kumar S, Phogat JB, et al. Polymorphism of melatonin receptor (MTNR1A) gene and its association with seasonal reproduction in water buffalo (*Bubalus bubalis*). *Anim Reprod Sci* (2018) 199:51–9. doi: 10.1016/j.anireprosci.2018.10.006
- Fathy HA, Gouda EM, Gafer JA, Galal MK, Nowier AM. Genetic polymorphism in melatonin receptor 1A and arylalkylamine n-acetyltransferase and its impact on seasonal reproduction in Egyptian sheep breeds. *Arch Anim Breed* (2018) 61:505–16. doi: 10.5194/aab-61-505-2018
- Mura MC, Luridiana S, Bodano S, Daga C, Cosso G, Diaz ML, et al. Influence of melatonin receptor 1A gene polymorphisms on seasonal reproduction in sarda ewes with different body condition scores and ages. *Anim Reprod Sci* (2014) 149:173–7. doi: 10.1016/j.anireprosci.2014.07.022
- do Nascimento Marinho KS, Lapa Neto CJC, de Sousa Coelho IDD, da Silva MA, Gomes Melo ME, Dos Santos KRP, et al. Genotoxic and mutagenic evaluation of the protective effect of exogenous melatonin in adult rats and their offspring exposed to the insecticides methomyl and cypermethrin during pregnancy. *Mutat Res Genet Toxicol Environ Mutagen* (2019) 848:503107. doi: 10.1016/j.mrgentox.2019.503107
- Huang J, Shan W, Li N, Zhou B, Guo E, Xia M, et al. Melatonin provides protection against cisplatin-induced ovarian damage and loss of fertility in mice. *Reprod BioMed Online* (2021) 42:505–19. doi: 10.1016/j.rbmo.2020.10.001
- de Almeida LL, Teixeira AAC, Soares AF, da Cunha FM, da Silva VA, Vieira Filho LD, et al. Effects of melatonin in rats in the initial third stage of pregnancy exposed to sub-lethal doses of herbicides. *Acta Histochem* (2017) 119:220–7. doi: 10.1016/j.acthis.2017.01.003
- de Sousa Coelho IDD, Lapa Neto CJC, Souza TGDS, da Silva MA, Chagas CA, Santos KRPD, et al. Protective effect of exogenous melatonin in rats and their offspring on the genotoxic response induced by the chronic consumption of alcohol during pregnancy. *Mutat Res Genet Toxicol Environ Mutagen* (2018) 832–833:52–60. doi: 10.1016/j.mrgentox.2018.06.018
- Wang Z, Teng Z, Wang Z, Song Z, Zhu P, Li N, et al. Melatonin ameliorates paclitaxel-induced mice spermatogenesis and fertility defects. *J Cell Mol Med* (2022) 26:1219–28. doi: 10.1111/jcmm.17177

47. Abo El Gheit RE, Soliman NA, Nagla SA, El-Sayed RM, Badawi GA, Emam MN, et al. Melatonin epigenetic potential on testicular functions and fertility profile in varicocele rat model is mediated by silent information regulator 1. *Br J Pharmacol* (2022) 179:3363–81. doi: 10.1111/bph.15804
48. Xu Y-M, Hao G-M, Gao B-L. Application of growth hormone in *in vitro* fertilization. *Front Endocrinol* (2019) 10:502. doi: 10.3389/fendo.2019.00502
49. Lainez NM, Coss D. Obesity, neuroinflammation, and reproductive function. *Endocrinology* (2019) 160:2719–36. doi: 10.1210/en.2019-00487
50. Kurowska P, Mlyczyńska E, Dawid M, Jurek M, Klimczyk D, Dupont J, et al. Review: Vaspin (SERPINA2) expression and function in endocrine cells. *Cells* (2021) 10:1710. doi: 10.3390/cells10071710
51. Roberts RE, Farahani L, Webber L, Jayasena C. Current understanding of hypothalamic amenorrhoea. *Ther Adv Endocrinol Metab* (2020) 11:2024018820945854. doi: 10.1177/2042018820945854
52. Shufelt CL, Torbati T, Dutra E. Hypothalamic amenorrhea and the long-term health consequences. *Semin Reprod Med* (2017) 35:256–62. doi: 10.1055/s-0037-1603581
53. Saleem M, Martin H, Coates P. Prolactin biology and laboratory measurement: An update on physiology and current analytical issues. *Clin Biochem Rev* (2018) 39:3–16.
54. Koyada A, Orsu P. Role of hypothyroidism and associated pathways in pregnancy and infertility: Clinical insights. *Tzu-Chi Med J* (2020) 32:312–7. doi: 10.4103/tcmj.tcmj_255_19
55. Evanson J. Radiology of the pituitary. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al, editors. *Endotext*. South Dartmouth (MA: MDText.com, Inc (2000). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK279161/>.
56. Bachmeier CAE, Snell C, Morton A. Visual loss in pregnancy. *BMJ Case Rep* (2019) 12:e228323. doi: 10.1136/bcr-2018-228323
57. Kelleher AM, DeMayo FJ, Spencer TE. Uterine glands: Developmental biology and functional roles in pregnancy. *Endocr Rev* (2019) 40:1424–45. doi: 10.1210/er.2018-00281
58. Ergun-Longmire B, Wajrajch MP. Growth and growth disorders. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al, editors. *Endotext*. South Dartmouth (MA: MDText.com, Inc (2000). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK279142/>.
59. Kumar S, Kumar KH. Pituitary diseases in the tropics. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al, editors. *Endotext*. South Dartmouth (MA: MDText.com, Inc (2000). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK568567/>.
60. Miljic D, Pekic S, Popovic V. Empty sella. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al, editors. *Endotext*. South Dartmouth (MA: MDText.com, Inc (2000). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK532084/>.
61. Somers EC, Marder W. Infertility – prevention and management. *Rheum Dis Clin North Am* (2017) 43:275–85. doi: 10.1016/j.rdc.2016.12.007
62. Pekic S, Miljic D, Popovic V. Infections of the hypothalamic-pituitary region. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al, editors. *Endotext*. South Dartmouth (MA: MDText.com, Inc (2000). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK532083/>.
63. Prete A, Salvatori R. Hypophysitis. In: *Endotext* (2000). South Dartmouth (MA: MDText.com, Inc. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK519842/> (Accessed June 4, 2022).
64. Sciarra F, Franceschini E, Campolo F, Gianfrilli D, Pallotti F, Paoli D, et al. Disruption of circadian rhythms: A crucial factor in the etiology of infertility. *Int J Mol Sci* (2020) 21:3943. doi: 10.3390/ijms21113943
65. Rattan S, Zhou C, Chiang C, Mahalingam S, Brehm E, Flaws JA. Exposure to endocrine disruptors during adulthood: consequences for female fertility. *J Endocrinol* (2017) 233:R109–29. doi: 10.1530/JOE-17-0023
66. Borowska M, Czarnywojtek A, Sawicka-Gutaj N, Woliński K, Płazińska MT, Mikołajczak P, et al. The effects of cannabinoids on the endocrine system. *Endokrynol Pol* (2018) 69:705–19. doi: 10.5603/EP.a2018.0072
67. Darbandi M, Darbandi S, Agarwal A, Sengupta P, Durairajanayagam D, Henkel R, et al. Reactive oxygen species and male reproductive hormones. *Reprod Biol Endocrinol RBE* (2018) 16:87. doi: 10.1186/s12958-018-0406-2
68. Childs GV, Odle AK, MacNicol MC, MacNicol AM. The importance of leptin to reproduction. *Endocrinology* (2021) 162:bqaa204. doi: 10.1210/endo/bqaa204
69. Lateef OM, Akintubosun MO. Sleep and reproductive health. *J Circadian Rhythms* (2020) 18:1. doi: 10.5334/jcr.190
70. Selvaraj K, Ravichandran S, Krishnan S, Radhakrishnan RK, Manickam N, Kandasamy M. Testicular atrophy and hypothalamic pathology in COVID-19: Possibility of the incidence of Male infertility and HPG axis abnormalities. *Reprod Sci Thousand Oaks Calif* (2021) 28:2735–42. doi: 10.1007/s43032-020-00441-x
71. Youssef J, Sadara R, Mital D, Ahmed MH. HIV And the pituitary gland: Clinical and biochemical presentations. *J Lab Physicians* (2021) 13:84–90. doi: 10.1055/s-0041-1723055
72. Møller M, Baeres FM. The anatomy and innervation of the mammalian pineal gland. *Cell Tissue Res* (2002) 309:139–50. doi: 10.1007/s00441-002-0580-5
73. Amaral FGdo, Cipolla-Neto J. A brief review about melatonin, a pineal hormone. *Arch Endocrinol Metab* (2018) 62:472–9. doi: 10.20945/2359-399700000066
74. Arendt J, Aulinas A. Physiology of the pineal gland and melatonin. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al, editors. *Endotext*. South Dartmouth (MA: MDText.com, Inc (2000). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK550972/>.
75. Reiter RJ, Tan D-X, Tamura H, Cruz MHC, Fuentes-Broto L. Clinical relevance of melatonin in ovarian and placental physiology: a review. *Gynecol Endocrinol Off J Int Soc Gynecol Endocrinol* (2014) 30:83–9. doi: 10.3109/09513590.2013.849238
76. Srinivasan V, Pandi-Perumal SR, Trahlt I, Spence DW, Poeggeler B, Hardeland R, et al. Melatonin and melatonergic drugs on sleep: possible mechanisms of action. *Int J Neurosci* (2009) 119:821–46. doi: 10.1080/00207450802328607
77. Crowley SJ, Acebo C, Carskadon MA. Human puberty: Salivary melatonin profiles in constant conditions. *Dev Psychobiol* (2012) 54:468–73. doi: 10.1002/dev.20605
78. Waldhauser F, Weissenbacher G, Frisch H, Zeitlhuber U, Waldhauser M, Wurtman RJ. Fall in nocturnal serum melatonin during prepuberty and pubescence. *Lancet Lond Engl* (1984) 1:362–5. doi: 10.1016/s0140-6736(84)90412-4
79. Roy D, Belsham DD. Melatonin receptor activation regulates GnRH gene expression and secretion in GT1-7 GnRH neurons. *Signal transduction mechanisms. J Biol Chem* (2002) 277:251–8. doi: 10.1074/jbc.M108890200
80. Gingerich S, Wang X, Lee PKP, Dhillion SS, Chalmers JA, Koletar MM, et al. The generation of an array of clonal, immortalized cell models from the rat hypothalamus: analysis of melatonin effects on kisspeptin and gonadotropin-inhibitory hormone neurons. *Neuroscience* (2009) 162:1134–40. doi: 10.1016/j.neuroscience.2009.05.026
81. van Geijlswijk IM, Mol RH, Egberts TCG, Smits MG. Evaluation of sleep, puberty and mental health in children with long-term melatonin treatment for chronic idiopathic childhood sleep onset insomnia. *Psychopharmacol (Berl)* (2011) 216:111–20. doi: 10.1007/s00213-011-2202-y
82. Madalinski MH. Does a melatonin supplement alter the course of gastro-esophageal reflux disease? *World J Gastrointest Pharmacol Ther* (2011) 2:50–1. doi: 10.4292/wjgpt.v2.i6.50
83. Biran V, Phan Duy A, Decobert F, Bednarek N, Alberti C, Baud O. Is melatonin ready to be used in preterm infants as a neuroprotectant? *Dev Med Child Neurol* (2014) 56:717–23. doi: 10.1111/dmcn.12415
84. Bai J, Dong L, Song Z, Ge H, Cai X, Wang G, et al. The role of melatonin as an antioxidant in human lens epithelial cells. *Free Radic Res* (2013) 47:635–42. doi: 10.3109/10715762.2013.808743
85. Nawrot-Porąbka K, Jaworek J, Leja-Szpak A, Szklarczyk J, Konturek SJ, Reiter RJ. Luminal melatonin stimulates pancreatic enzyme secretion via activation of serotonin-dependent nerves. *Pharmacol Rep PR* (2013) 65:494–504. doi: 10.1016/s1734-1140(13)71025-9
86. Markus RP, Fernandes PA, Kinker GS, Cruz-Machado S da S, Marçola M. Immune-pineal axis – acute inflammatory responses coordinate melatonin synthesis by pinealocytes and phagocytes. *Br J Pharmacol* (2018) 175:3239. doi: 10.1111/bph.14083
87. Reiter RJ, Rosales-Corral SA, Manchester LC, Tan D-X. Peripheral reproductive organ health and melatonin: ready for prime time. *Int J Mol Sci* (2013) 14:7231–72. doi: 10.3390/ijms14047231
88. Reiter RJ, Tan DX, Osuna C, Gitto E. Actions of melatonin in the reduction of oxidative stress. A review. *J BioMed Sci* (2000) 7:444–58. doi: 10.1007/BF02253360
89. Chen C-Q, Fichna J, Bashashati M, Li Y-Y, Storr M. Distribution, function and physiological role of melatonin in the lower gut. *World J Gastroenterol WJG* (2011) 17:3888–98. doi: 10.3748/wjg.v17.i34.3888
90. Bubenik GA. Localization, physiological significance and possible clinical implication of gastrointestinal melatonin. *Biol Signals Recept* (2001) 10:350–66. doi: 10.1159/000046903
91. Carrillo-Vico A, Lardone PJ, Fernández-Santos JM, Martín-Lacave I, Calvo JR, Karasek M, et al. Human lymphocyte-synthesized melatonin is involved in the regulation of the interleukin-2/interleukin-2 receptor system. *J Clin Endocrinol Metab* (2005) 90:992–1000. doi: 10.1210/jc.2004-1429
92. Lewy AJ, Tetsuo M, Markey SP, Goodwin FK, Kopin IJ. Pinelectomy abolishes plasma melatonin in the rat. *J Clin Endocrinol Metab* (1980) 50:204–5. doi: 10.1210/jcem-50-1-204
93. Korkmaz A, Rosales-Corral S, Reiter RJ. Gene regulation by melatonin linked to epigenetic phenomena. *Gene* (2012) 503:1–11. doi: 10.1016/j.gene.2012.04.040
94. Li W, Kong A-N. Molecular mechanisms of Nrf2-mediated antioxidant response. *Mol Carcinog* (2009) 48:91–104. doi: 10.1002/mc.20465
95. Galano A, Tan DX, Reiter RJ. Cyclic 3-hydroxymelatonin, a key metabolite enhancing the peroxyl radical scavenging activity of melatonin. *RSC Adv* (2014) 4:5220–7. doi: 10.1039/C3RA44604B
96. Berga SL, Mortola JF, Yen SSC. Amplification of nocturnal melatonin secretion in women with functional hypothalamic amenorrhea. *J Clin Endocrinol Metab* (1988) 66:242–4. doi: 10.1210/jcem-66-1-242
97. Tang PL, Chan TY, Tang GWK, Pang SF. Plasma melatonin profile and hormonal interactions in the menstrual cycles of anovulatory infertile women treated with gonadotropins. *Gynecol Obstet Invest* (1998) 45:247–52. doi: 10.1159/00009977
98. Tamura H, Takasaki A, Taketani T, Tanabe M, Lee L, Tamura I, et al. Melatonin and female reproduction. *J Obstet Gynaecol Res* (2014) 40:1–11. doi: 10.1111/jog.12177
99. Jamilian M, Foroozanfar F, Mirhosseini N, Kavossian E, Aghadavod E, Bahmani F, et al. Effects of melatonin supplementation on hormonal, inflammatory, genetic, and oxidative stress parameters in women with polycystic ovary syndrome. *Front Endocrinol* (2019) 10:273. doi: 10.3389/fendo.2019.00273
100. Taub RL, Ellis SA, Neal-Perry G, Magaret AS, Prager SW, Micks EA. The effect of testosterone on ovulatory function in transmasculine individuals. *Am J Obstet Gynecol* (2020) 223:229.e1–8. doi: 10.1016/j.ajog.2020.01.059

101. Palacio JR, Iborra A, Ulcova-Gallova Z, Badia R, Martínez P. The presence of antibodies to oxidative modified proteins in serum from polycystic ovary syndrome patients. *Clin Exp Immunol* (2006) 144:217–22. doi: 10.1111/j.1365-2249.2006.03061.x
102. Ejaz H, Figaro JK, Woolner AMF, Thottakam BMV, Galley HF. Maternal serum melatonin increases during pregnancy and falls immediately after delivery implicating the placenta as a major source of melatonin. *Front Endocrinol* (2020) 11:623038. doi: 10.3389/fendo.2020.623038
103. Zheng M, Tong J, Li W-P, Chen Z-J, Zhang C. Melatonin concentration in follicular fluid is correlated with antral follicle count (AFC) and *in vitro* fertilization (IVF) outcomes in women undergoing assisted reproductive technology (ART) procedures. *Gynecol Endocrinol Off J Int Soc Gynecol Endocrinol* (2018) 34:446–50. doi: 10.1080/09513590.2017.1409713
104. Yang L, Wang Q, Cui M, Li Q, Mu S, Zhao Z. Effect of melatonin on the *In vitro* maturation of porcine oocytes, development of parthenogenetically activated embryos, and expression of genes related to the oocyte developmental capability. *Anim Open Access J MDPI* (2020) 10:E209. doi: 10.3390/ani10020209
105. Fathi M, Salama A, El-Shahat KH, El-Sherbiny HR, Abdelnaby EA. Effect of melatonin supplementation during IVM of dromedary camel oocytes (*Camelus dromedarius*) on their maturation, fertilization, and developmental rates. *vitro. Theriogenology* (2021) 172:187–92. doi: 10.1016/j.theriogenology.2021.05.021
106. Shu-Qin Z, Yong Z, Yuan G, Xiao-Pu Y, Zhen Y, Zhi-Jie Y. The *in vitro* effects of melatonin and cry gene on the secretion of estradiol from camel ovarian granulosa cells. *Domest Anim Endocrinol* (2021) 74:106497. doi: 10.1016/j.domaniend.2020.106497
107. Anderson RA, Lincoln GA, Wu FC. Melatonin potentiates testosterone-induced suppression of luteinizing hormone secretion in normal men. *Hum Reprod Oxf Engl* (1993) 8:1819–22. doi: 10.1093/oxfordjournals.humrep.a137940
108. Aghaz F, Vaisi-Raygani A, Khazaei M, Arkan E. Enhanced cryoprotective effect of melatonin and resveratrol by coencapsulation: Improved *In vitro* development of vitrified-warmed mouse germinal vesicle oocytes. *Biopreservation Biobanking* (2021) 19:184–93. doi: 10.1089/bio.2020.0102
109. Zhang Y, Li W, Ma Y, Wang D, Zhao X, Zeng C, et al. Improved development by melatonin treatment after vitrification of mouse metaphase II oocytes. *Cryobiology* (2016) 73:335–42. doi: 10.1016/j.cryobiol.2016.09.171
110. Tang Y, Zhang Y, Liu L, Yang Y, Wang Y, Xu B. Glycine and melatonin improve preimplantation development of porcine oocytes vitrified at the germinal vesicle stage. *Front Cell Dev Biol* (2022) 10:856486. doi: 10.3389/fcell.2022.856486
111. Awad H, Halawa F, Mostafa T, Atta H. Melatonin hormone profile in infertile males. *Int J Androl* (2006) 29:409–13. doi: 10.1111/j.1365-2605.2005.00624.x
112. Sharbatoghli M, Rezazadeh Valojerdi M, Bahadori MH, Salman Yazdi R, Ghaleno LR. The relationship between seminal melatonin with sperm parameters, DNA fragmentation and nuclear maturity in intra-cytoplasmic sperm injection candidates. *Cell J* (2015) 17:547–53. doi: 10.22074/cellj.2015.15
113. Ortiz A, Espino J, Bejarano I, Lozano GM, Monllor F, García JF, et al. High endogenous melatonin concentrations enhance sperm quality and short-term *in vitro* exposure to melatonin improves aspects of sperm motility. *J Pineal Res* (2011) 50:132–9. doi: 10.1111/j.1600-079X.2010.00822.x
114. Luboshitzky R, Shen-Orr Z, Nave R, Lavi S, Lavie P. Melatonin administration alters semen quality in healthy men. *J Androl* (2002) 23:572–8.
115. Lu X-L, Liu J-J, Li J-T, Yang Q-A, Zhang J-M. Melatonin therapy adds extra benefit to varicocelelectomy in terms of sperm parameters, hormonal profile and total antioxidant capacity: A placebo-controlled, double-blind trial. *Andrologia* (2018) 50: e13033. doi: 10.1111/and.13033
116. Cebi Sen C, Yumusak N, Atilgan HI, Sadic M, Koca G, Korkmaz M. The protective effect of melatonin on sperm quality in rat after radioiodine treatment. *Andrologia* (2018). doi: 10.1111/and.12962
117. Khalil SS, Aziz JA, Ismail KA, El-Malkey NF. Comparative protective effects of n-acetylcysteine and melatonin against obesity-induced testicular dysfunction in rats. *Can J Physiol Pharmacol* (2021) 99:708–19. doi: 10.1139/cjpp-2020-0499
118. Moradi M, Goodarzi N, Faramarzi A, Cheraghi H, Hashemian AH, Jalili C. Melatonin protects rats testes against bleomycin, etoposide, and cisplatin-induced toxicity via mitigating nitro-oxidative stress and apoptosis. *BioMed Pharmacother Biomedicine Pharmacother* (2021) 138:111481. doi: 10.1016/j.biopha.2021.111481
119. Venditti M, Ben Rhouma M, Romano MZ, Messaoudi I, Reiter RJ, Minucci S. Evidence of melatonin ameliorative effects on the blood-testis barrier and sperm quality alterations induced by cadmium in the rat testis. *Ecotoxicol Environ Saf* (2021) 226:112878. doi: 10.1016/j.ecoenv.2021.112878
120. Tijmes M, Pedraza R, Valladares L. Melatonin in the rat testis: evidence for local synthesis. *Steroids* (1996) 61:65–8. doi: 10.1016/0039-128x(95)00197-x
121. Nasiraei-Moghadam SN-M, Parivar K, Ahmadiani A, Movahhedin M, Vaez Mahdavi MR. Protective effect of melatonin against inequality-induced damage on testicular tissue and sperm parameters. *Int J Fertil Steril* (2014) 7:313–22.
122. Deng S-L, Chen S-R, Wang Z-P, Zhang Y, Tang J-X, Li J, et al. Melatonin promotes development of haploid germ cells from early developing spermatogenic cells of Suffolk sheep under *in vitro* condition. *J Pineal Res* (2016) 60:435–47. doi: 10.1111/jpi.12327
123. Karimfar MH, Niazvand F, Haghani K, Ghafourian S, Shirazi R, Bakhtiyari S. The protective effects of melatonin against cryopreservation-induced oxidative stress in human sperm. *Int J Immunopathol Pharmacol* (2015) 28:69–76. doi: 10.1177/0394632015572080
124. Syrenicz A, Syrenicz M, Sworcak K, Garanty-Bogacka B, Zimnicka A, Walczak M. [Hashimoto disease and hypothyroidism in child-bearing period—essential problem for woman and her child]. *Endokrynol Pol* (2005) 56:1008–15.
125. Poppe K, Glinoe D. Thyroid autoimmunity and hypothyroidism before and during pregnancy. *Hum Reprod Update* (2003) 9:149–61. doi: 10.1093/humupd/dmg012
126. Vaquero E, Lazzarin N, De Carolis C, Valensise H, Moretti C, Ramanini C. Mild thyroid abnormalities and recurrent spontaneous abortion: diagnostic and therapeutic approach. *Am J Reprod Immunol N Y N* 1989 (2000) 43:204–8. doi: 10.1111/j.8755-8920.2000.430404.x
127. Kakita-Kobayashi M, Murata H, Nishigaki A, Hashimoto Y, Komiya S, Tsubokura H, et al. Thyroid hormone facilitates *in vitro* decidualization of human endometrial stromal cells via thyroid hormone receptors. *Endocrinology* (2020) 161: bqaa049. doi: 10.1210/endo/bqaa049
128. Barber KJ, Franklyn JA, McCabe CJ, Khanim FL, Bulmer JN, Whitley GSJ, et al. The *in vitro* effects of triiodothyronine on epidermal growth factor-induced trophoblast function. *J Clin Endocrinol Metab* (2005) 90:1655–61. doi: 10.1210/jc.2004-0785
129. Giuliani C, Bucci I, Di Santo S, Rossi C, Grassadonia A, Piantelli M, et al. The flavonoid quercetin inhibits thyroid-restricted genes expression and thyroid function. *Food Chem Toxicol Int J Publ Br Ind Biol Res Assoc* (2014) 66:23–9. doi: 10.1016/j.fct.2014.01.016
130. Condorelli RA, La Vignera S, Mongioi LM, Alamo A, Giaccone F, Cannarella R, et al. Thyroid hormones and spermatozoa: *In Vitro* Effects on sperm mitochondria, viability and DNA integrity. *J Clin Med* (2019) 8:E756. doi: 10.3390/jcm8050756
131. Dallas JS. Autoimmune thyroid disease and pregnancy: relevance for the child. *Autoimmunity* (2003) 36:339–50. doi: 10.1080/08916930310001602993
132. Krassas GE. Thyroid disease and female reproduction. *Fertil Steril* (2000) 74:1063–70. doi: 10.1016/s0015-0282(00)01589-2
133. Colicchia M, Campagnolo L, Baldini E, Ulisse S, Valensise H, Moretti C. Molecular basis of thyrotropin and thyroid hormone action during implantation and early development. *Hum Reprod Update* (2014) 20:884–904. doi: 10.1093/humupd/dmu028
134. Stavreus Evers A. Paracrine interactions of thyroid hormones and thyroid stimulation hormone in the female reproductive tract have an impact on female fertility. *Front Endocrinol* (2012) 3:50. doi: 10.3389/fendo.2012.00050
135. Benvenega S, Ferrari SM, Elia G, Ragusa F, Patrizio A, Paparo SR, et al. Nutraceuticals in thyrology: A review of *in vitro*, and *in vivo* animal studies. *Nutrients* (2020) 12:E1337. doi: 10.3390/nu12051337
136. Giuliani C, Noguchi Y, Harii N, Napolitano G, Tatone D, Bucci I, et al. The flavonoid quercetin regulates growth and gene expression in rat FRTL-5 thyroid cells. *Endocrinology* (2008) 149:84–92. doi: 10.1210/en.2007-0618
137. Tran L, Hammuda M, Wood C, Xiao CW. Soy extracts suppressed iodine uptake and stimulated the production of autoimmunogen in rat thyrocytes. *Exp Biol Med Maywood NJ* (2013) 238:623–30. doi: 10.1177/1535370213489488
138. Chandra AK, De N. Goitrogenic/antithyroidal potential of green tea extract in relation to catechin in rats. *Food Chem Toxicol Int J Publ Br Ind Biol Res Assoc* (2010) 48:2304–11. doi: 10.1016/j.fct.2010.05.064
139. Cito G, Cocci A, Micelli E, Gabutti A, Russo GI, Coccia ME, et al. Vitamin d and Male fertility: An updated review. *World J Mens Health* (2020) 38:164–77. doi: 10.5534/wjmh.190057
140. Brenza HL, DeLuca HF. Regulation of 25-hydroxyvitamin D3 1 α -hydroxylase gene expression by parathyroid hormone and 1,25-dihydroxyvitamin D3. *Arch Biochem Biophys* (2000) 381:143–52. doi: 10.1006/abbi.2000.1970
141. Fukumoto S. Phosphate metabolism and vitamin d. *BoneKey Rep* (2014) 3:497. doi: 10.1038/bonekey.2013.231
142. Zanatta L, Zamoner A, Gonçalves R, Zanatta AP, Bouraïma-Lelong H, Bois C, et al. Effect of 1 α ,25-dihydroxyvitamin D3 in plasma membrane targets in immature rat testis: ionic channels and gamma-glutamyl transpeptidase activity. *Arch Biochem Biophys* (2011) 515:46–53. doi: 10.1016/j.abb.2011.09.001
143. Mahmoudi AR, Zarnani AH, Jeddi-Tehrani M, Katouzian L, Tavakoli M, Soltanghoraei H, et al. Distribution of vitamin d receptor and 1 α -hydroxylase in male mouse reproductive tract. *Reprod Sci Thousand Oaks Calif* (2013) 20:426–36. doi: 10.1177/1933719112459235
144. Fu L, Chen Y-H, Xu S, Ji Y-L, Zhang C, Wang H, et al. Vitamin d deficiency impairs testicular development and spermatogenesis in mice. *Reprod Toxicol Elmsford N* (2017) 73:241–9. doi: 10.1016/j.reprotox.2017.06.047
145. Kourtis AP, Read JS, Jamieson DJ. Pregnancy and infection. *N Engl J Med* (2014) 370:2211–8. doi: 10.1056/NEJMr1213566
146. Saito S, Nishikawa K, Morii T, Narita N, Enomoto M, Ichijo M. Expression of activation antigens CD69, HLA-DR, interleukin-2 receptor-alpha (IL-2R alpha) and IL-2R beta on T cells of human decidua at an early stage of pregnancy. *Immunology* (1992) 75:710–2.
147. Watanabe M, Iwatani Y, Kaneda T, Hidaka Y, Mitsuda N, Morimoto Y, et al. Changes in T, b, and NK lymphocyte subsets during and after normal pregnancy. *Am J Reprod Immunol N Y N* 1989 (1997) 37:368–77. doi: 10.1111/j.1600-0897.1997.tb00246.x
148. Li N, Wang L, Cao N, Zhang L, Han X, Yang L. Early pregnancy affects the expression of toll-like receptor pathway in ovine thymus. *Reprod Biol* (2020) 20:547–54. doi: 10.1016/j.repbio.2020.10.003

149. Zhang L, Zhao Z, Mi H, Liu B, Wang B, Yang L. Modulation of helper T cytokines in thymus during early pregnancy in ewes. *Anim Open Access J MDPI* (2019) 9:E245. doi: 10.3390/ani9050245
150. Yang L, Lv W, Liu Y, Chen K, Xue J, Wang Q, et al. Effect of early pregnancy on the expression of prostaglandin synthases in the ovine thymus. *Theriogenology* (2019) 136:166–71. doi: 10.1016/j.theriogenology.2019.06.040
151. Zhang L, Zhao Z, Wang Y, Li N, Cao N, Yang L. Changes in expression of interferon-stimulated genes and ubiquitin activating enzyme E1-like in ovine thymus during early pregnancy. *Anim Reprod* (2020) 17:e20190134. doi: 10.1590/1984-3143-AR2019-0134
152. Wise T. *In vitro* and *in vivo* effects of thymulin on rat testicular steroid synthesis. *J Steroid Biochem Mol Biol* (1998) 66:129–35. doi: 10.1016/s0960-0760(98)00045-4
153. Wise T, Ford JJ. Effects of the thymic peptide thymulin on *in vitro* and *in vivo* testicular steroid concentrations in white composite and meishan boars. *J Anim Sci* (1999) 77:2240–51. doi: 10.2527/1999.7782240x
154. Jacobo P, Guazzone VA, Jarazo-Dietrich S, Theas MS, Lustig L. Differential changes in CD4+ and CD8+ effector and regulatory T lymphocyte subsets in the testis of rats undergoing autoimmune orchitis. *J Reprod Immunol* (2009) 81:44–54. doi: 10.1016/j.jri.2009.04.005
155. Wegmann TG, Lin H, Guilbert L, Mosmann TR. Bidirectional cytokine interactions in the maternal-fetal relationship: is successful pregnancy a Th2 phenomenon? *Immunol Today* (1993) 14:353–6. doi: 10.1016/0167-5699(93)90235-D
156. Wegmann TG. Foetal protection against abortion: is it immunosuppression or immunostimulation? *Ann Immunol* (1984) 135D:309–12. doi: 10.1016/s0769-2625(84)81196-4
157. Szekeres-Bartho J, Wegmann TG. A progesterone-dependent immunomodulatory protein alters the Th1/Th2 balance. *J Reprod Immunol* (1996) 31:81–95. doi: 10.1016/0165-0378(96)00964-3
158. Piccinni MP, Beloni L, Livi C, Maggi E, Scarselli G, Romagnani S. Defective production of both leukemia inhibitory factor and type 2 T-helper cytokines by decidual T cells in unexplained recurrent abortions. *Nat Med* (1998) 4:1020–4. doi: 10.1038/2006
159. Saito S. Cytokine network at the feto-maternal interface. *J Reprod Immunol* (2000) 47:87–103. doi: 10.1016/s0165-0378(00)00060-7
160. Saito S, Sakai M, Sasaki Y, Tanebe K, Tsuda H, Michimata T. Quantitative analysis of peripheral blood Th0, Th1, Th2 and Th1:Th2 cell ratio during normal human pregnancy and preeclampsia. *Clin Exp Immunol* (1999) 117:550–5. doi: 10.1046/j.1365-2249.1999.00997.x
161. Mao H, Gao W, Ma C, Sun J, Liu J, Shao Q, et al. Human placental trophoblasts express the immunosuppressive cytokine IL-35. *Hum Immunol* (2013) 74:872–7. doi: 10.1016/j.humimm.2013.04.010
162. Niedbala W, Wei X-Q, Cai B, Hueber AJ, Leung BP, McInnes IB, et al. IL-35 is a novel cytokine with therapeutic effects against collagen-induced arthritis through the expansion of regulatory T cells and suppression of Th17 cells. *Eur J Immunol* (2007) 37:3021–9. doi: 10.1002/eji.200737810
163. Komorowska B. Autoimmune premature ovarian failure. *Przeegląd Menopauzalny Menopause Rev* (2016) 15:210–4. doi: 10.5114/pm.2016.65666
164. Lin H, Mosmann TR, Guilbert L, Tuntipopipat S, Wegmann TG. Synthesis of T helper 2-type cytokines at the maternal-fetal interface. *J Immunol Baltim Md 1950* (1993) 151:4562–73.
165. Wang W, Sung N, Gilman-Sachs A, Kwak-Kim J. T Helper (Th) cell profiles in pregnancy and recurrent pregnancy losses: Th1/Th2/Th9/Th17/Th22/Th cells. *Front Immunol* (2020) 11:2025. doi: 10.3389/fimmu.2020.02025
166. Kahn DA, Baltimore D. Pregnancy induces a fetal antigen-specific maternal T regulatory cell response that contributes to tolerance. *Proc Natl Acad Sci U.S.A.* (2010) 107:9299–304. doi: 10.1073/pnas.1003909107
167. Shima T, Inada K, Nakashima A, Ushijima A, Ito M, Yoshino O, et al. Paternal antigen-specific proliferating regulatory T cells are increased in uterine-draining lymph nodes just before implantation and in pregnant uterus just after implantation by seminal plasma-priming in allogeneic mouse pregnancy. *J Reprod Immunol* (2015) 108:72–82. doi: 10.1016/j.jri.2015.02.005
168. García L, Hinojosa L, Dominguez R, Chavira R, Rosas P. Effects of infantile thymectomy on ovarian functions and gonadotrophin-induced ovulation in prepubertal mice: role of thymulin. *J Endocrinol* (2000) 166:381–7. doi: 10.1677/joe.0.1660381
169. Hinojosa L, Chavira R, Dominguez R, Rosas P. Effects of thymulin on spontaneous puberty and gonadotrophin-induced ovulation in prepubertal normal and hypothyroid mice. *J Endocrinol* (1999) 163:255–60. doi: 10.1677/joe.0.1630255
170. Brown OA, Sosa YE, Dardenne M, Pléau JM, Goya RG. Studies on the gonadotropin-releasing activity of thymulin: changes with age. *J Gerontol A Biol Sci Med Sci* (2000) 55:B170–176. doi: 10.1093/gerona/55.4.b170
171. Hinojosa L, García L, Dominguez R, Romano MC, Damián-Matsumura PG, Castillo L, et al. Effects of thymulin and GnRH on the release of gonadotropins by *in vitro* pituitary cells obtained from rats in each day of estrous cycle. *Life Sci* (2004) 76:795–804. doi: 10.1016/j.lfs.2004.07.017
172. Duan Y-G, Yu C-F, Novak N, Bieber T, Zhu C-H, Schuppe H-C, et al. Immunodeviation towards a Th17 immune response associated with testicular damage in azoospermic men. *Int J Androl* (2011) 34:e536–545. doi: 10.1111/j.1365-2605.2010.01137.x
173. Wang H, Zhang Y, Fang X, Kwak-Kim J, Wu L. Insulin resistance adversely affect IVF outcomes in lean women without PCOS. *Front Endocrinol* (2021) 12:734638. doi: 10.3389/fendo.2021.734638
174. Mekar K, Oishi S, Akamine K, Kinjo T, Heshiki C, Masamoto H, et al. The effect of insulin resistance on in-vitro fertilization-embryo transfer in women without polycystic ovary syndrome. *Open J Obstet Gynecol* (2016) 6:157–66. doi: 10.4236/ojog.2016.63020
175. Li Y, Lin H, Pan P, Yang D, Zhang Q. Impact of central obesity on women with polycystic ovary syndrome undergoing *In vitro* fertilization. *Biores Open Access* (2018) 7:116–22. doi: 10.1089/biores.2017.0040
176. Issa CM, Abu Khuzam RH. *In vitro* fertilization-induced hypertriglyceridemia with secondary acute pancreatitis and diabetic ketoacidosis. *SAGE Open Med Case Rep* (2017) 5:2050313X16689209. doi: 10.1177/2050313X16689209
177. Vuguin PM, Kedeas MH, Cui L, Guz Y, Gelling RW, Nejjathaim M, et al. Ablation of the glucagon receptor gene increases fetal lethality and produces alterations in islet development and maturation. *Endocrinology* (2006) 147:3995–4006. doi: 10.1210/en.2005-1410
178. Saleh AAEW, Amin EM, Elfallah AA, Hamed AM. Insulin resistance and idiopathic infertility: A potential possible link. *Andrologia* (2020) 52:e13773. doi: 10.1111/and.13773
179. Izzi-Engbeaya C, Jones S, Crustna Y, Machenahalli PC, Papadopoulou D, Modi M, et al. Acute effects of glucagon on reproductive hormone secretion in healthy men. *J Clin Endocrinol Metab* (2020) 105:dgaal64. doi: 10.1210/clinem/dgaa164
180. Sakumoto T, Tokunaga Y, Tanaka H, Nohara M, Motegi E, Shinkawa T, et al. Insulin resistance/hyperinsulinemia and reproductive disorders in infertile women. *Reprod Med Biol* (2010) 9:185–90. doi: 10.1007/s12522-010-0062-5
181. Gill L, Mackey S. Obstetrician-gynecologists' strategies for patient initiation and maintenance of antiobesity treatment with glucagon-like peptide-1 receptor agonists. *J Womens Health* (2021) 30:1016–27. doi: 10.1089/jwh.2020.8683
182. Jensterle M, Janez A, Fliers E, DeVries JH, Vrtacnik-Bokal E, Siegelar SE. The role of glucagon-like peptide-1 in reproduction: from physiology to therapeutic perspective. *Hum Reprod Update* (2019) 25:504–17. doi: 10.1093/humupd/dmz019
183. Eddy JJ, Gideonsen MD, Song JY, Grobman WA, O'Halloran P. Pancreatitis in pregnancy. *Obstet Gynecol* (2008) 112:1075–81. doi: 10.1097/AOG.0b013e318185a032
184. Seo D, Suh H, Lee JK, Jang DK, Kwon HY, Lee CH, et al. Estrogen-induced acute pancreatitis: A case report and literature review. *Obstet Gynecol Sci* (2017) 60:485–9. doi: 10.5468/ogs.2017.60.5.485
185. Badalov N, Baradarjan R, Iswara K, Li J, Steinberg W, Tenner S. Drug-induced acute pancreatitis: an evidence-based review. *Clin Gastroenterol Hepatol Off Clin Pract J Am Gastroenterol Assoc* (2007) 5:648–61. doi: 10.1016/j.cgh.2006.11.023. quiz 644.
186. Gan S-I, Edwards A-L, Symonds C-J, Beck P-L. Hypertriglyceridemia-induced pancreatitis: A case-based review. *World J Gastroenterol* (2006) 12:7197–202. doi: 10.3748/wjg.v12.i44.7197
187. Ouhilal S, Vuguin P, Cui L, Du X-Q, Gelling RW, Reznik SE, et al. Hypoglycemia, hyperglucagonemia, and fetoplacental defects in glucagon receptor knockout mice: a role for glucagon action in pregnancy maintenance. *Am J Physiol Endocrinol Metab* (2012) 302:E522–531. doi: 10.1152/ajpendo.00420.2011
188. Kumar S, Agrawal D, Sharma K, Swain TR. Association of Male infertility to metabolic syndrome and other related disorders. *J Integr Nephrol Androl* (2015) 2:107. doi: 10.4103/2225-1243.168524
189. Csemiczky G, Landgren BM, Collins A. The influence of stress and state anxiety on the outcome of IVF-treatment: psychological and endocrinological assessment of Swedish women entering IVF-treatment. *Acta Obstet Gynecol Scand* (2000) 79:113–8. doi: 10.1034/j.1600-0412.2000.079002113.x
190. Smeenk JMJ, Verhaak CM, Vingerhoets AJJM, Sweep CGJ, Merkus JMW, Willemsen SJ, et al. Stress and outcome success in IVF: the role of self-reports and endocrine variables. *Hum Reprod Oxf Engl* (2005) 20:991–6. doi: 10.1093/humrep/deh739
191. Piquer B, Fonseca JL, Lara HE. Gestational stress, placental norepinephrine transporter and offspring fertility. *Reprod Camb Engl* (2017) 153:147–55. doi: 10.1530/REP-16-0312
192. Kapoor A, Matthews SG. Short periods of prenatal stress affect growth, behaviour and hypothalamic-pituitary-adrenal axis activity in male guinea pig offspring. *J Physiol* (2005) 566:967–77. doi: 10.1113/jphysiol.2005.090191
193. Mayerhofer A, Bartke A, Steger RW. Catecholamine effects on testicular testosterone production in the gonadally active and the gonadally regressed adult golden hamster. *Biol Reprod* (1989) 40:752–61. doi: 10.1095/biolreprod40.4.752
194. Rehman R, Amjad S, Tariq H, Zahid N, Akhter M, Ashraf M. Oxidative stress and male infertility: a cross sectional study. *JPMA J Pak Med Assoc* (2020) 70:461–6. doi: 10.5455/JPMA.12992
195. Whirlledge S, Cidlowski JA. Glucocorticoids, stress, and fertility. *Minerva Endocrinol* (2010) 35:109–25.
196. Ziaja J, Cholewa K, Mazurek U, Cierpka L. [Molecular basics of aldosterone and cortisol synthesis in normal adrenals and adrenocortical adenomas]. *Endokrynol Pol* (2008) 59:330–9.
197. Palomba S, Daolio J, Romeo S, Battaglia FA, Marci R, La Sala GB. Lifestyle and fertility: the influence of stress and quality of life on female fertility. *Reprod Biol Endocrinol RBE* (2018) 16:113. doi: 10.1186/s12958-018-0434-y
198. Pepin É, Dehboneh SS, Raguema N, Esfandarani MT, Lavoie JL. *Role of the renin-angiotensin system in healthy and pathological pregnancies*. IntechOpen (2017). doi: 10.5772/66748
199. Sabbadin C, Andrisani A, Ambrosini G, Bordin L, Donà G, Manso J, et al. Aldosterone in gynecology and its involvement on the risk of hypertension in pregnancy. *Front Endocrinol* (2019) 10:575. doi: 10.3389/fendo.2019.00575

200. Aranceta J, Serra-Majem L, Ribas L, Pérez-Rodrigo C. Breakfast consumption in Spanish children and young people. *Public Health Nutr* (2001) 4:1439–44. doi: 10.1079/phn2001235
201. Spergel DJ. Modulation of gonadotropin-releasing hormone neuron activity and secretion in mice by non-peptide neurotransmitters, gasotransmitters, and gliotransmitters. *Front Endocrinol* (2019) 10:329. doi: 10.3389/fendo.2019.00329
202. Ferlin A, Arredi B, Foresta C. Genetic causes of male infertility. *Reprod Toxicol Elmsford N* (2006) 22:133–41. doi: 10.1016/j.reprotox.2006.04.016
203. Whirlledge S, Cidlowski JA. Glucocorticoids and reproduction: Traffic control on the road to reproduction. *Trends Endocrinol Metab TEM* (2017) 28:399–415. doi: 10.1016/j.tem.2017.02.005
204. Hamilton KJ, Hewitt SC, Arao Y, Korach KS. Estrogen hormone biology. *Curr Top Dev Biol* (2017) 125:109–46. doi: 10.1016/bs.ctdb.2016.12.005
205. Mueller SO, Korach KS. Estrogen receptors and endocrine diseases: lessons from estrogen receptor knockout mice. *Curr Opin Pharmacol* (2001) 1:613–9. doi: 10.1016/s1471-4892(01)00105-9
206. Lu Q, Schnitzler GR, Ueda K, Iyer LK, Diomedea OI, Andrade T, et al. ER alpha rapid signaling is required for estrogen induced proliferation and migration of vascular endothelial cells. *PLoS One* (2016) 11:e0152807. doi: 10.1371/journal.pone.0152807
207. Findlay JK, Liew SH, Simpson ER, Korach KS. Estrogen signaling in the regulation of female reproductive functions. *Handb Exp Pharmacol* (2010) 198:29–35. doi: 10.1007/978-3-642-02062-9_2
208. Palaniappan M, Nguyen L, Grimm SL, Xi Y, Xia Z, Li W, et al. The genomic landscape of estrogen receptor α binding sites in mouse mammary gland. *PLoS One* (2019) 14:e0220311. doi: 10.1371/journal.pone.0220311
209. Stefkovich ML, Arao Y, Hamilton KJ, Korach KS. Experimental models for evaluating non-genomic estrogen signaling. *Steroids* (2018) 133:34–7. doi: 10.1016/j.steroids.2017.11.001
210. Wu S-P, Li R, DeMayo FJ. Progesterone receptor regulation of uterine adaptation for pregnancy. *Trends Endocrinol Metab TEM* (2018) 29:481–91. doi: 10.1016/j.tem.2018.04.001
211. Kumar P, Magon N. Hormones in pregnancy. *Niger Med J J Niger Med Assoc* (2012) 53:179–83. doi: 10.4103/0300-1652.107549
212. Taraborrelli S. Physiology, production and action of progesterone. *Acta Obstet Gynecol Scand* (2015) 94 Suppl 161:8–16. doi: 10.1111/aogs.12771
213. De Vos M, Devroey P, Fauser BCJM. Primary ovarian insufficiency. *Lancet Lond Engl* (2010) 376:911–21. doi: 10.1016/S0140-6736(10)60355-8
214. European Society for Human Reproduction and Embryology (ESHRE) Guideline Group on POI, Webber L, Davies M, Anderson R, Bartlett J, Braat D, et al. ESHRE guideline: management of women with premature ovarian insufficiency. *Hum Reprod Oxf Engl* (2016) 31:926–37. doi: 10.1093/humrep/dew027
215. Rebar RW, Connolly HV. Clinical features of young women with hypergonadotropic amenorrhea. *Fertil Steril* (1990) 53:804–10.
216. Chitayat D, Wyatt PRSOCIETY OF OBSTETRICIANS AND GYNAECOLOGISTS OF CANADA GENETICS COMMITTEE and CANADIAN COLLEGE OF MEDICAL GENETICISTS PRENATAL DIAGNOSIS COMMITTEE. Fragile X testing in obstetrics and gynaecology in Canada. *J Obstet Gynaecol Can JOGC J Obstet Gynecol Can JOGC* (2008) 30:837–41. doi: 10.1016/S1701-2163(16)32949-8
217. Bachelot A, Rouxel A, Massin N, Dulon J, Courtillot C, Matuchansky C, et al. Phenotyping and genetic studies of 357 consecutive patients presenting with premature ovarian failure. *Eur J Endocrinol* (2009) 161:179–87. doi: 10.1530/EJE-09-0231
218. Lakhil B, Braham R, Berguigua R, Bouali N, Zaouali M, Chaieb M, et al. Cytogenetic analyses of premature ovarian failure using karyotyping and interphase fluorescence *in situ* hybridization (FISH) in a group of 1000 patients. *Clin Genet* (2010) 78:181–5. doi: 10.1111/j.1399-0004.2009.01359.x
219. Rudnicka E, Kruszewska J, Klicka K, Kowalczyk J, Grymowicz M, Skórka J, et al. Premature ovarian insufficiency - aetiopathology, epidemiology, and diagnostic evaluation. *Przegląd Menopauzalny Menopause Rev* (2018) 17:105–8. doi: 10.5114/pm.2018.78550
220. Varanasi LC, Subasinghe A, Jayasinghe YL, Callegari ET, Garland SM, Gorelik A, et al. Polycystic ovarian syndrome: Prevalence and impact on the wellbeing of Australian women aged 16–29 years. *Aust N Z J Obstet Gynaecol* (2018) 58:222–33. doi: 10.1111/ajo.12730
221. Rotterdam ESHRE/ASRM-Sponsored PCOS consensus workshop group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Hum Reprod Oxf Engl* (2004) 19:41–7. doi: 10.1093/humrep/deh098
222. Papadakis G, Kandaraki EA, Tseniklidi E, Papanlou O, Diamanti-Kandaraki E. Polycystic ovary syndrome and NC-CAH: Distinct characteristics and common findings. *A Systematic Review. Front Endocrinol* (2019) 10:388. doi: 10.3389/fendo.2019.00388
223. Norman RJ, Teede HJ. A new evidence-based guideline for assessment and management of polycystic ovary syndrome. *Med J Aust* (2018) 209:299–300. doi: 10.5694/mja18.00635
224. Norman RJ, Noakes M, Wu R, Davies MJ, Moran L, Wang JX. Improving reproductive performance in overweight/obese women with effective weight management. *Hum Reprod Update* (2004) 10:267–80. doi: 10.1093/humupd/dmh018
225. Carvalho LML, Dos Reis FM, Candido AL, Nunes FFC, Ferreira CN, Gomes KB. Polycystic ovary syndrome as a systemic disease with multiple molecular pathways: a narrative review. *Endocr Regul* (2018) 52:208–21. doi: 10.2478/enr-2018-0026
226. Richards JS, Ren YA, Candelaria N, Adams JE, Rajkovic A. Ovarian follicular theca cell recruitment, differentiation, and impact on fertility: 2017 update. *Endocr Rev* (2018) 39:1–20. doi: 10.1210/er.2017-00164
227. Rosenfield RL. Current concepts of polycystic ovary syndrome pathogenesis. *Curr Opin Pediatr* (2020) 32:698–706. doi: 10.1097/MOP.0000000000000945
228. Chaudhari N, Dawalbhakta M, Nampoothiri L. GnRH dysregulation in polycystic ovarian syndrome (PCOS) is a manifestation of an altered neurotransmitter profile. *Reprod Biol Endocrinol RBE* (2018) 16:37. doi: 10.1186/s12958-018-0354-x
229. Ashraf S, Nabi M, Rasool S ul A, Rashid F, Amin S. Hyperandrogenism in polycystic ovarian syndrome and role of CYP gene variants: a review. *Egypt J Med Hum Genet* (2019) 20:25. doi: 10.1186/s43042-019-0031-4
230. Teng F, Ma X, Yu X, Yan Y, Zhao J, Gao J, et al. High serum androgen and insulin concentrations increase the tendency of endometrial carcinoma. *J Cancer* (2020) 11:5656–64. doi: 10.7150/jca.46391
231. Rumi MAK, Dhakal P, Kubota K, Chakraborty D, Lei T, Larson MA, et al. Generation of Esr1-knockout rats using zinc finger nuclease-mediated genome editing. *Endocrinology* (2014) 155:1991–9. doi: 10.1210/en.2013-2150
232. Rumi MAK, Singh P, Roby KF, Zhao X, Iqbal K, Ratri A, et al. Defining the role of estrogen receptor β in the regulation of female fertility. *Endocrinology* (2017) 158:2330–43. doi: 10.1210/en.2016-1916
233. Hipp HS, Charen KH, Spencer JB, Allen EG, Sherman SL. Reproductive and gynecologic care of women with fragile X primary ovarian insufficiency (FXPOI). *Menopause N Y N* (2016) 23:993–9. doi: 10.1097/GME.0000000000000658
234. Mohammed K, Abu Dabrh AM, Benkhadra K, Al Nofal A, Carranza Leon BG, Prokop LJ, et al. Oral vs transdermal estrogen therapy and vascular events: A systematic review and meta-analysis. *J Clin Endocrinol Metab* (2015) 100:4012–20. doi: 10.1210/jc.2015-2237
235. Canonico M, Oger E, Plu-Bureau G, Conard J, Meyer G, Lévesque H, et al. Hormone therapy and venous thromboembolism among postmenopausal women: impact of the route of estrogen administration and progestogens: the ESTHER study. *Circulation* (2007) 115:840–5. doi: 10.1161/CIRCULATIONAHA.106.642280
236. Langrish JP, Mills NL, Bath LE, Warner P, Webb DJ, Kelnar CJ, et al. Cardiovascular effects of physiological and standard sex steroid replacement regimens in premature ovarian failure. *Hypertens Dallas Tex 1979* (2009) 53:805–11. doi: 10.1161/HYPERTENSIONAHA.108.126516
237. Piedade KC, Spencer H, Persani L, Nelson LM. Optimizing fertility in primary ovarian insufficiency: Case report and literature review. *Front Genet* (2021) 12:676262. doi: 10.3389/fgene.2021.676262
238. Pinelli S, Artini PG, Basile S, Obino MER, Sergiampietri C, Giannarelli D, et al. Estrogen treatment in infertile women with premature ovarian insufficiency in transitional phase: a retrospective analysis. *J Assist Reprod Genet* (2018) 35:475–82. doi: 10.1007/s10815-017-1096-y
239. Zhang X, Zheng Y, Guo Y, Lai Z. The effect of low carbohydrate diet on polycystic ovary syndrome: A meta-analysis of randomized controlled trials. *Int J Endocrinol* (2019) 2019:4386401. doi: 10.1155/2019/4386401
240. Glintborg D, Altinok ML, Mumm H, Hermann AP, Ravn P, Andersen M. Body composition is improved during 12 months' treatment with metformin alone or combined with oral contraceptives compared with treatment with oral contraceptives in polycystic ovary syndrome. *J Clin Endocrinol Metab* (2014) 99:2584–91. doi: 10.1210/jc.2014-1135
241. Takasaki A, Tamura I, Okada-Hayashi M, Orita T, Tanabe M, Maruyama S, et al. Usefulness of intermittent clomiphene citrate treatment for women with polycystic ovarian syndrome that is resistant to standard clomiphene citrate treatment. *Reprod Med Biol* (2018) 17:454–8. doi: 10.1002/rmb2.12219
242. Homburg R, Hendriks ML, König TE, Anderson RA, Balen AH, Brincat M, et al. Clomifene citrate or low-dose FSH for the first-line treatment of infertile women with anovulation associated with polycystic ovary syndrome: a prospective randomized multinational study. *Hum Reprod Oxf Engl* (2012) 27:468–73. doi: 10.1093/humrep/der401
243. Waanbah BD, Joseph T, Rebekah G, Kunjummen AT, Kamath MS. Letrozole as first-line drug for ovulation induction in treatment-naïve infertile polycystic ovarian syndrome women. *J Obstet Gynaecol Res* (2021) 47:3583–9. doi: 10.1111/jog.14926
244. Ganie MA, Khurana ML, Nisar S, Shah PA, Shah ZA, Kulshrestha B, et al. Improved efficacy of low-dose spironolactone and metformin combination than either drug alone in the management of women with polycystic ovary syndrome (PCOS): a six-month, open-label randomized study. *J Clin Endocrinol Metab* (2013) 98:3599–607. doi: 10.1210/jc.2013-1040
245. Glintborg D, Mumm H, Holst JJ, Andersen M. Effect of oral contraceptives and/or metformin on GLP-1 secretion and reactive hypoglycaemia in polycystic ovary syndrome. *Endocr Connect* (2017) 6:267–77. doi: 10.1530/EC-17-0034
246. Devin JK, Nian H, Celedonio JE, Wright P, Brown NJ. Sitagliptin decreases visceral fat and blood glucose in women with polycystic ovarian syndrome. *J Clin Endocrinol Metab* (2020) 105:dgz028. doi: 10.1210/clinem/dgz028
247. Javed Z, Papageorgiou M, Deshmukh H, Rigby AS, Qamar U, Abbas J, et al. Effects of empagliflozin on metabolic parameters in polycystic ovary syndrome: A randomized controlled study. *Clin Endocrinol (Oxf)* (2019) 90:805–13. doi: 10.1111/cen.13968
248. Benito E, Gómez-Martin JM, Vega-Piñero B, Priego P, Galindo J, Escobar-Morreale HF, et al. Fertility and pregnancy outcomes in women with polycystic ovary syndrome following bariatric surgery. *J Clin Endocrinol Metab* (2020) 105:dgaa439. doi: 10.1210/clinem/dgaa439

249. Toulis KA, Goulis DG, Kolibianakis EM, Venetis CA, Tarlatzis BC, Papadimas I. Risk of gestational diabetes mellitus in women with polycystic ovary syndrome: a systematic review and a meta-analysis. *Fertil Steril* (2009) 92:667–77. doi: 10.1016/j.fertnstert.2008.06.045
250. De Frène V, Vansteelandt S, T'Sjoen G, Gerris J, Somers S, Vercruyse L, et al. A retrospective study of the pregnancy, delivery and neonatal outcome in overweight versus normal weight women with polycystic ovary syndrome. *Hum Reprod Oxf Engl* (2014) 29:2333–8. doi: 10.1093/humrep/deu154
251. Wierman ME, Basson R, Davis SR, Khosla S, Miller KK, Rosner W, et al. Androgen therapy in women: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* (2006) 91:3697–710. doi: 10.1210/jc.2006-1121
252. Sullivan SD, Sarrel PM, Nelson LM. Hormone replacement therapy in young women with primary ovarian insufficiency and early menopause. *Fertil Steril* (2016) 106:1588–99. doi: 10.1016/j.fertnstert.2016.09.046
253. Renoux C, Dell'aniello S, Garbe E, Suissa S. Transdermal and oral hormone replacement therapy and the risk of stroke: a nested case-control study. *BMJ* (2010) 340:c2519. doi: 10.1136/bmj.c2519
254. Talaulikar VS, Arulkumar S. Reproductive outcomes after assisted conception. *Obstet Gynecol Surv* (2012) 67:566–83. doi: 10.1097/OGX.0b013e31826a5d4a
255. Holloch KJ, Savaris RF, Forstein DA, Miller PB, Higdon HL, Likes CE, et al. Coexistence of polycystic ovary syndrome and endometriosis in women with infertility. *J Endometr Pelvic Pain Disord* (2014) 6:79–83. doi: 10.5301/je.5000181
256. Niafar M, Pourafkari L, Porhomayon J, Nader N. A systematic review of GLP-1 agonists on the metabolic syndrome in women with polycystic ovaries. *Arch Gynecol Obstet* (2016) 293:509–15. doi: 10.1007/s00404-015-3976-7
257. Titi-Lartey OA, Khan YS. *Embryology, testicle*. Treasure Island (FL: StatPearls Publishing (2022)). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK557763/>.
258. Amory JK, Bremner WJ. Regulation of testicular function in men: implications for male hormonal contraceptive development. *J Steroid Biochem Mol Biol* (2003) 85:357–61. doi: 10.1016/s0960-0760(03)00205-x
259. Tanaka T, Kanatsu-Shinohara M, Lei Z, Rao CV, Shinohara T. The luteinizing hormone-testosterone pathway regulates mouse spermatogonial stem cell self-renewal by suppressing WNT5A expression in sertoli cells. *Stem Cell Rep* (2016) 7:279–91. doi: 10.1016/j.stemcr.2016.07.005
260. Tyagi V, Scordo M, Yoon RS, Liporace FA, Greene LW. Revisiting the role of testosterone: Are we missing something? *Rev Urol* (2017) 19:16–24. doi: 10.3909/riu0716
261. Walker WH. Testosterone signaling and the regulation of spermatogenesis. *Spermatogenesis* (2011) 1:116–20. doi: 10.4161/spmg.1.2.16956
262. O'Shaughnessy PJ, Monteiro A, Verhoeven G, De Gendt K, Abel MH. Effect of FSH on testicular morphology and spermatogenesis in gonadotrophin-deficient hypogonadal mice lacking androgen receptors. *Reprod Camb Engl* (2010) 139:177–84. doi: 10.1530/REP-09-0377
263. Nottelmann ED, Susman EJ, Inoff-Germain G, Cutler GB, Loriaux DL, Chrousos GP. Developmental processes in early adolescence: relationships between adolescent adjustment problems and chronologic age, pubertal stage, and puberty-related serum hormone levels. *J Pediatr* (1987) 110:473–80. doi: 10.1016/s0022-3476(87)80521-8
264. Rowe R, Maughan B, Worthman CM, Costello EJ, Angold A. Testosterone, antisocial behavior, and social dominance in boys: pubertal development and biosocial interaction. *Biol Psychiatry* (2004) 55:546–52. doi: 10.1016/j.biopsych.2003.10.010
265. Turcu A, Smith JM, Auchus R, Rainey WE. Adrenal androgens and androgen precursors—definition, synthesis, regulation and physiologic actions. *Compr Physiol* (2014) 4:1369–81. doi: 10.1002/cphy.c140006
266. de Menezes KJ, Peixoto C, Nardi AE, Carta MG, Machado S, Veras AB. Dehydroepiandrosterone, its sulfate and cognitive functions. *Clin Pract Epidemiol Ment Health CP EMH* (2016) 12:24–37. doi: 10.2174/1745017901612010024
267. Gray PB, McHale TS, Carré JM. A review of human male field studies of hormones and behavioral reproductive effort. *Horm Behav* (2017) 91:52–67. doi: 10.1016/j.yhbeh.2016.07.004
268. Verhoeven G, Willems A, Denolet E, Swinnen JV, De Gendt K. Androgens and spermatogenesis: lessons from transgenic mouse models. *Philos Trans R Soc Lond B Biol Sci* (2010) 365:1537–56. doi: 10.1098/rstb.2009.0117
269. Mohanty G, Jena SR, Kar S, Samanta L. Paternal factors in recurrent pregnancy loss: an insight through analysis of non-synonymous single-nucleotide polymorphism in human testis-specific chaperone HSPA2 gene. *Environ Sci Pollut Res Int* (2021) 29:62219–34. doi: 10.1007/s11356-021-17799-3
270. Choobineh H, Kazemi M, Sadighi Gilani MA, Heydari T, Shokri S, Bazrafkan M, et al. Testosterone reduces spinal cord injury-induced effects on Male reproduction by preventing CADM1 defect. *Cell J* (2018) 20:138–49. doi: 10.22074/cellj.2018.5003
271. Snyder E, Chukrallah L, Seltzer K, Goodwin L, Braun RE. ADAD1 and ADAD2, testis-specific adenosine deaminase domain-containing proteins, are required for male fertility. *Sci Rep* (2020) 10:11536. doi: 10.1038/s41598-020-67834-5
272. Chi X, Luo W, Song J, Li B, Su T, Yu M, et al. Kindlin-2 in sertoli cells is essential for testis development and male fertility in mice. *Cell Death Dis* (2021) 12:604. doi: 10.1038/s41419-021-03885-4
273. Wei Y, Gao Q, Niu P, Xu K, Qiu Y, Hu Y, et al. Integrative proteomic and phosphoproteomic profiling of testis from Wip1 phosphatase-knockout mice: Insights into mechanisms of reduced fertility. *Mol Cell Proteomics MCP* (2019) 18:216–30. doi: 10.1074/mcp.RA117.000479
274. Xiang F, Cui B, Gao Q, Zhang J, Zhang J, Li W. Decreased levels of Ca²⁺-calmodulin-dependent protein kinase IV in the testis as a contributing factor to reduced fertility in male Crybb2^{-/-} mice. *Int J Mol Med* (2012) 30:1145–51. doi: 10.3892/ijmm.2012.1116
275. Sun J, Lu Y, Nozawa K, Xu Z, Morohoshi A, Castaneda JM, et al. CRISPR/Cas9-based genome editing in mice uncovers 13 testis- or epididymis-enriched genes individually dispensable for male reproduction†. *Biol Reprod* (2020) 103:183–94. doi: 10.1093/biolre/ioaa083
276. Xia M, Xia J, Niu C, Zhong Y, Ge T, Ding Y, et al. Testis-expressed protein 33 is not essential for spermiogenesis and fertility in mice. *Mol Med Rep* (2021) 23:317. doi: 10.3892/mmr.2021.11956
277. Shah B, Khan R, Shah W, Aftab A, Khan M, Dil S, et al. Inactivation of testis-specific gene C4orf46 is dispensable for spermatogenesis and fertility in mouse. *Mamm Genome Off J Int Mamm Genome Soc* (2021) 32:364–70. doi: 10.1007/s00335-021-09879-z
278. Holcomb RJ, Oura S, Nozawa K, Kent K, Yu Z, Robertson MJ, et al. The testis-specific serine proteases PRSS44, PRSS46, and PRSS54 are dispensable for male mouse fertility†. *Biol Reprod* (2020) 102:84–91. doi: 10.1093/biolre/ioz158
279. He X, Xie W, Li H, Cui Y, Wang Y, Guo X, et al. The testis-specifically expressed gene Trim69 is not essential for fertility in mice. *J BioMed Res* (2020) 35:47–60. doi: 10.7555/JBR.34.20200069
280. Jia X, Liu Z, Lu X, Tang J, Wu Y, Du Q, et al. Effects of MCLR exposure on sex hormone synthesis and reproduction-related genes expression of testis in male rana nigromaculata. *Environ Pollut Barking Essex 1987* (2018) 236:12–20. doi: 10.1016/j.envpol.2018.01.057
281. Yang X, Zhang X, Yao Q, Song M, Han Y, Shao B, et al. T-2 toxin impairs male fertility by disrupting hypothalamic-pituitary-testis axis and declining testicular function in mice. *Chemosphere* (2019) 234:909–16. doi: 10.1016/j.chemosphere.2019.06.145
282. Zhang X, Yue Z, Zhang H, Liu L, Zhou X. Repeated administrations of Mn3O4 nanoparticles cause testis damage and fertility decrease through PPAR-signaling pathway. *Nanotoxicology* (2020) 14:326–40. doi: 10.1080/17435390.2019.1695976
283. Domínguez-Salazar E, Hurtado-Alvarado G, Medina-Flores F, Dorantes J, González-Flores O, Contis-Montes de Oca A, et al. Chronic sleep loss disrupts blood-testis and blood-epididymis barriers, and reduces male fertility. *J Sleep Res* (2020) 29:e12907. doi: 10.1111/jsr.12907
284. Ye J, Luo D, Xu X, Sun M, Su X, Tian Z, et al. Metformin improves fertility in obese males by alleviating oxidative stress-induced blood-testis barrier damage. *Oxid Med Cell Longev* (2019) 2019:9151067. doi: 10.1155/2019/9151067
285. Akomolafe SF, Odeniyi IA, Oyetayo FL, Ajayi OB. African Star apple fruit pulp-supplemented diet modulates fertility-related biomolecules in the testis and epididymis of high-fat diet/streptozotocin-induced diabetic rats. *J Food Biochem* (2019) 43:e12969. doi: 10.1111/jfbc.12969
286. Silva JV, Yoon S, Domingues S, Guimarães S, Goltsev AV, da Cruz E Silva EF, et al. Amyloid precursor protein interaction network in human testis: sentinel proteins for male reproduction. *BMC Bioinf* (2015) 16:12. doi: 10.1186/s12859-014-0432-9
287. Miyata H, Castaneda JM, Fujihara Y, Yu Z, Archambeault DR, Isotani A, et al. Genome engineering uncovers 54 evolutionarily conserved and testis-enriched genes that are not required for male fertility in mice. *Proc Natl Acad Sci U.S.A.* (2016) 113:7704–10. doi: 10.1073/pnas.1608458113
288. Lu Y, Oura S, Matsumura T, Oji A, Sakurai N, Fujihara Y, et al. CRISPR/Cas9-mediated genome editing reveals 30 testis-enriched genes dispensable for male fertility in mice†. *Biol Reprod* (2019) 101:501–11. doi: 10.1093/biolre/iox103
289. Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *Reprod Biol Endocrinol RBE* (2015) 13:37. doi: 10.1186/s12958-015-0032-1
290. Kurkowska W, Bogacz A, Janiszewska M, Gabryś E, Tiszler M, Bellanti F, et al. Oxidative stress is associated with reduced sperm motility in normal semen. *Am J Mens Health* (2020) 14:1557988320939731. doi: 10.1177/1557988320939731
291. Kumar N, Singh AK. Trends of male factor infertility, an important cause of infertility: A review of literature. *J Hum Reprod Sci* (2015) 8:191–6. doi: 10.4103/0974-1208.170370
292. Tiepolo L, Zuffardi O. Localization of factors controlling spermatogenesis in the nonfluorescent portion of the human y chromosome long arm. *Hum Genet* (1976) 34:119–24. doi: 10.1007/BF00278879
293. Sharma N, Singh M, Acharya N, Singh SK, Thapa BR, Kaur G, et al. Implication of the cystic fibrosis transmembrane conductance regulator gene in infertile family members of Indian CF patients. *Biochem Genet* (2008) 46:847–56. doi: 10.1007/s10528-008-9199-x
294. Liu K, Zhao R, Shen M, Ye J, Li X, Huang Y, et al. Role of genetic mutations in folate-related enzyme genes on Male infertility. *Sci Rep* (2015) 5:15548. doi: 10.1038/srep15548
295. Xiao Q, Wang L, Supekar S, Shen T, Liu H, Ye F, et al. Structure of human steroid 5 α -reductase 2 with anti-androgen drug finasteride. *Res Sq* (2020), rs.3.rs-40159. doi: 10.21203/rs.3.rs-40159/v1
296. Wang T, Gao H, Li W, Liu C. Essential role of histone replacement and modifications in Male fertility. *Front Genet* (2019) 10:962. doi: 10.3389/fgene.2019.00962
297. Choy JT, Eisenberg ML. Male Infertility as a window to health. *Fertil Steril* (2018) 110:810–4. doi: 10.1016/j.fertnstert.2018.08.015
298. Aitken RJ, Roman SD. Antioxidant systems and oxidative stress in the testes. *Oxid Med Cell Longev* (2008) 1:15–24. doi: 10.4161/oxim.1.1.6843
299. Torres-Arce E, Vizmanos B, Babio N, Márquez-Sandoval F, Salas-Huetos A. Dietary antioxidants in the treatment of Male infertility: Counteracting oxidative stress. *Biology* (2021) 10:241. doi: 10.3390/biology10030241

300. Di Guardo F, Vloeberghs V, Bardhi E, Blockeel C, Verheyen G, Tournaye H, et al. Low testosterone and semen parameters in Male partners of infertile couples undergoing IVF with a total sperm count greater than 5 million. *J Clin Med* (2020) 9:E3824. doi: 10.3390/jcm9123824
301. Crosnoe LE, Grober E, Ohl D, Kim ED. Exogenous testosterone: a preventable cause of male infertility. *Transl Androl Urol* (2013) 2:106–13. doi: 10.3978/j.issn.2223-4683.2013.06.01
302. Shiming Z, Mak K-K, Balijepalli MK, Chakravarthi S, Pichika MR. Swietenine potentiates the antihyperglycemic and antioxidant activity of metformin in streptozotocin induced diabetic rats. *BioMed Pharmacother Biomedecine Pharmacother* (2021) 139:111576. doi: 10.1016/j.biopha.2021.111576
303. Sjaarda LA, Mumford SL, Kuhr DL, Holland TL, Silver RM, Plowden TC, et al. Association of testosterone and antimüllerian hormone with time to pregnancy and pregnancy loss in fecund women attempting pregnancy. *Fertil Steril* (2018) 109:540–548.e1. doi: 10.1016/j.fertnstert.2017.11.014
304. Lathi RB, Dahan MH, Reynolds-May MF, Milki AA, Behr B, Westphal LM. The role of serum testosterone in early pregnancy outcome: a comparison in women with and without polycystic ovary syndrome. *J Obstet Gynaecol Can JOGC J Obstet Gynecol Can JOGC* (2014) 36:811–6. doi: 10.1016/S1701-2163(15)30483-7
305. Valdimarsdóttir R, Wikström A-K, Kallak TK, Elenis E, Axelsson O, Preissl H, et al. Pregnancy outcome in women with polycystic ovary syndrome in relation to second-trimester testosterone levels. *Reprod BioMed Online* (2021) 42:217–25. doi: 10.1016/j.rbmo.2020.09.019
306. Chinnathambi V, Blesson CS, Vincent KL, Saade GR, Hankins GD, Yallampalli C, et al. Elevated testosterone levels during rat pregnancy cause hypersensitivity to angiotensin II and attenuation of endothelium-dependent vasodilation in uterine arteries. *Hypertens Dallas Tex 1979* (2014) 64:405–14. doi: 10.1161/HYPERTENSIONAHA.114.03283
307. Sun B, Wang F, Sun J, Yu W, Sun Y. Basal serum testosterone levels correlate with ovarian response but do not predict pregnancy outcome in non-PCOS women undergoing IVF. *J Assist Reprod Genet* (2014) 31:829–35. doi: 10.1007/s10815-014-0246-8
308. Chen Z, Zhang D, Sun Z, Yu Q. A proper increasing in the testosterone level may be associated with better pregnancy outcomes for patients with tubal or Male infertility during *in vitro* Fertilization/Intracytoplasmic sperm injection. *Front Physiol* (2021) 12:696854. doi: 10.3389/fphys.2021.696854
309. Saharkhiz N, Zademoares S, Salehpour S, Hosseini S, Nazari L, Tehrani HG. The effect of testosterone gel on fertility outcomes in women with a poor response in *in vitro* fertilization cycles: A pilot randomized clinical trial. *J Res Med Sci Off J Isfahan Univ Med Sci* (2018) 23:3. doi: 10.4103/jrms.JRMS_864_17
310. Burger HG. Androgen production in women. *Fertil Steril* (2002) 77 Suppl 4:S3–5. doi: 10.1016/s0015-0282(02)02985-0
311. Sowers MF, Beebe JL, McConnell D, Randolph J, Jannausch M. Testosterone concentrations in women aged 25–50 years: associations with lifestyle, body composition, and ovarian status. *Am J Epidemiol* (2001) 153:256–64. doi: 10.1093/aje/153.3.256
312. Hirschberg AL. Female hyperandrogenism and elite sport. *Endocr Connect* (2020) 9:R81–92. doi: 10.1530/EC-19-0537