



OPEN ACCESS

EDITED AND REVIEWED BY
Jiali Zeng,
Syracuse University, Syracuse, United States

*CORRESPONDENCE

Haibo Xu
✉ xuhaibo@whu.edu.cn
Zhe Dai
✉ daizhe@znhospital.cn

[†]These authors have contributed equally to this work

RECEIVED 20 October 2024
ACCEPTED 29 October 2024
PUBLISHED 20 November 2024

CITATION

Deng M, Li Z, Chen S, Wang H, Sun L, Tang J, Luo L, Zhang X, Xu H and Dai Z (2024) Corrigendum: Exploring the heterogeneity of hepatic and pancreatic fat deposition in obesity: implications for metabolic health. *Front. Endocrinol.* 15:1514364. doi: 10.3389/fendo.2024.1514364

COPYRIGHT

© 2024 Deng, Li, Chen, Wang, Sun, Tang, Luo, Zhang, Xu and Dai. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Exploring the heterogeneity of hepatic and pancreatic fat deposition in obesity: implications for metabolic health

Ming Deng^{1,2,3†}, Zhen Li^{4†}, Shangyu Chen^{5,6†}, Huawei Wang⁵, Li Sun⁵, Jun Tang⁵, Liman Luo⁵, Xiaoxiao Zhang⁷, Haibo Xu^{1,2,3*} and Zhe Dai^{5,8*}

¹Department of Radiology, Zhongnan Hospital of Wuhan University, Wuhan University, Wuhan, China,

²Hubei Provincial Engineering Research Center of Multimodal Medical Imaging Technology and Clinical Application, Wuhan, China, ³Wuhan Clinical Research and Development Center of Brain Resuscitation and Functional Imaging, Wuhan, China, ⁴Department of Hepatobiliary and Pancreatic Surgery, Zhongnan Hospital of Wuhan University, Wuhan University, Wuhan, China, ⁵Department of Endocrinology, Zhongnan Hospital of Wuhan University, Wuhan University, Wuhan, China, ⁶Department of Endocrinology and Metabolism, Guangxi Academy of Medical Sciences and the People's Hospital of Guangxi Zhuang Autonomous Region, Guangxi, Nanning, China, ⁷Department of MSC Clinical & Technical Solutions, Philips Healthcare, Beijing, China, ⁸Department of Clinical Nutrition, Zhongnan Hospital of Wuhan University, Wuhan University, Wuhan, China

KEYWORDS

obesity, non-alcoholic fatty liver disease, non-alcoholic fatty pancreas disease, insulin resistance, fat deposition heterogeneity

A Corrigendum on

Exploring the heterogeneity of hepatic and pancreatic fat deposition in obesity: implications for metabolic health


By Deng M, Li Z, Chen S, Wang H, Sun L, Tang J, Luo L, Zhang X, Xu H and Dai Z (2024). *Front. Endocrinol.* 15:1447750. doi: 10.3389/fendo.2024.1447750

In the published article, there was an error in the legend for **Figure 1** as published. In the original article, there was a misplacement of data in the text description for **Figure 1** as published. The corrected caption and its figure appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

**FIGURE 1**

Four categories of MR images illustrate different levels of fat deposition in obesity. **(A)** Represents the first category, showing no fat deposition in either the liver or pancreas. The average fat fraction (FF) of the liver and pancreas were 4.37% and 1%, respectively. **(B)** Represents the second category, with high-fat deposition in the liver and minimal fat deposition in the pancreas. The average FF of the liver and pancreas were 28% and 4%, respectively. **(C)** Illustrates the third category, with less fat deposition in the liver but higher fat deposition in the pancreas. The average FF of the liver and pancreas were 3% and 13.1%, respectively. **(D)** Demonstrates the fourth category, characterized by fatty deposits in both the liver and pancreas. The average FF of the liver and pancreas were 31% and 22.6%, respectively.