



OPEN ACCESS

EDITED AND REVIEWED BY
Terry Francis Davies,
Icahn School of Medicine at Mount Sinai,
United States

*CORRESPONDENCE

Aldo Ferreira-Hermosillo
✉ aldo.nagisa@gmail.com

RECEIVED 08 November 2024
ACCEPTED 13 December 2024
PUBLISHED 10 January 2025

CITATION

Ferreira-Hermosillo A, Toledo JO and
Cordoba K (2025) Corrigendum: Evaluation of
the risk of hypothyroidism and its clinical
manifestations using the Zulewski scale.
Front. Endocrinol. 15:1525162.
doi: 10.3389/fendo.2024.1525162

COPYRIGHT

© 2025 Ferreira-Hermosillo, Toledo and
Cordoba. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Corrigendum: Evaluation of the risk of hypothyroidism and its clinical manifestations using the Zulewski scale

Aldo Ferreira-Hermosillo^{1*}, Juan Omar Toledo²
and Karla Cordoba²

¹Unidad de Investigación Médica en Enfermedades Endocrinas, Centro Médico Nacional Siglo XXI,
Instituto Mexicano del Seguro Social, Mexico City, Mexico, ²Medical Affairs México, MERCK,
Naucalpan de Juárez, Mexico

KEYWORDS

hypothyroidism, population, Zulewski scale, screening, e-health

A Corrigendum on

**Evaluation of the risk of hypothyroidism and its clinical manifestations
using the Zulewski scale**

By Ferreira-Hermosillo A, Toledo JO and Cordoba K (2024) *Front. Endocrinol.* 15:1416663.
doi: 10.3389/fendo.2024.1416663

In the published article, there was an error in **Table 2** as published. In the column ≤ 45 years ($n = 25,159$), the percentage of “Tingling or burning sensation” and “weight gain” were displayed as “4.5%”. The correct percentage is “45%”. In the column of Men ($n = 3868$), the percentage of “Constipation” was displayed as “3.4%” and the correct percentage is “34%”. The corrected **Table 2** and its caption Comparison of the prevalence of symptoms and signs of hypothyroidism according to sex and age group in the study participants appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

TABLE 2 Comparison of the prevalence of symptoms and signs of hypothyroidism according to sex and age group in the study participants.

	Women (n=27581)	Men (n=3868)	<i>p-value</i>	≤45 years (n=25,159)	>45 years (n=6290)	<i>p-value</i>
Risk						
No/low	24%	50%	<0.001	25%	31%	<0.001
Mediate	38%	32%		39%	32%	
High	38%	18%		36%	37%	
Symptoms						
Little sweating	40%	30%	<0.001	40%	37%	<0.001
Need to clear throat	41%	38%	<0.001	41%	42%	<0.001
Tingling or burning sensation	47%	33%	<0.001	45%	44%	<0.001
Dry Skin	49%	32%	<0.001	47%	48%	<0.001
Constipation	64%	34%	<0.001	62%	54.5%	<0.001
Decreased hearing capacity	22%	20%	<0.001	20%	30%	<0.001
Weight gain	47%	21%	<0.001	45%	41%	<0.001
Signs						
Slow movements	40%	28%	<0.001	36%	47%	<0.001
Slow reaction speed	29%	19%	<0.001	27%	31%	<0.001
Skin thickening	23%	18%	<0.001	23%	20%	<0.001
Facial edema	31%	14%	<0.001	30%	26%	<0.001
Cold intolerance	44%	22%	<0.001	44%	33%	<0.001