



Policy Recommendation for the Conservation of the Suweon Treefrog (*Dryophytes suweonensis*) in the Republic of Korea

Amaël Borzée* and Yikweon Jang

Division of EcoScience and Department of Life Science, Ewha Womans University, Seoul, South Korea

OPEN ACCESS

Edited by:

Franco Andreone,
Museo Regionale di Scienze Naturali,
Italy

Reviewed by:

Nicolas Urbina-Cardona,
Pontificia Universidad Javeriana,
Colombia
Attila D. Sándor,
University of Agricultural Sciences and
Veterinary Medicine of Cluj-Napoca,
Romania

*Correspondence:

Amaël Borzée
amaelborzee@gmail.com

Specialty section:

This article was submitted to
Conservation,
a section of the journal
Frontiers in Environmental Science

Received: 27 September 2018

Accepted: 13 March 2019

Published: 04 April 2019

Citation:

Borzée A and Jang Y (2019) Policy
Recommendation for the
Conservation of the Suweon Treefrog
(*Dryophytes suweonensis*) in the
Republic of Korea.
Front. Environ. Sci. 7:39.
doi: 10.3389/fenvs.2019.00039

The Suweon treefrog (*Dryophytes suweonensis*) is an endangered species with an important evolutionary significance. However, the current rate of decline projects the extinction of the species within a few decades in the Republic of Korea. The species is presently exclusively inhabiting rice paddies and is not present in any protected area, although it would be relatively easy to designate protected area under the RAMSAR convention and correct the current negative population dynamics. Through three policy recommendations, we present the points of importance for the conservation of the Suweon Treefrog, list the site of importance for its conservation, and introduce an agricultural model allowing for the conservation of the species as well as continued rice production and economic income.

Actionable recommendations:

- Establishment of protected areas within the range of *D. suweonensis*
- Pesticide and herbicide ban in rice-paddy complexes where *D. suweonensis* occurs
- Vegetation mowing limit for surroundings of rice paddies where *D. suweonensis* occurs.

Keywords: conservation, policy, Suweon Treefrog, Republic of Korea, *Dryophytes suweonensis*

INTRODUCTION

The Suweon Treefrog (*Dryophytes [Hyla] suweonensis*; Duellman et al., 2016) is an endangered species in the Republic of Korea (IUCN, 2018) due to a low extent of occurrence (Borzée et al., 2017a), fragmented population (Borzée, 2018) and continuing decline in area of occurrence (Borzée et al., 2018a). About 2,500 calling males are estimated to be present in the Republic of Korea (Borzée et al., 2018a), segregated into 14 disconnected areas (Borzée, 2018). The species is also known to occur at a minimum of three sites in the Democratic People's Republic of Korea, with at least one seemingly healthy sub-population (Chun et al., 2012; Borzée and Seliger, 2018); although population sizes are not available for the country. Under the current threat levels, the species could become extinct within decades (Borzée, 2018).

In the Republic of Korea the species ranges from the Mangyeong River in the south to the Imjin River in the north (Kim et al., 2012; Borzée et al., 2016c, 2017a). Population sizes are generally decreasing because of habitat loss through encroachment and urbanization (Lee and Miller-Rushing, 2014), although some rare populations displayed low increases between 2015 and 2017 (Borzée et al., 2018a). Despite landscape modifications being the first reason for the endangered status of *D. suweonensis* (Borzée and Jang, 2015; Borzée, 2018),

the species is under multiple threats including hybridization (Kuramoto, 1984; Borzée and Jang, 2018b), competition with *D. japonicus* (Borzée et al., 2016a,b, 2018e; Borzée and Jang, 2018a), environmental pollution (Borzée et al., 2018d), crop management and harvest (Borzée and Jang, 2017; Borzée et al., 2018b), emergent diseases such as the Chytrid fungus (Borzée et al., 2017b), and invasive species (Kim, 2016; Borzée et al., 2017b), among the main and defined threats.

It is important to protect the species due to its clear evolutionary potential, differentiated from other Hylid species by a switch in heterogamety resulting in a ZW system (Dufresnes et al., 2015), and in regard to the international conservation crisis involving amphibians worldwide (Stuart et al., 2004; Beebee and Griffiths, 2005; Bishop et al., 2012; Wake, 2012; Howard and Bickford, 2014; Pimm et al., 2014). Despite the current situation, *D. suweonensis* is not present in any protected area (Borzée et al., 2017a) and is only known to breed in rice paddies (Borzée and Jang, 2015), while original potential habitats might have included wetlands with low vegetation including *Alopecurus* sp. and *Typha* spp. (Borzée and Jang, 2015). The only conservation project to reintroduce the species was unsuccessful (Borzée et al., 2018c) and despite plans for *ex-situ* breeding facilities having been drafted (Noh, 2017), conservation of populations in the wild would simplify the protection of the species (Snyder et al., 1996). Additionally, the species (Borzée et al., 2018b) and other anuran species can adequately take advantage of the artificial flooding of rice paddies for agricultural purposes (Fujioka and Lane, 1997; Machado and Maltchik, 2010; Naito et al., 2013; Holzer, 2014; Orchard et al., 2019).

Amphibian conservation plans benefit from articulating local policies around the prioritized actions of the Amphibian Conservation Action Plan (Wren et al., 2015). Out of the 12 chapters of the ACAP, some sections on habitat protection and ecotoxicology are relevant to the policy recommendations recommended here. Regarding habitat protection, the ACAP recommends first to refine the definition of “Critical Habitat” for the target species and develop a list of standards for the identification of important sites. In the long term, the ACAP recommends developing “Amphibian Critical Habitat” keys to identify priority sites. Regarding ecotoxicology, short term recommendations include identifying organic agriculture movements to explore potential development and working with authorities to change legislation. They also include developing outreach material on the effects of locally applied pollutants on amphibian and human health at the stakeholder level, here rice farmers. Long term recommendations include long-term partnerships with organic agriculture organizations and the social marketing of the issue through the establishment of links with grassroots organizations. The policy recommendations made here are in-line with these global recommendations.

POLICY AND PRACTICE

Consequently, we recommend three policies for the *in-situ* conservation of the Suweon Treefrog (*D. suweonensis*). These

policies are independent of each other and need not be implemented in the order presented here.

Designation of Protected Areas Within the Range of the Species

We strongly urge the preservation of the current rice paddy system and the continued agricultural practices by farmers such as current timing of flooding, tilling and planting (Borzée et al., 2018b). However, avoiding burning straws left-over from the harvest before the fallow phase would avoid lethal consequences to individuals brumating in the vicinity of the rice paddies (Borzée et al., 2019). The designation of agricultural wetlands as protected area and their continued harvest may be conducted under the guidelines stipulated in the RAMSAR convention (see Classification of Wetland Type, §4.3.4; Ramsar Convention Secretariat, 2013) and governmental funding following those provided for “agri-environment” support programmes in the UK, where funding is matched to a score based on defined environmental benefits (Baker et al., 2011). This recommendation is based on the fact that the breeding activity of *D. suweonensis* is synchronized with the regulated hydroperiod resulting from the flooding of rice paddies for rice cultivation (Borzée et al., 2018b), which has so far resulted in adequate population recruitment. The transformation of rice paddies into natural wetlands is not recommended at the moment as the data available on the natural requirements of the species are still limited and transformations may result in the local extirpation of the species if conducted inadequately. The knowledge currently available only includes partial vegetation data, habitat size, and distribution models (Borzée and Jang, 2015; Song, 2015; Borzée et al., 2017a). In this regard, we recommend the implementation of protected area at the sites of Sihung, Iksan, Paju, Asan, and Chungju based on genetic variability (Borzée et al., 2018c; **Figure 1**), large source populations, buffer populations and important populations for connectivity (Borzée, 2018).

Habitat Protection

We recommend the ban of pesticides and herbicides in the rice-paddy complexes where *D. suweonensis* is known to occur (Borzée et al., 2017a; **Figure 1**), as these are related to lower population sizes (Borzée et al., 2018d). The chemicals to be avoided are the ones resulting in high levels of phosphates and nitrates in the environment, and their application should be avoided during the breeding season (May and June) and tadpole development (June and July). We recommend a shift to ecological agriculture, a demonstrated way to maintain sustainability (Bellon and Penvern, 2014), including for rice harvest (Mendoza, 2008; Lestari and Suryana, 2013), as recommended by the government of the Republic of Korea (Jeong and Moon, 2013; Ministry of Agriculture Food and Rural Affairs, 2013) and already locally demonstrated (Lee et al., 2016). This would enable the conservation of the *D. suweonensis* in rice agricultural landscapes. We recommend the policy shift to be acknowledged by a special “Suweon Treefrog friendly” label, which would be economically sustainable (Jačudová et al., 2018) and in line with the rising environment and conservation awareness worldwide



FIGURE 1 | Distribution of *Dryophytes suweonensis* in the Republic of Korea. Each black patch represents an area within which sites where the species is present are connected (Borzée et al., 2018a), but disconnected from the nearest rice-paddy complex (Borzée, 2018). The species range is extracted from Borzée et al. (2017a). Map generated through ArcMap 10.5 (Environmental Systems Resource Institute, Redlands, California, USA).

(Sutherland et al., 2018). This is the case of “Gangwha Maewhamalleum rice,” set up to protect the endangered *Ranunculus kazusensis makino* in a Ramsar site composed of rice paddies only and managed by the Korea National Trust Foundation (Ramsar Convention, 2018).

Microhabitat Management

Finally, we recommend a limit in the cutting of grass surrounding rice paddies at all sites where *D. suweonensis* occurs. Mowing grass present on ditches and delimitation banks between rice paddies is negatively impacting the species, as this environment is the preferred diurnal resting habitat (Borzée et al., 2016a; Groffen et al., 2018) and is also used for feeding, microhabitat segregation with *D. japonicus* and overwintering (Borzée et al., 2016a, 2019). We acknowledge the need for weed control and shade prevention for rice growth, and when the vegetation needs to be cut, it should not be done below 30 cm as frogs diurnal activity is below this height (Borzée et al., 2016a), and as this is unlikely to impact rice production due to a lower height than the rice itself. When possible, vegetation including Korean willows (*Salix koreensis*), and occasionally high grasses such as *Elymus repens* and *Phragmites communis* should be maintained in the vicinity of rice paddies to allow for population connectivity (Driscoll et al., 2013; Schneider-Maunoury et al., 2016), as well as foraging and overwintering by *D. suweonensis* (Borzée et al., 2016a, 2019).

REFERENCES

- Baker, J., Beebe, T., Buckley, J., Gent, T., and Orchard, D. (2011). *Amphibian Habitat Management Handbook*. Bournemouth: Amphibian and Reptile Conservation.
- Beebe, T. J. C., and Griffiths, R. A. (2005). The amphibian decline crisis: a watershed for conservation biology? *Biol. Cons.* 125, 271–285. doi: 10.1016/j.biocon.2005.04.009
- Bellon, S., and Penvern, S. (2014). “Organic food and farming as a prototype for sustainable agricultures,” in *Organic Farming, Prototype for Sustainable Agricultures*, eds S. Bellon and S. Penvern (Dordrecht: Springer), 1–19.
- Bishop, P. J., Angulo, A., Lewis, J. P., Moore, R. D., Rabb, G. B., and Garcia Moreno, J. (2012). The amphibian extinction crisis—what will it take to put the action into the Amphibian Conservation Action Plan? *Surveys Perspect. Integr. Environ. Soc.* 5, 97–111.
- Borzée, A. (2018). *Why are Anurans Threatened? The case of Dryophytes suweonensis*. Ph.D. Seoul National University, Seoul, Republic of Korea.
- Borzée, A., Andersen, D., and Jang, Y. (2018a). Population trend inferred from aural surveys for calling anurans in Korea. *PeerJ*. 6:e5568. doi: 10.7717/peerj.5568
- Borzée, A., Choi, Y., Kim, Y. E., Jablonski, P. G., and Jang, Y. (2019). Interspecific variation in seasonal migration and brumation behavior in two closely related species of treefrogs. *Front. Ecol. Evol.* 7:55. doi: 10.3389/fevo.2019.00055
- Borzée, A., Heo, K., and Jang, Y. (2018b). Relationship between agroenvironmental variables and breeding Hylids in rice paddies. *Sci. Rep.* 8:8049. doi: 10.1038/s41598-018-26222-w
- Borzée, A., and Jang, Y. (2015). Description of a seminatural habitat of the endangered Suweon treefrog, *Hyla suweonensis*. *Anim Cells Syst.* 19, 1–5. doi: 10.1080/19768354.2015.1028442
- Borzée, A., and Jang, Y. (2017). Impact of rice and bean harvests on the Suweon Treefrog (*Dryophytes suweonensis*). *Int. J. Curr. Res.* 9, 59620–59623.
- Borzée, A., and Jang, Y. (2018a). Interference competition driven by hydric stress in Korean Hylids. *Nat. Conserv. Res.* 3, 120–124. doi: 10.24189/ncr.2018.008
- Borzée, A., and Jang, Y. (2018b). “Large-scale hybridisation as an extinction threat to the Suweon treefrog (*Dryophytes suweonensis*),” in *Conservation Asia 2018, vol. 1*, (Bishkek: Society for Conservation Biology – Asia section).
- Borzée, A., Kim, J. Y., Cunha, M. A. M., Lee, D., Sin, E., Oh, S., et al. (2016a). Temporal and spatial differentiation in microhabitat use: Implications for reproductive isolation and ecological niche specification. *Integr. Zool.* 11, 375–387. doi: 10.1111/1749-4877.12200
- Borzée, A., Kim, J. Y., and Jang, Y. (2016b). Asymmetric competition over calling sites in two closely related treefrog species. *Sci. Rep.* 6:32569. doi: 10.1038/srep32569
- Borzée, A., Kim, K., Heo, K., Jablonski, P. G., and Jang, Y. (2017a). Impact of land reclamation and agricultural water regime on the distribution and conservation status of the endangered *Dryophytes suweonensis*. *PeerJ*. 5:e3872. doi: 10.7717/peerj.3872
- Borzée, A., Kim, Y. I., Kim, Y. E., and Jang, Y. (2018c). Translocation of the endangered and endemic Korean treefrog *Dryophytes suweonensis*. *Conserv. Evid.* 15, 6–11.
- Borzée, A., Kosch, T. A., Kim, M., and Jang, Y. (2017b). Introduced bullfrogs are associated with increased *Batrachochytrium dendrobatidis* prevalence and reduced occurrence of Korean treefrogs. *PLoS ONE*. 12:e0177860. doi: 10.1371/journal.pone.0177860
- Borzée, A., Kyong, C. N., Kil, H. K., and Jang, Y. (2018d). Impact of water quality on the occurrence of two endangered Korean anurans: *Dryophytes suweonensis* and *Pelophylax chosonicus*. *Herpetologica*. 74, 1–7. doi: 10.1655/Herpetologica-D-17-00011
- Borzée, A., and Seliger, B. (2018). *Dryophytes suweonensis* (Suweon Treefrog). *Herpetol. Rev.* 49:707.
- Borzée, A., Yu, A.-Y., and Jang, Y. (2018e). Variations in boldness, behavioural and physiological traits of an endangered and a common hylid species from Korea. *Ethol. Ecol. Evol.* 30, 515–533. doi: 10.1080/03949370.2018.1441192
- Borzée, A., Yu, S. H., and Jang, Y. (2016c). *Dryophytes suweonensis* (Suweon Treefrog). *Herpetol. Rev.* 47:418.
- Chun, S., Chung, E., Voloshina, I., Chong, J. R., Lee, H., and Min, M.-S. (2012). Genetic diversity of Korean tree frogs (*Hyla suweonensis* and *Hyla japonica*):

The implementation of these policies will enable the survival of the Suweon treefrog (*D. suweonensis*) and the continued production of rice, highlighting the importance of the Republic of Korea in biological conservation and organic sustainable development. Moreover, our recommendations are based on a review of the literature presenting such cases and can be adapted to protect any species relying on rice paddies during their life cycle.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

This policy results from the research supported by numerous Small Grants for Science and Conservation from The Biodiversity Foundation, with special thanks to Sanha Kim and Jaeha Ahn, and a grant from the National Geographic Society Asia (Young Explorer #17-15) to AB; and Research Grants from the National Research Foundation of Korea (#2017R1A2B2003579) and the Rural Development Administration (PJ012285) to YJ. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

- assessed by mitochondrial cytochrome b gene and cytochrome oxidase subunit I gene. *Kor. J. Herpetol.* 4, 31–41. doi: 10.1080/23802359.2017.1292475
- Driscoll, D. A., Banks, S. C., Barton, P. S., Lindenmayer, D. B., and Smith, A. L. (2013). Conceptual domain of the matrix in fragmented landscapes. *Trends Ecol. Evol.* 28, 605–613. doi: 10.1016/j.tree.2013.06.010
- Duellman, W. E., Marion, A. B., and Hedges, S. B. (2016). Phylogenetics, classification, and biogeography of the treefrogs (Amphibia: Anura: Arboranae). *Zootaxa.* 4104, 1–109. doi: 10.11646/zootaxa.4104.1.1
- Dufresnes, C., Borzée, A., Horn, A., Stöck, M., Ostini, M., Sermier, R., et al. (2015). Sex-chromosome homomorphy in Palearctic tree frogs proceeds from both turnovers and X-Y recombination. *Mol. Biol. Evol.* 32, 2328–2337. doi: 10.1093/molbev/msv113
- Fujioka, M., and Lane, S. J. (1997). The impact of changing irrigation practices in rice fields on frog populations of the Kanto Plain, central Japan. *Ecol. Res.* 12, 101–108. doi: 10.1007/BF02523615
- Groffen, J., Borzée, A., and Jang, Y. (2018). Positioning of two treefrog species within rice paddies in relation to different habitat borders. *Anim. Cells Syst.* 22, 205–211. doi: 10.1080/19768354.2018.1475301
- Holzer, K. A. (2014). *Amphibian-Human Coexistence in Urban Areas*. San Diego, CA: University of California Davis.
- Howard, S. D., and Bickford, D. P. (2014). Amphibians over the edge: silent extinction risk of Data Deficient species. *Divers Distrib.* 20, 837–846. doi: 10.1111/ddi.12218
- IUCN (2018). *Dryophytes Suweonensis*. Gland: IUCN.
- Jačudová, J., Marková, I., Hroncová, E., and Vicianová, J. H. (2018). An assessment of regional sustainability through quality labels for small farmers' products: a Slovak case study. *Sustainability* 10:1273. doi: 10.3390/su10041273
- Jeong, H.-K., and Moon, D.-H. (2013). *Response Atrategy to the Abolishment of Low-Pesticide Agricultural Product Certification*. Seoul: Korea Rural Economic Institute.
- Kim, I.-H., Son, S.-H., Kang, S.-W., and Kim, J.-B. (2012). Distribution and habitat characteristics of the endangered Suweon-Tree Frog (*Hyla suweonensis*). *Kor. J. Herpetol.* 4, 15–22.
- Kim, Y. E. (2016). *Differential Antipredator Behavior Between Hyla Japonica and H. suweonensis Suggests Separate Evolution*. Ms.C. Ewha Womans University, Seoul.
- Kuramoto, M. (1984). Systematic implications of hybridization experiments with some eurasian treefrogs (genus *Hyla*). *Copeia* 3, 609–616. doi: 10.2307/1445141
- Lee, S., Nguyen, T. T., Poppenborg, P., Shin, H.-J., and Koellner, T. (2016). Conventional, partially converted and environmentally friendly farming in South Korea: Profitability and factors affecting farmers' choice. *Sustainability* 8:704. doi: 10.3390/su8080704
- Lee, S.-D., and Miller-Rushing, A. J. (2014). Degradation, urbanization, and restoration: a review of the challenges and future of conservation on the Korean Peninsula. *Biol Conserv.* 176, 262–276. doi: 10.1016/j.biocon.2014.05.010
- Lestari, Y. K., and Suryana, A. T. (2013). "Sustainability of organic rice farming in indonesia," in *The 10th Hokkaido Indonesia Student Association Scientific Meeting (HISAS 10)*. (Hokkaido).
- Machado, I. F., and Maltchik, L. (2010). Can management practices in rice fields contribute to amphibian conservation in southern Brazilian wetlands? *Aquatic Conserv.* 20, 39–46. doi: 10.1002/aqc.1070
- Mendoza, T. C. (2008). Evaluating the benefits of organic farming in rice agroecosystems in the Philippines. *J. Sustain. Agric.* 2, 93–115. doi: 10.1300/J064v24n02_09
- Ministry of Agriculture Food and Rural Affairs (2013). *The 3rd Environment-Friendly Agriculture Promotion 5-year Plan*. Sejong: MAFRA - Environment-friendly agriculture division.
- Naito, R., Sakai, M., Natuhara, Y., Morimoto, Y., and Shibata, S. (2013). Microhabitat use by *Hyla japonica* and *Pelophylax porosa brevipoda* at levees in rice paddy areas of Japan. *Zool. Sci.* 30, 386–391. doi: 10.2108/zsj.30.386
- Noh, H. K. (2017). "Endangered wildlife restoration road map (2017–2021)," in *Focus on Medium and Large Mammal Species. vol. 1*, ed S.O.R.O.E. Wildlife (Wonju: Korea National Park Services & Ministry of Environment), 11.
- Orchard, D., Tessa, G., and Jehle, R. (2019). Age and growth in a European flagship amphibian: equal performance at agricultural ponds and favourably managed aquatic sites. *Aquat. Ecol.* 53: 37–48. doi: 10.1007/s10452-018-09671-3
- Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., et al. (2014). The biodiversity of species and their rates of extinction, distribution, and protection. *Science* 344, 987–997. doi: 10.1126/science.1246752
- Ramsar Convention (2018). "Annotated list of wetlands of international importance - republic of korea. ganghwa maehwamareum habitat," in *Ganghwa Maehwamareum Habitat, vol. 1*. ed R. S. I. Service (Gland: Ramsar Convention), 3.
- Ramsar Convention Secretariat (2013). *The Ramsar Convention Manual: a Guide to the Convention on Wetlands (Ramsar, Iran, 1971), 6th edn*. Gland: Ramsar Convention Secretariat.
- Schneider-Maunoury, L., Lefebvre, V., Ewers, R. M., Medina-Rangel, G. F., Peres, C. A., Somarriba, E., et al. (2016). Abundance signals of amphibians and reptiles indicate strong edge effects in Neotropical fragmented forest landscapes. *Biol. Conserv.* 200, 207–215. doi: 10.1016/j.biocon.2016.06.011
- Snyder, N. F., Derrickson, S. R., Beissinger, S. R., Wiley, J. W., Smith, T. B., Toone, W. D., et al. (1996). Limitations of captive breeding in endangered species recovery. *Conserv. Biol.* 10, 338–348. doi: 10.1046/j.1523-1739.1996.10020338.x
- Song, W. (2015). Habitat analysis of *Hyla suweonensis* in the breeding season using species distribution modeling. *J. Kor. Soc. Environ. Restor. Technol.* 18, 71–82. doi: 10.13087/kosert.2015.18.1.71
- Stuart, S. N., Chanson, J. S., Cox, N. A., Young, B. E., Rodrigues, A. S., Fischman, D. L., et al. (2004). Status and trends of amphibian declines and extinctions worldwide. *Science* 306, 1783–1786. doi: 10.1126/science.1103538
- Sutherland, W. J., Dicks, L. V., Ockendon, N., Petrovan, S. O., and Smith, R. K. (2018). *WhatWorks in Conservation: 2018*. Cambridge, UK: Open Book Publishers.
- Wake, D. B. (2012). Facing extinction in real time. *Science* 335, 1052–1053. doi: 10.1126/science.1218364
- Wren, S., Angulo, A., Meredith, H., Kielgast, J., and Santos, M. D., Bishop, P. (2015). *Amphibian Conservation Action Plan*. IUCN SSC Amphibian Specialist Group. Available online at: <http://www.amphibians.org/acap/> (accessed March 24, 2019).

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Borzée and Jang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.