



OPEN ACCESS

EDITED AND REVIEWED BY
Paolo Perona,
EPFL, Switzerland

*CORRESPONDENCE
Devanita Ghosh,
✉ devnita.rs@gmail.com

RECEIVED 21 December 2023
ACCEPTED 22 December 2023
PUBLISHED 05 January 2024

CITATION

Ghosh D, Haq I, Isa MH, Maity JP, Bhattacharya P and Ahmad A (2024), Editorial: Assessment of environmental risk and challenges in addressing emerging pollutants: current and future perspectives. *Front. Environ. Sci.* 11:1359376. doi: 10.3389/fenvs.2023.1359376

COPYRIGHT

© 2024 Ghosh, Haq, Isa, Maity, Bhattacharya and Ahmad. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Editorial: Assessment of environmental risk and challenges in addressing emerging pollutants: current and future perspectives

Devanita Ghosh^{1*}, Izharul Haq², Mohamed Hasnain Isa³, Jyoti Prakash Maity⁴, Prosun Bhattacharya⁵ and Akil Ahmad⁶

¹Sanitary Engineering Section, Water Management Department, Delft University of Technology, Delft, Netherlands, ²Department of Civil Engineering, Indian Institute of Technology Guwahati, Guwahati, India, ³Faculty of Engineering, University of Technology Brunei Gadong, Gadong, Brunei, ⁴Department of Chemistry, School of Applied Sciences, KIIT Deemed to Be University, Bhubaneswar, Odisha, India, ⁵Department of Sustainable Development, Environmental Science and Engineering, Royal Institute of Technology, Stockholm, Sweden, ⁶College of Sciences and Humanities, Prince Sattam Bin Abdulaziz University, Al-Kharj, Saudi Arabia

KEYWORDS

micropollutants, wastewater, treatment technologies, toxic effects, sustainability, toxicology, biological treatment, industrial wastewater

Editorial on the Research Topic

[Assessment of environmental risk and challenges in addressing emerging pollutants: current and future perspectives](#)

Industrial wastewater is a pressing issue due to significant water and chemical usage in production, leading to the discharge of harmful micropollutants, toxins, and lethal substances into the environment without proper treatment. Employing eco-friendly biological treatment systems involving microorganisms like bacteria, fungi, and enzymes is crucial for detoxifying such contaminated water. Hence, urgent measures are needed to develop sustainable methods for treating emerging pollutants in wastewater to safeguard the environment. The toxicity of emerging pollutants poses a global environmental concern, including rare Earth elements, micro/nano plastics, antibiotics, and pharmaceuticals. Inadequate treatment systems discharge many of these hazardous compounds, challenging their detection due to their adverse effects even at low concentrations. Nano/micro-scale physical pollutants add complexity, while the fate, transport, and toxicity of these contaminants remain unclear. Understanding their behaviours and advancing remediation techniques is critical to mitigating the ecosystem exposure risks.

Emerging pollutants, largely unregulated, are found in drainage systems, medication residues, household products, and agricultural runoff. Predominantly organic compounds, their fate, toxicity, and mitigation strategies remain ambiguous. Research aimed to uncover the occurrence, sources, and treatment of emerging pollutants in industrial wastewater through conventional and advanced technologies. Five state-of-the-art publications were generated under this flagship.

Gao et al., reported the presence of pharmaceuticals in the aqueous environment has had adverse effects on both human health and the balance of ecosystems. Removing

pharmaceutical contaminants using conventional treatment technologies has proven challenging. To effectively degrade carbamazepine (CBZ), a common antiepileptic drug, three plasma discharge modes were investigated: liquid-phase discharge, air gas-liquid jet, and dielectric barrier discharge. Their results showed the highest levels of degradation efficiency and energy yield (0.11 g/kWh), achieved while operating at 225 W input power under liquid-phase discharge, surpassing the other discharge modes. In this research article, parameters such as concentration, electrode distance, volume, input power, and discharge frequency were systematically evaluated and optimized. Plasma emission spectroscopy revealed that the degradation process is primarily attributed to reactive oxygen species (ROS), with the role of reactive nitrogen species being insignificant. The potential degradation intermediates and pathways of CBZ were investigated and analysed using liquid chromatography-mass spectrometry (LC-MS). Considering aspects such as cleanliness, degradation efficiency, and application feasibility, the liquid-phase discharge treatment emerges as a promising technology for water treatment. Their comprehensive approach highlights the potential of this method for effectively addressing pharmaceutical contamination in aquatic environments.

Wang et al., further reported the contamination of per- and polyfluoroalkyl substances (PFAS) in the rivers of Shandong, China. They investigated the spatiotemporal variations, mass discharges, and ecological risks of PFAS by collecting surface waters seasonally at the estuaries of 13 major rivers in Shandong. Among the compounds that were studied, two novel perfluoroalkyl ether carboxylic acids, hexafluoropropylene oxide-dimer acid (HFPO-DA) and hexafluoropropylene oxide-trimer acid (HFPO-TA), along with 10 legacy PFAS. All were detectable, and their total concentrations (\sum PFAS) exhibited a wide range from 23 to 25,400 ng/L. Elevated levels of all target compounds were notably observed in the Xiaoqing River, although the Majia and Jiaolai Rivers also raised concerns. Perfluorooctanoic acid (PFOA) stood out as dominant across all rivers, with its concentration in the Xiaoqing River estuary exhibiting a six-fold increase from 2011 to 2014 and stabilizing during 2014–2020. The absence of significant seasonality in PFAS concentrations in rivers with seasonal flows in their study suggests different emission patterns, possibly higher emissions during the wet season and lower emissions during the dry season. A proposed parameter, the ratio of maximum to minimum concentration (C_{\max}/C_{\min}), aimed to characterize the influence of local point sources, with a wide range observed from 10 to 14,000. The Xiaoqing and Yellow Rivers were identified as discharging approximately 98% of the \sum PFAS flux into the sea, estimated at 28.0 tons in 2020. Their study highlighted the need to focus on industrial emission patterns of PFAS and emphasizes the necessity for further investigations with higher time resolution to comprehend the emission dynamics of PFAS in Shandong.

Li et al., highlighted the viability of immobilized microbial technology in combatting water pollution. They stressed the importance of understanding microorganisms in immobilized biochar for effective contaminant removal. Several challenges were identified, including microporous structure blockage hindering microbe-pollutant contact, unstable microbial

loading, and extended cycle times. Their study focused on *A. faecalis* immobilization on rice hull biochar for phenol degradation. *A. faecalis* JH1 effectively removed 300 mg/L of phenol in 24 h. Phenol adsorption by rice husk biochar increased with higher pyrolysis temperatures (700°C > 500°C > 300°C). Using 700°C biochar, JH1 achieved 249.45 nmol P/g of immobilized biomass. Remarkably, JH1 consistently eliminated 300 mg/L of phenol within 12 h after six cycles. Bacterial growth formed a robust biofilm, accelerating phenol removal. Their study emphasized the strong adherence of *A. faecalis* to rice hull biochar and its efficient phenol degradation. Overall, their findings demonstrate *A. faecalis* potential for effective, durable phenol removal when immobilized on rice hull biochar.

Rana et al., used nuclear and conventional methods to analyse metal pollutants in Islamabad-Rawalpindi's groundwater (the "twin cities"). Of the 122 samples they have studied, most met aesthetic and WHO/PSQCA standards for parameters like pH, copper, and zinc. However, elevated electrical conductivity and dissolved solids made 9.83% and 4.09% of samples unsuitable for drinking. Lead and iron exceeded safe limits. Anthropogenic and geogenic factors, like waste disposal and over abstraction, contributed to metal presence. The water quality index revealed poor groundwater quality due to high metal levels. Isotopic data highlighted vulnerability to various sources. This underscores the impact of urbanization, waste mismanagement, and lack of water policies on groundwater quality. Regulating abstraction and improving treatment systems are urgent. Their findings aid in strategizing to improve water quality in the "twin cities."

Ibrahim presented a review elaborating on how the COVID-19 pandemic has significantly affected public health, economies, social life, and the environment worldwide. While much research has focused on medications like antivirals and vaccines, studies on their environmental impact, especially in the water sector, are limited. The review provided a comprehensive summary of COVID-19's effects on water, covering detection methods, virus presence in wastewater, transmission probabilities, variant detection, and both positive and negative impacts on the water sector. It offers recommendations for governments, policymakers, water treatment plants, the public, and researchers to minimize its effects and prepare for future pandemics. Their aim is to highlight the pandemic's various aspects and suggest measures to reduce its impact and that of potential future pandemics.

Overall, this Research Topic provided valuable insights into environmental challenges stemming from industrial wastewater, emphasizing the urgency of sustainable treatment methods for emerging pollutants. The focus on the COVID-19 pandemic highlighted its multifaceted impact on public health, the economy, and the environment, particularly in the water sector. Additionally, it delved into research concerning microbial technology in water pollution mitigation and the need for a holistic approach to understand and address environmental concerns. Overall, these Research Topic underscore the significance of proactive measures, interdisciplinary research, and innovative solutions to tackle contemporary global challenges.

Author contributions

DG: Writing–original draft, Writing–review and editing. IH: Writing–original draft, Writing–review and editing. MI: Writing–original draft, Writing–review and editing. JM: Writing–original draft, Writing–review and editing. PB: Writing–original draft, Writing–review and editing. AA: Writing–original draft, Writing–review and editing.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.