



# An Upstream Open Reading Frame Represses Translation of Chicken PPAR $\gamma$ Transcript Variant 1

Yankai Chu<sup>1,2,3</sup>, Jiaxin Huang<sup>1,2,3</sup>, Guangwei Ma<sup>1,2,3</sup>, Tingting Cui<sup>1,2,3</sup>, Xiaohong Yan<sup>1,2,3</sup>, Hui Li<sup>1,2,3</sup> and Ning Wang<sup>1,2,3\*</sup>

<sup>1</sup> Key Laboratory of Chicken Genetics and Breeding, Ministry of Agriculture and Rural Affairs, Harbin, China, <sup>2</sup> Key Laboratory of Animal Genetics, Breeding and Reproduction, Education Department of Heilongjiang Province, Harbin, China, <sup>3</sup> College of Animal Science and Technology, Northeast Agricultural University, Harbin, China

## OPEN ACCESS

### Edited by:

Graziano Pesole,  
University of Bari Aldo Moro, Italy

### Reviewed by:

Petar Ozretić,  
Rudjer Boskovic Institute, Croatia  
Chi-Ming Wong,  
The University of Hong Kong,  
Hong Kong

### \*Correspondence:

Ning Wang  
wangning@neau.edu.cn

### Specialty section:

This article was submitted to  
RNA,  
a section of the journal  
Frontiers in Genetics

Received: 05 December 2019

Accepted: 12 February 2020

Published: 28 February 2020

### Citation:

Chu Y, Huang J, Ma G, Cui T,  
Yan X, Li H and Wang N (2020) An  
Upstream Open Reading Frame  
Represses Translation of Chicken  
PPAR $\gamma$  Transcript Variant 1.  
Front. Genet. 11:165.  
doi: 10.3389/fgene.2020.00165

Peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) is a master regulator of adipogenesis. The PPAR $\gamma$  gene produces various transcripts with different 5'-untranslated regions (5' UTRs) because of alternative promoter usage and splicing. The 5' UTR plays important roles in posttranscriptional gene regulation. However, to date, the regulatory role and underlying mechanism of 5' UTRs in the posttranscriptional regulation of PPAR $\gamma$  expression remain largely unclear. In this study, we investigated the effects of 5' UTRs on posttranscriptional regulation using reporter assays. Our results showed that the five PPAR $\gamma$  5' UTRs exerted different effects on reporter gene activity. Bioinformatics analysis showed that chicken PPAR $\gamma$  transcript 1 (PPAR $\gamma$ 1) possessed an upstream open reading frame (uORF) in its 5' UTR. Mutation analysis showed that a mutation in the uORF led to increased *Renilla* luciferase activity and PPAR $\gamma$  protein expression, but decreased *Renilla* luciferase and PPAR $\gamma$ 1 mRNA expression. mRNA stability analysis using real-time RT-PCR showed that the uORF mutation did not interfere with mRNA stability, but promoter activity analysis of the cloned 5' UTR showed that the uORF mutation reduced promoter activity. Furthermore, *in vitro* transcription/translation assays demonstrated that the uORF mutation markedly increased the translation of PPAR $\gamma$ 1 mRNA. Collectively, our results indicate that the uORF represses the translation of chicken PPAR $\gamma$ 1 mRNA.

**Keywords:** PPAR $\gamma$ , 5'-untranslated region, upstream open reading frame, translational repression, gene expression

## INTRODUCTION

Peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) is a member of the PPAR subfamily of ligand-activated transcription factors. *In vitro* and *in vivo* studies have demonstrated that PPAR $\gamma$  is essential for adipocyte differentiation, adipocyte survival, adipocyte function, insulin sensitivity, and lipogenesis (Lehrke and Lazar, 2005; Lefterova et al., 2014). Synthetic PPAR $\gamma$  agonists have been used as therapeutic agents for diabetes and insulin insensitivity (Cariou et al., 2012).

The PPAR $\gamma$  gene is controlled by multiple alternative promoters (Aprile et al., 2014; Chandra et al., 2017). Because of alternative promoter usage and splicing, the PPAR $\gamma$  gene can produce multiple transcript variants, resulting in expression of two PPAR $\gamma$  protein isoforms that differ

in the N-terminal. All PPAR $\gamma$  transcript variants differ in their 5'-untranslated regions (5' UTRs) (McClelland et al., 2014). These PPAR $\gamma$  5' UTR isoforms have distinct tissue distributions (Ahmadian et al., 2013), suggesting that the 5' UTRs may be involved in posttranscriptional and translational regulation of the PPAR $\gamma$  gene.

The 5' UTRs of mRNAs exert crucial roles in posttranscriptional and translational regulation. Several *cis*-regulatory elements within the 5' UTRs have been identified, such as the 5' cap structure (Mitchell et al., 2010), upstream open reading frames (uORFs) (Hood et al., 2009; Barbosa et al., 2013), internal ribosome entry sites (IRES) (Xia and Holcik, 2009), terminal oligo-pyrimidine tracts, secondary structures, and G-quadruplexes (Yamashita et al., 2008; Bugaut and Balasubramanian, 2012). These *cis*-regulatory elements can function via various mechanisms, controlling mRNA stability (Nasif et al., 2018), nuclear export, localization, and translation efficiency (Araujo et al., 2012). Of these *cis*-regulatory elements, uORFs have been widely studied. Bioinformatics analysis showed that about 50% of human transcripts contain uORFs (Suzuki et al., 2000; Iacono et al., 2005; Calvo et al., 2009), and experimental studies have revealed that a number of uORFs can affect the expression of the main downstream ORFs by inducing mRNA decay or by regulating translation (Iacono et al., 2005; Crowe et al., 2006; Sathirapongasuti et al., 2011).

Given the importance of PPAR $\gamma$  in various physiological and pathological processes, PPAR $\gamma$  gene regulation has been extensively studied at the genomic and transcriptional levels in recent decades (Lee and Ge, 2014). The half-life of PPAR $\gamma$  mRNA and protein is short and PPAR $\gamma$  protein can be post-translationally modified in various ways (van Beekum et al., 2009; Katsura et al., 2014), suggesting that posttranscriptional regulation is crucial for its function. However, to date, posttranscriptional regulation by the 5' UTR has been mostly unexplored. In the present study, we investigated the posttranscriptional regulation of chicken PPAR $\gamma$  by the 5' UTR. Of interest, we demonstrated that translation of chicken PPAR $\gamma$  transcript variant 1 (PPAR $\gamma$ 1) is repressed by a uORF that is absent in human and mouse PPAR $\gamma$  transcripts.

## MATERIALS AND METHODS

### Cell Culture

Chicken embryo fibroblast (DF1) cell line was purchased from the Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences, and the immortalized chicken preadipocyte cell line 1 (ICP1) was generated in our laboratory (Wang et al., 2017). All cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin, at 37°C and 5% CO<sub>2</sub>. The culture medium was changed two to three times per week and cells were passaged 1:3 or 1:5 as needed.

### Plasmid Construction

For PPAR $\gamma$  5' UTR reporter constructs, the reporter vector psi-CHECK2 (Invitrogen, Carlsbad, CA, United States) was

mutated using the Site-directed Gene Mutagenesis Kit (Beyotime Institute of Biotechnology, Shanghai, China) according to the manufacturer's instructions, in which the initiation start codon ATG of *Renilla* luciferase was mutated to TTG, and the resultant vector was named psi-CHECK2-Mut. Then, the DNA sequences corresponding to the five different PPAR $\gamma$  5' UTRs plus initiation codon ATG were synthesized and inserted into the *Nhe*I restriction site upstream of the *Renilla* luciferase gene to create the five chicken PPAR $\gamma$  5' UTR reporter constructs: PPAR $\gamma$ 1-5'UTR, PPAR $\gamma$ 2-5'UTR, PPAR $\gamma$ 3-5'UTR, PPAR $\gamma$ 4-5'UTR, and PPAR $\gamma$ 5-5'UTR, respectively. In these five 5' UTR reporter constructs, the *Renilla* luciferase expression was driven by SV40 early enhancer/promoter, and these PPAR $\gamma$  5' UTRs were expressed as the respective 5' UTRs of *Renilla* luciferase mRNAs.

To test the promoter activity of the DNA sequences corresponding to the PPAR $\gamma$ 1 wild-type and uORF-mutant 5' UTRs, which the uAUG was mutated to a stop codon UAG (AUG > UAG), the DNA sequences were synthesized and subcloned into the *Bam*HI and *Xho*I restriction sites of pGL3-basic vector and named pGL3-PPAR $\gamma$ -WT and pGL3-PPAR $\gamma$ -Mut, respectively. Site-directed mutagenesis of the uORF was performed by DNA synthesis (GENEWIZ, Suzhou, China).

For PPAR $\gamma$  expression constructs, the full-length coding sequence of PPAR $\gamma$ 1 was PCR amplified from the cDNA derived from DF1 cells with a set of primers (forward primer: 5'-GAATTCATGGTTGACACAGAAATGCCGT-3' and reverse primer: 5'-CCTCGAGGAGGATAAGAACTACTATCGCC-3') and cloned into the *Bam*HI and *Eco*RI restriction sites of the pcDNA3.1 expression vector. The synthesized DNA fragments corresponding to the wild-type and uORF mutated 5' UTR (AUG > UAG) of PPAR $\gamma$ 1 were inserted with *Bam*HI and *Nhe*I restriction sites upstream of PPAR $\gamma$  ORF in pcDNA3.1 vector. The resultant vectors were named pcDNA3.1-PPAR $\gamma$ -WT and pcDNA3.1-PPAR $\gamma$ -Mut, respectively. All constructs were confirmed by DNA sequencing and restriction enzyme digestion.

### Quantitative Real-Time PCR Assays (qRT-PCRs)

Total RNA was isolated from ICP1 or DF1 cells using the RNeasy Plus Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol, and the first-strand cDNA was synthesized from 1  $\mu$ g of total RNA with oligo dT or random primers using ImProm-II reverse transcriptase (Promega, Madison, WI, United States). The qPCR reactions were performed in a 20  $\mu$ L reaction mixture using FastStart Universal SYBR Green Master [Rox] (Roche, Madison, WI, United States). The primers were as follows: *Renilla* luciferase (*Rluc*) (forward 5'-TGATCGAGTCCTGGGACGA-3', reverse 5'-ACAATCTGGACGACGTCGGG-3'); wild-type and uORF-mutated PPAR $\gamma$ 1 (forward 5'-GGAGTTTATCCCACCAGAAG-3', reverse 5'-AATCAACAGTGGTAAATGGC-3'); *NONO* (forward 5'-AGAAGCAGCAGCAAGAAC-3', reverse 5'-TCCTCCATCCTCCTCAGT-3'). qPCR was carried out in an ABI 7500 real-time PCR system (Applied Biosystems, Foster City, CA, United States), and PCR results were recorded as threshold cycle numbers (Ct). The fold change in the target gene

expression, normalized to the expression of an internal control gene (*NONO*) and relative to the expression at time point 0 (Normann et al., 2016), was calculated using the  $2^{-\Delta\Delta Ct}$  method (Livak and Schmittgen, 2001). The results are presented as the mean  $\pm$  SEM of three independent experiments.

## Protein Isolation and Western Blot Analysis

The ICP1 cells were transfected with either pcDNA3.1-PPAR $\gamma$ -WT or pcDNA-PPAR $\gamma$ -Mut vector. At 48 h post-transfection, cells were washed twice with PBS and lysed using radioimmunoprecipitation assay (RIPA) buffer (Beyotime Institute of Biotechnology, Shanghai, China) supplemented with 1% protease inhibitor mixture. Equal amounts of protein extracts were separated by sodium dodecyl sulfate-PAGE, transferred onto Immobilon-P PVDF membranes (Millipore, Billerica, MA, United States). The membrane was blocked for 1 to 2 h at room temperature with Tris-buffered saline containing 0.1% Tween and 5% non-fat dry milk, and immunoblotted with rabbit polyclonal antibody to chicken PPAR $\gamma$  (1:1000 dilution) or  $\beta$ -actin (1:1000 dilution, ZSGB-BIO, Beijing, China) at room temperature for 1 h. Horseradish peroxidase-conjugated goat anti-mouse or goat anti-rabbit IgG (Promega, Madison, WI, United States; 1:10,000) was incubated for 1 h at room temperature and then washed four times with PBS-Tween for 20 min. The immunoreactive bands were visualized using an ECL Plus detection kit (HaiGene Biotechnology, Harbin, China). Immunoreactive protein levels were determined semi-quantitatively by densitometric analysis using the UVP system Labworks TM software 3.0 (UVP, Upland, CA, United States). Each western blot analysis was performed at least three times.

## Dual-Luciferase Reporter Assays

Transient transfections were performed using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, United States). Cells were plated at  $1.0$  to  $1.5 \times 10^5$  cells per well in 24-well plates. For the 5' UTR reporter gene assay, 1  $\mu$ g of the indicated reporter constructs was transfected into each well. For the promoter reporter gene assay, 0.8  $\mu$ g of the indicated reporter constructs and 0.4  $\mu$ g of pRL-TK (Invitrogen, Carlsbad, CA, United States), as an internal control of transfection efficiency, were co-transfected into each well. Luciferase activity was analyzed at 48 h post-transfection, using a dual-luciferase reporter kit (Promega, Madison, WI, United States) as per the manufacturer's instructions. All luciferase reporter assays were performed at least three times in quadruplicates.

## In vitro Transcription and Translation

Plasmids pcDNA3.1-PPAR $\gamma$ -WT and pcDNA3.1-PPAR $\gamma$ -Mut were linearized, purified by agarose gel electrophoresis, eluted with diethylpyrocarbonate-treated H<sub>2</sub>O, and quantified. Equal amounts (1  $\mu$ g) of linearized DNA were used as templates for *in vitro* transcription in the T7 RiboMAX Large Scale RNA Production System (Promega, Madison, WI, United States) according to the manufacturer's protocol. Capped mRNAs were generated using the Ribo m<sup>7</sup>G Cap Analog (Promega, Madison,

WI, United States). The capped mRNAs were digested with DNase I and purified with the RNeasy kit (Qiagen, Hilden, Germany) and quantified. The size and integrity of the purified mRNAs were assessed by gel electrophoresis. The mRNA outputs of pcDNA3.1-PPAR $\gamma$ -WT and pcDNA3.1-PPAR $\gamma$ -Mut were analyzed by absolute qRT-PCR. *In vitro* translation reactions were performed in nuclease-treated Rabbit Reticulocyte Lysate (Promega, Madison, WI, United States) as described by the manufacturer. Equal amounts of the capped mRNA (2  $\mu$ g) derived from pcDNA3.1-PPAR $\gamma$ -WT or pcDNA3.1-PPAR $\gamma$ -Mut construct were used as the template for *in vitro* translation, which was performed for 60 min at 30°C, and the reactions were stopped by transferring the tubes to ice. Biotinylated lysine residues were added to the translation reaction as a precharged  $\epsilon$ -labeled biotinylated lysine-tRNA complex (Transcend tRNA; Promega, Madison, WI, United States) and incorporated into nascent proteins during translation. The translated protein was analyzed using a Transcend Non-Radioactive Translation Detection System (Promega, Madison, WI, United States).

## RNA Stability Assay

The stability of luciferase mRNA transcripts from the indicated constructs (PPAR $\gamma$ 1-5'UTR-WT and PPAR $\gamma$ 1-5'UTR-Mut) was determined by measuring the amount of *Rluc* luciferase mRNA at selected intervals: 0 (control), 3, 6, 9, and 12 h, following the addition of 5 mg/mL actinomycin D (Sigma-Aldrich, St. Louis, MO, United States) at 48 h post-transfection. Time-course intervals were chosen based on the manufacturer's data of *luc2* mRNA half-life (approximately 2 h). For mRNA expression analysis, total RNA (1  $\mu$ g) was reverse-transcribed into cDNA using the PrimeScript RT reagent Kit with gDNA Eraser (Takara, Shiga, Japan), and relative mRNA expression was determined by real-time PCR using FastStart Universal SYBR Green Master [Rox] (Roche, Madison, WI, United States) with *Rluc* primers as described above. Relative mRNA levels were normalized to the *NONO* gene and to expression at time point 0 (Normann et al., 2016) and calculated using the  $2^{-\Delta\Delta Ct}$  method (Livak and Schmittgen, 2001).

## Bioinformatics Analysis

Online software programs used to predict the potential *cis*-regulatory elements of PPAR $\gamma$  5' UTR: StarORF<sup>1</sup> (Ceraj et al., 2009), UTRscan<sup>2</sup> (Grillo et al., 2010), and Reg RNA2.0<sup>3</sup> (Chang et al., 2013). Preliminary RNA secondary structures were predicted using Vienna RNAfold 2.0<sup>4</sup> (Hofacker, 2003). Intrinsic protein disorder analyses were made using PSIPRED protein sequence analysis workbench<sup>5</sup> (Buchan et al., 2013). All bioinformatic computations were performed using default prediction parameters.

<sup>1</sup><http://star.mit.edu/index.html>

<sup>2</sup><http://itbtools.ba.itb.cnr.it/utrscan>

<sup>3</sup><http://regrna2.mbc.nctu.edu.tw>

<sup>4</sup><http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi>

<sup>5</sup><http://bioinf.cs.ucl.ac.uk/psipred/>

## Statistical Analysis

Experimental data were analyzed using GraphPad Prism software (GraphPad Inc., San Diego, CA, United States). The results were presented as mean  $\pm$  SEM. For comparison of two groups, statistical analysis was performed using two-tailed Student's *t*-test and linear regression. *P* values < 0.05 (\*) were considered significant, *P* values < 0.01 (\*\*) were considered highly significant. For multiple comparisons, one-way analysis of variance (ANOVA) was used to determine significance, followed by Tukey's *post hoc* test.

## Data Availability Statement

Strains, plasmids and cell lines are available upon request. The authors affirm that all data necessary for confirming the conclusions of the article are present within the article, figures, and Supplementary Material.

## RESULTS

### The Effects of PPAR $\gamma$ 5' UTRs on Reporter Gene Expression


We previously identified five different chicken PPAR $\gamma$  transcript variants (PPAR $\gamma$  1 to 5) by 5' rapid amplification of cDNA ends (5' RACE) in chicken abdominal adipose tissue (Duan et al., 2015). These chicken PPAR $\gamma$  transcript variants encode two protein isoforms (PPAR $\gamma$ 1 and PPAR $\gamma$ 2) that differ in their N-terminal extension. Chicken PPAR $\gamma$ 2 contains six additional amino acids at the N-terminus compared with PPAR $\gamma$ 1. These five chicken PPAR $\gamma$  transcript variants differed in 5' UTR sequence and length, and had different tissue distribution patterns (Duan et al., 2015), suggesting that 5' UTRs may play a role in the posttranscriptional regulation of PPAR $\gamma$  gene expression. To investigate the posttranscriptional regulatory

roles of the five 5' UTR isoforms on chicken PPAR $\gamma$  gene expression, we constructed their respective 5' UTR reporter constructs. Firstly, the ATG start codon of *Renilla* luciferase gene in the psi-CHECK2 vector was mutated to TTG by site-directed mutagenesis, and the resultant vector was named psi-CHECK2-Mut. Then, the five DNA fragments corresponding to the five different PPAR $\gamma$  5' UTRs plus the start codon ATG were synthesized and inserted at the *Nhe*I restriction site upstream of the *Renilla* luciferase gene in psi-CHECK2-Mut to yield the five chicken PPAR $\gamma$  5' UTR reporter constructs.

We transfected these five 5' UTR reporters into an ICP1 cell line and a chicken embryo fibroblast (DF1) cell line and measured *Renilla* luciferase activity. The reporter gene assay showed that these five 5' UTR reporters displayed different luciferase activities. As shown in **Figures 1A,B**, PPAR $\gamma$ 1-5'UTR exhibited the highest luciferase activity in ICP1 cells and the second-highest activity in DF1 cells. PPAR $\gamma$ 5-5'UTR exhibited the highest activity in DF1 cells and the second-highest activity in ICP1 cells. PPAR $\gamma$ 3-5'UTR exhibited the lowest activity in both ICP1 and DF1 cells. PPAR $\gamma$ 2-5'UTR and PPAR $\gamma$ 4-5'UTR exhibited similar reporter activity in both ICP1 and DF1 cells. These results support our speculation that the 5' UTR regulates chicken PPAR $\gamma$  gene expression.

### Bioinformatics Analysis of Chicken PPAR $\gamma$ 5' UTRs

To gain insight into the molecular mechanisms by which the 5' UTRs regulate gene expression, we performed a bioinformatics analysis of these five 5' UTR sequences using the online software programs StarORE, UTRscan, and RegRNA 2.0. Bioinformatics analysis showed that PPAR $\gamma$ 1 5' UTR contains a 54-nucleotide (nt)-long uORF (PPAR $\gamma$ 1 uORF), PPAR $\gamma$ 3 5' UTR has a 12-nt-long uORF (PPAR $\gamma$ 3 uORF) and a putative IRES element, and PPAR $\gamma$ 5 5' UTR has two uORFs, which are 15 and 51 nt long,



**FIGURE 1 |** Effects of PPAR $\gamma$  5' UTR isoforms on reporter gene activity. **(A)** The luciferase activity of each of the PPAR $\gamma$  5' UTR reporter constructs was measured in DF1 cells. **(B)** The luciferase activity of each of the PPAR $\gamma$  5' UTR reporter constructs was measured in ICP1 cells. Data are expressed as mean  $\pm$  SEM (*n*  $\geq$  3 independent experiments). Lower case letters above bars indicate the order of expression levels, "a" represents the highest expression level, "e" represents the lowest expression level. Bars with different superscripts are mutually statistically different. ANOVA followed by Tukey's multiple comparison test was used to determine significance.


respectively. No putative *cis*-regulatory elements were predicted in PPAR $\gamma$ 2 and PPAR $\gamma$ 4 5' UTR sequences. The PPAR $\gamma$ 1 uORF is located in its 5' UTR from nucleotides -78 to -25 (relative to the start codon AUG of the PPAR $\gamma$  protein-coding ORF, where A is +1; **Figure 2A**), and the uORF AUG (uAUG) resides in a favorable Kozak consensus context, suggesting that there is a high probability that scanning ribosomes consistently initiate the translation at this uAUG codon to encode a 17-amino acid peptide (MGRPGEFIPPEGNSFSG; **Figure 2A**).


### The Effect of PPAR $\gamma$ 1 uORF on Gene Expression

Upstream ORFs have emerged as a major posttranscriptional regulatory element in eukaryotic species (Wen et al., 2009). The above bioinformatics analysis showed that, of these five PPAR $\gamma$  5'UTR isoforms, three contained uORFs, which led

us to speculate that these uORFs may be implicated in posttranscriptional regulation of chicken PPAR $\gamma$ . Herein, we focused our attention on the PPAR $\gamma$ 1 uORF. Of these five chicken PPAR $\gamma$  transcript variants, PPAR $\gamma$  transcript variant 1 (PPAR $\gamma$ 1) is highly expressed in various chicken tissues, including abdominal adipose, spleen, and liver (Duan et al., 2015), which is consistent with our results showing that PPAR $\gamma$ 1 5' UTR had high reporter activity (**Figures 1A,B**). Unlike the other two uORF-containing 5' UTR isoforms, PPAR $\gamma$ 1 5' UTR presented the largest uORF, and its uAUG was in a favorable Kozak consensus context.

To test our speculation, we investigated the effect of PPAR $\gamma$ 1 uORF on posttranscriptional regulation of the *Renilla* luciferase reporter gene. We generated a uORF-mutated reporter construct, named PPAR $\gamma$ 1-uORF-Mut, by mutating the uAUG of the PPAR $\gamma$ 1 uORF to a stop codon UAG (AUG > UAG) by site-directed mutagenesis. Transient transfection and reporter gene






assays showed that the luciferase activities of the mutant reporter construct (PPAR $\gamma$ 1-uORF-Mut) were 3- and 2.5-fold higher, respectively, than those of the wild-type PPAR $\gamma$ 1 5' UTR reporter construct (PPAR $\gamma$ 1-5'UTR-WT) in ICP1 and DF1 cells ( $P$  < 0.01, **Figure 2B**). These results indicate that this uORF functions as an intrinsic repressor for downstream ORF expression. To further understand the molecular mechanism underlying the repressive effect of this uORF, we quantified the relative mRNA levels of *Renilla luciferase (Rluc)* in cells transfected with the same amount of PPAR $\gamma$ 1-uORF-Mut and PPAR $\gamma$ 1-5'UTR-WT, respectively. Surprisingly, in contrast to the reporter gene assay results (**Figure 2B**), quantitative real-time RT-PCR showed that transfection of PPAR $\gamma$ 1-uORF-Mut resulted in lower *Rluc* mRNA expression compared with the PPAR $\gamma$ 1-5'UTR-WT in both ICP1 ( $P$  < 0.05) and DF1 cells ( $P$  < 0.01) (**Figure 2C**). Thus, the luciferase reporter gene assay and quantitative RT-PCR results together allow us to conclude that PPAR $\gamma$ 1 uORF inhibits *Rluc* translation.

### Inhibition of PPAR $\gamma$ 1 Translation by the uORF

To exclude the possibility that the effect of this uORF is reporter gene-specific, we generated full-length PPAR $\gamma$ 1 expression constructs with either the wild-type or uORF-mutated 5' UTR (AUG > UAG), termed pcDNA3.1-PPAR $\gamma$ -WT and pcDNA3.1-PPAR $\gamma$ -Mut, respectively. Then, ICP1 cells were transfected with pcDNA3.1-PPAR $\gamma$ -WT or pcDNA3.1-PPAR $\gamma$ -Mut alone, and PPAR $\gamma$  protein expression was assayed by western blot. The western blot analysis showed that PPAR $\gamma$ 1 protein levels were significantly higher in the ICP1

cells transfected with pcDNA3.1-PPAR $\gamma$ -Mut than with the pcDNA3.1-PPAR $\gamma$ -WT ( $P$  < 0.01, **Figures 3A,B**). In parallel, we investigated the PPAR $\gamma$ 1 mRNA expression. Real-time



RT-PCR analysis showed that *PPAR $\gamma$ 1* mRNA expression levels were significantly lower in both ICP1 and DF1 cells transfected with pcDNA3.1-PPAR $\gamma$ -Mut than with pcDNA3.1-PPAR $\gamma$ -WT (Figure 3C). These results are consistent with those of the reporter gene assay (Figure 2C). Collectively, these results indicate that this uORF represses downstream PPAR $\gamma$ 1 translation.

### No Effect of PPAR $\gamma$ 1 uORF on mRNA Stability


Our results showed that the uORF mutation resulted in reduced mRNA expression levels of *Rluc* and *PPAR $\gamma$ 1* (Figures 2C, 3C). There are two possibilities to explain this. First, the uORF mutation may affect mRNA stability. Previous studies have indicated that uORF can reduce mRNA expression via mRNA destabilization (Dikstein, 2012; Dvir et al., 2013). The other possibility is that the cloned chicken PPAR $\gamma$ 1 5' UTR in our 5' UTR reporters and PPAR $\gamma$  expression vectors may contain promoter activity, and uORF mutation may lead to reduced promoter activity. To test whether this uORF mutation affected mRNA stability, using real-time RT-PCR, we determined the mRNA decay rate of *Rluc* in cells transfected with PPAR $\gamma$ 1-5'UTR-WT or PPAR $\gamma$ 1-5'UTR-Mut at 0, 3, 6, 9, and 12 h following treatment with actinomycin D. As shown in Figure 4, no significant difference in *Rluc* mRNA half-life was observed over a 12-h period between cells transfected with PPAR $\gamma$ 1-5'UTR-WT or PPAR $\gamma$ 1-5'UTR-Mut (PPAR $\gamma$ 1-5'UTR-WT: 6.43 h; PPAR $\gamma$ 1-5'UTR-Mut: 5.90 h ( $P = 0.4146$ ). These results indicate that this uORF had no obvious effect on mRNA stability.

### The Effect of uORF Mutation on Promoter Activity

The genomic region corresponding to the 5' UTR is usually part of the promoter. Our previous study demonstrated that the 108-bp sequence downstream of the transcription start site of *PPAR $\gamma$ 1*, which is part of the 5' UTR, had the highest promoter activity (Cui et al., 2018). To test whether the cloned PPAR $\gamma$ 1 5' UTR had promoter activity and whether the uORF mutation reduced it, we cloned DNA sequences corresponding to wild-type and uORF-mutated 5' UTRs of PPAR $\gamma$ 1 into luciferase reporter vector pGL3-Basic, named pGL3-PPAR $\gamma$ 1-WT and pGL3-PPAR $\gamma$ 1-Mut, respectively. A reporter gene assay showed that the pGL3-PPAR $\gamma$ 1-WT and pGL3-PPAR $\gamma$ 1-Mut displayed 111- and 90-fold higher luciferase reporter activity, respectively, than the pGL3-Basic empty vector in DF1 cells, and 180- and 120-fold higher luciferase reporter activity, respectively, than the pGL3-Basic empty vector in ICP1 cells, suggesting that the cloned PPAR $\gamma$ 1 5' UTR has promoter activity. By comparison, pGL3-PPAR $\gamma$ 1-Mut showed significantly lower luciferase activity than pGL3-PPAR $\gamma$ 1-WT in DF1 cells (Figure 5A,  $P < 0.05$ ) and ICP1 cells (Figure 5B,  $P < 0.01$ ). These results demonstrated that the cloned PPAR $\gamma$ 1 5' UTR had strong promoter activity and that the uORF mutation can result in reduced promoter activity. These findings explain why the mRNA expression levels of *Rluc* and *PPAR $\gamma$ 1* were reduced in the above study (Figures 2C, 3C).

### The Effect of the uORF on *in vitro* Translation of PPAR $\gamma$ 1

The results reported here suggest that at the mRNA level, the uORF represses PPAR $\gamma$ 1 translation (Figures 3A,B). To further



**FIGURE 5 |** The promoter activity analysis of the DNA sequences corresponding wild-type and uORF-mutated 5' UTRs of PPAR $\gamma$ 1. The DNA sequences corresponding wild-type and uORF-mutated 5' UTRs of PPAR $\gamma$ 1 were cloned into luciferase reporter vector pGL3-basic to yield pGL3-PPAR $\gamma$ 1-WT and pGL3-PPAR $\gamma$ 1-Mut, respectively. The indicated reporters along with the pRL-TK *Renilla* luciferase vector were transiently transfected into DF1 (A) and ICP1 cells (B), and the luciferase activity was determined at 48 h after transfection. The pRL-TK vector was used for normalization of transfection efficiency. All data represent the mean  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$ , Student's *t*-test.

validate this finding, we performed an *in vitro* transcription and translation assay. For *in vitro* transcription, equal amounts (1  $\mu$ g) of linearized pcDNA3.1-PPAR $\gamma$ 1-WT or pcDNA3.1-PPAR $\gamma$ 1-Mut were used as templates to produce the PPAR $\gamma$ 1 mRNA with wild-type and uORF-mutated 5' UTR using the T7 RiboMAX Large Scale RNA Production System. The results showed that, as expected, pcDNA3.1-PPAR $\gamma$ 1-WT and pcDNA3.1-PPAR $\gamma$ 1-Mut produced almost the same amount of PPAR $\gamma$ 1 mRNA (Figure 6A). Equal amounts of the transcribed PPAR $\gamma$ 1 mRNA produced from pcDNA3.1-PPAR $\gamma$ 1-WT and pcDNA3.1-PPAR $\gamma$ 1-Mut were used for the *in vitro* translation assay. The *in vitro* translation assay results showed that more PPAR $\gamma$ 1 protein was synthesized with PPAR $\gamma$ 1 mRNA from pcDNA3.1-PPAR $\gamma$ 1-Mut than from pcDNA3.1-PPAR $\gamma$ 1-WT (Figure 6B). Together, these results indicate that the uORF represses PPAR $\gamma$ 1 translation.

### The PPAR $\gamma$ 1 uORF Can Be Translated

To gain insight into the molecular mechanism by which the uORF represses translation, we tested whether the uAUG of PPAR $\gamma$ 1 uORF was used for translation initiation. We generated a construct in which the uORF was fused in frame with the enhanced green fluorescent protein (EGFP)

coding sequences, with no intervening in-frame stop codons, and named it pcDNA3.1-uORF-EGFP; pcDNA3.1-EGFP was used as a positive control. The ICP1 cells were transiently transfected with pcDNA3.1-uORF-EGFP or pcDNA3.1-EGFP and examined by microscopy and western blotting. Microscopy showed that the cells transfected with either pcDNA3.1-uORF-EGFP or pcDNA3.1-EGFP displayed GFP fluorescence (Figure 7A). Comparatively, GFP fluorescence intensity was lower in the cells transfected with pcDNA3.1-uORF-EGFP than with pcDNA3.1-EGFP (Figure 7A). Consistent with these findings, western blot analysis showed that the uORF-EGFP fusion protein was expressed but at a lower level in the cells transfected with pcDNA3.1-uORF-EGFP compared with that in the cells transfected with pcDNA3.1-EGFP (Figure 7B). Collectively, these data suggest that translation can indeed be initiated at the uAUG.


### DISCUSSION

Investigating the molecular mechanisms that control PPAR $\gamma$  expression is critical for understanding adipogenesis, as well as pathological conditions such as obesity and diabetes. In the present study, we investigated PPAR $\gamma$  posttranscriptional regulation by 5' UTR. We demonstrated that a uORF, which is absent in human and mouse PPAR $\gamma$  transcripts, represses chicken PPAR $\gamma$  transcript variant 1 translation. To our knowledge, this is the first report of a uORF regulating PPAR $\gamma$  gene expression.

In the present study, we demonstrated that the five chicken PPAR $\gamma$  5' UTR isoforms exerted different effects on the reporter gene activity (Figures 1A,B), and further study showed the 5' UTR uORF of PPAR $\gamma$ 1 represses reporter gene and PPAR $\gamma$ 1 translation. Sequence analysis revealed that the uAUG of the PPAR $\gamma$ 1 uORF resides in a favorable Kozak sequence context, which is the most efficient context for ribosome recognition and initiation of translation (Figure 2A). In agreement with the bioinformatics prediction, we demonstrated that the uAUG could serve as a translation start site (Figures 7A,B). Furthermore, secondary structural analysis showed there was a stable loop structure within the uORF (Supplementary Figure 1), and the uORF mutation (AUG > UAG) was not able to alter the secondary structure of PPAR $\gamma$ 1 5' UTR (Supplementary Figure 1). Thus, we could rule out an effect of secondary structure alteration on PPAR $\gamma$ 1 expression.


The 17-amino acid PPAR $\gamma$ 1 uORF peptide was analyzed using the PSIPRED protein sequence analysis workbench. It was predicted to be a disordered peptide (Romero et al., 2001; Buchan et al., 2013). Disordered peptides are enriched with residues Gly, Pro, Arg, and Ser, which are potential targets for phosphorylation that could promote ribosome stalling during translation elongation or termination (Hayden and Jorgensen, 2007; Johansson et al., 2011; Koutmou et al., 2015), which may explain why EGFP expression was lower in the cells transfected with pcDNA3.1-uORF-EGFP than with pcDNA3.1-EGFP (Figures 7A,B).

An increasing number of uORF-encoded peptides have been identified and shown to repress the downstream ORF expression



**FIGURE 6 |** The uORF represses *in vitro* PPAR $\gamma$ 1 translation. **(A)** *In vitro* transcribed PPAR $\gamma$ 1 mRNAs from the wild-type and uORF-mutant PPAR $\gamma$ 1 expression vectors (pcDNA3.1-PPAR $\gamma$ 1-WT and pcDNA3.1-PPAR $\gamma$ 1-Mut) were analyzed by quantitative real-time RT-PCR. No difference in PPAR $\gamma$ 1 mRNA was observed. Data were expressed as the mean  $\pm$  SEM, n.s., not significant, Student's *t*-test. **(B)** Equal amounts of the *in vitro* transcribed mRNAs (2  $\mu$ g) were used for *in vitro* translation. Note that the uORF strongly represses PPAR $\gamma$ 1 translation. A *in vitro* translation reaction without RNA template was used as a negative control.






**FIGURE 7 |** Translation can be initiated at the uAUG of the PPAR $\gamma$ 1 uORF. **(A)** The pcDNA3.1-EGFP and pcDNA3.1-uORF-EGFP were respectively transiently transfected into ICP1 cells, 48 h post-transfection, the green fluorescence signal was visualized under a fluorescence microscope. **(B)** Lysates from the cells transfected with pcDNA3.1-EGFP and pcDNA3.1-uORF-EGFP, EGFP or uORF-EGFP fusion protein was immunoblotted with an anti-EGFP antibody. Actin was used as a loading control.

by triggering ribosome stalling or suppressing reinitiation (Wilson and Beckmann, 2011; Ito and Chiba, 2013; Starck et al., 2016). Our data demonstrated that the PPAR $\gamma$ 1 uORF repressed downstream ORF expression and that it could be translated. This raised the question of whether this uORF-encoded peptide represses downstream ORF translation. To this end, we constructed a uORF expression vector, pcDNA3.1-uORF, and co-transfected pcDNA3.1-uORF and PPAR $\gamma$ 1-5'UTR-WT or PPAR $\gamma$ 1-5'UTR-Mut into DF1 cells. Unexpectedly, reporter gene assays showed that transfection of pcDNA3.1-uORF increased reporter gene activities of both PPAR $\gamma$ 1-5'UTR-WT and PPAR $\gamma$ 1-5'UTR-Mut (**Supplementary Figure 2**). This result suggested that this uORF-encoded peptide may repress the downstream PPAR $\gamma$ 1 translation in *cis*, but not in *trans*.

It has been reported that several uORF-encoded peptides act in *cis* on the ribosome during their own translation to stall translation. Arrest of translation can occur either during translation elongation, as seen for SecM (Tsai et al., 2014) and VemP (Vazquez-Laslop et al., 2008), or during translation termination; for example, in the tryptophanase C (TnaC) (Gong et al., 2001) and S-adenosyl-methionine decarboxylase (SAM-DC) (Raney et al., 2002).

Based on our data, we speculated that uORF repressed PPAR $\gamma$ 1 translation by two possible mechanisms. The first was ribosome stalling (**Figure 8A**), in which uAUG is recognized by the scanning 40S ribosomal subunit and associated initiation factors, the uORF is translated, and the nascent peptide stalls the ribosome in the ribosome exit tunnel, thereby hampering



the progression of upstream ribosomes (Wilson, 2011; Brandman et al., 2012; Wilson et al., 2016). Only a tiny minority of ribosomes may leaky-scan the uORF start codon and translate the PPAR $\gamma$ 1 coding sequence. Consequently, the translational efficiency of PPAR $\gamma$ 1 is dramatically attenuated.

The other possible mechanism is translational reinitiation, in which ribosomes translate the uORF and remain associated with the mRNA, continue scanning, and reinitiate further downstream at either a proximal or distal AUG codon (Figure 8B). However, reinitiation efficiency is substantially reduced (Roy et al., 2010; Hinnebusch et al., 2016) and translation of PPAR $\gamma$ 1 inhibited. Recently, some nascent peptides of uORFs have been reported to be involved in the suppression of reinitiation (Ito and Chiba, 2013; Seefeldt et al., 2015). We speculate that the uORF-encoded peptide may contribute to suppression of PPAR $\gamma$ 1 translation.

In addition, several studies have implicated that uORF-containing mRNA has the potential to trigger the nonsense-mediated decay (NMD) pathway. NMD is one of the better characterized posttranscriptional control mechanisms, whereby transcripts harboring premature translation termination codons are degraded (Mendell et al., 2004). In the present study, we detected no significant effect of uORF mutation on *Rluc* mRNA stability (Figures 4, 6A). Therefore, we can rule out the possibility that PPAR $\gamma$ 1 uORF modulates PPAR $\gamma$ 1 expression by triggering the NMD pathway.

PPAR $\gamma$  is a master regulator of adipogenesis, whole-body lipid metabolism, and insulin sensitivity. Accumulating evidence shows that adipogenesis and lipid metabolism are different between mammals and chickens (Prigge and Grande, 1971; Ji et al., 2012). For example, unlike that in mammals, chicken adipocyte lipolysis is almost exclusively regulated by glucagon, and chicken adipose tissue is not sensitive to insulin (Dupont et al., 2012; Ji et al., 2012; Wang et al., 2017). In the present study, we demonstrated that the uORF represses chicken PPAR $\gamma$ 1 translation, and bioinformatics analysis showed that the 5' UTR sequences of PPAR $\gamma$  transcripts had low sequence similarity among humans, mice, and chickens, and no uORF element was identified in the 5' UTRs of human and mouse PPAR $\gamma$  transcripts. These data suggest that the

posttranscriptional regulation of the PPAR $\gamma$  gene by the 5' UTR differs between mammals and chickens. The identified uORF element is a unique regulatory element in chicken PPAR $\gamma$  gene, which may contribute to the differences in adipogenesis, adipose development, and insulin sensitivity between mammals and chickens. Our present study has two limitations. First, the scope of our present study is small; we just explored the regulatory role and possible mechanism of PPAR $\gamma$ 1 uORF. Second, we did not investigate the *in vivo* importance of the PPAR $\gamma$ 1 uORF in the regulation of PPAR $\gamma$  expression due to the fact that chicken gene knockout and transgenic approaches have not been well established. Given the importance of PPAR $\gamma$  in adipose development, obesity and related diseases, in the future we will use CRISPR-Cas9 technology to explore the *in vivo* importance and underlying molecular mechanism of PPAR $\gamma$ 1 uORF in physiological and pathological conditions. A better understanding of PPAR $\gamma$  5' UTRs may provide clues for controlling obesity, type 2 diabetes, and insulin resistance.

In summary, for the first time, we demonstrated that a uORF represses chicken PPAR $\gamma$ 1 translation.

## DATA AVAILABILITY STATEMENT

Publicly available datasets were analyzed in this study. This data can be found here: GenBank: KP736526.1 (<https://www.ncbi.nlm.nih.gov/nucleotide/KP736526.1/>).

## AUTHOR CONTRIBUTIONS

YC designed the experiments, collected and analyzed the data, and wrote the manuscript. JH, GM, TC, XY, and HL participated in scientific discussions and provided technical assistances. NW supervised the study and wrote the manuscript with YC.

## FUNDING

This work was supported by the National Natural Science Foundation of China (No. 31572392) and the China Agriculture Research System (No. CARS-41).

## ACKNOWLEDGMENTS

This manuscript has been released as a Pre-print at BioRxiv (<https://www.biorxiv.org/content/10.1101/858753v1>) (Chu et al., 2019).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2020.00165/full#supplementary-material>

## REFERENCES

- Ahmadian, M., Suh, J. M., Hah, N., Liddle, C., Atkins, A. R., Downes, M., et al. (2013). PPAR $\gamma$  signaling and metabolism: the good, the bad and the future. *Nat. Med.* 19, 557–566. doi: 10.1038/nm.3159
- Aprile, M., Ambrosio, M. R., D'Esposito, V., Beguinot, F., Formisano, P., Costa, V., et al. (2014). PPAR $\gamma$  in human adipogenesis: differential contribution of canonical transcripts and dominant negative isoforms. *PPAR Res.* 2014:537865. doi: 10.1155/2014/537865
- Araujo, P. R., Yoon, K., Ko, D., Smith, A. D., Qiao, M., Suresh, U., et al. (2012). Before it gets started: regulating translation at the 5' UTR. *Comp. Funct. Genomics* 2012:475731. doi: 10.1155/2012/475731
- Barbosa, C., Peixeiro, I., and Romao, L. (2013). Gene expression regulation by upstream open reading frames and human disease. *PLoS Genet.* 9:e1003529. doi: 10.1371/journal.pgen.1003529
- Brandman, O., Stewart-Ornstein, J., Wong, D., Larson, A., Williams, C. C., Li, G. W., et al. (2012). A ribosome-bound quality control complex triggers degradation of nascent peptides and signals translation stress. *Cell* 151, 1042–1054. doi: 10.1016/j.cell.2012.10.044
- Buchan, D. W., Minnici, F., Nugent, T. C., Bryson, K., and Jones, D. T. (2013). Scalable web services for the PSIPRED protein analysis workbench. *Nucleic Acids Res.* 41, W349–W357. doi: 10.1093/nar/gkt381
- Bugaut, A., and Balasubramanian, S. (2012). 5'-UTR RNA G-quadruplexes: translation regulation and targeting. *Nucleic Acids Res.* 40, 4727–4741. doi: 10.1093/nar/gks068
- Calvo, S. E., Pagliarini, D. J., and Mootha, V. K. (2009). Upstream open reading frames cause widespread reduction of protein expression and are polymorphic among humans. *Proc. Natl. Acad. Sci. U.S.A.* 106, 7507–7512. doi: 10.1073/pnas.0810916106
- Cariou, B., Charbonnel, B., and Staels, B. (2012). Thiazolidinediones and PPAR $\gamma$  agonists: time for a reassessment. *Trends Endocrinol. Metab.* 23, 205–215. doi: 10.1016/j.tem.2012.03.001
- Ceraj, I., T.J. Riley, and Shubert, C. (2009). "StarHPC – Teaching parallel programming within elastic compute cloud," in *Proceedings of the ITI 2009 31st International Conference on Information Technology Interfaces*, Cavtat.
- Chandra, M., Miriyala, S., and Panchatcharam, M. (2017). PPAR $\gamma$  and its role in cardiovascular diseases. *PPAR Res.* 2017:6404638. doi: 10.1155/2017/6404638
- Chang, T. H., Huang, H. Y., Hsu, B. K., Weng, S. L., Horng, J. T., and Huang, H. D. (2013). An enhanced computational platform for investigating the roles of regulatory RNA and for identifying functional RNA motifs. *BMC Bioinformatics* 14(Suppl. 2):S4–S4. doi: 10.1186/1471-2105-14-S2-S4
- Chu, Y., Huang, J., Ma, G., Cui, T., Yan, X., Li, H., et al. (2019). An upstream open reading frame represses translation of chicken PPAR $\gamma$  transcript variant 1. *bioRxiv* doi: 10.1101/858753
- Crowe, M. L., Wang, X. Q., and Rothnagel, J. A. (2006). Evidence for conservation and selection of upstream open reading frames suggests probable encoding of bioactive peptides. *BMC Genomics* 7:16. doi: 10.1186/1471-2164-7-16
- Cui, T., Xing, T., Huang, J., Mu, F., Jin, Y., You, X., et al. (2018). Nuclear respiratory factor 1 negatively regulates the p1 promoter of the peroxisome proliferator-activated receptor- $\gamma$  gene and inhibits chicken adipogenesis. *Front. Physiol.* 9:1823. doi: 10.3389/fphys.2018.01823
- Dikstein, R. (2012). Transcription and translation in a package deal: the TISU paradigm. *Gene* 491, 1–4. doi: 10.1016/j.gene.2011.09.013
- Duan, K., Sun, Y., Zhang, X., Zhang, T., Zhang, W., Zhang, J., et al. (2015). Identification and characterization of transcript variants of chicken peroxisome proliferator-activated receptor  $\gamma$ . *Poult Sci.* 94, 2516–2527. doi: 10.3382/ps/pev229
- Dupont, J., Metayer-Coustard, S., Ji, B., Rame, C., Gespach, C., Voy, B., et al. (2012). Characterization of major elements of insulin signaling cascade in chicken adipose tissue: apparent insulin refractoriness. *Gen. Compar. Endocrinol.* 176, 86–93. doi: 10.1016/j.ygcen.2011.12.030
- Dvir, S., Velten, L., Sharon, E., Zeevi, D., Carey, L. B., Weinberger, A., et al. (2013). Deciphering the rules by which 5'-UTR sequences affect protein expression in yeast. *Proc. Natl. Acad. Sci. U.S.A.* 110, E2792–E2801. doi: 10.1073/pnas.1222534110
- Gong, F., Ito, K., Nakamura, Y., Yanofsky, C., et al. (2001). The mechanism of tryptophan induction of tryptophanase operon expression: tryptophan inhibits release factor-mediated cleavage of TnaC-peptidyl-tRNA(Pro). 98, 8997–9001. doi: 10.1073/pnas.171299298
- Grillo, G., Turi, A., Licciulli, F., Mignone, F., Liuni, S., Banfi, S., et al. (2010). UTRdb and UTRsite (RELEASE 2010): a collection of sequences and regulatory motifs of the untranslated regions of eukaryotic mRNAs. *Nucleic Acids Res.* 38, D75–D80. doi: 10.1093/nar/gkp902
- Hayden, C. A., and Jorgensen, R. A. (2007). Identification of novel conserved peptide uORF homology groups in *Arabidopsis* and rice reveals ancient eukaryotic origin of select groups and preferential association with transcription factor-encoding genes. *BMC Biol.* 5:32. doi: 10.1186/1741-7007-5-32
- Hinnebusch, A. G., Ivanov, I. P., and Sonenberg, N. (2016). Translational control by 5'-untranslated regions of eukaryotic mRNAs. *Science* 352, 1413–1416. doi: 10.1126/science.aad9868
- Hofacker, I. L. (2003). Vienna RNA secondary structure server. *Nucleic Acids Res.* 31, 3429–3431. doi: 10.1093/nar/gkg599
- Hood, H. M., Neafsey, D. E., Galagan, J., and Sachs, M. S. (2009). Evolutionary roles of upstream open reading frames in mediating gene regulation in fungi. *Annu. Rev. Microbiol.* 63, 385–409. doi: 10.1146/annurev.micro.62.081307.162835
- Iacono, M., Mignone, F., and Pesole, G. (2005). uAUG and uORFs in human and rodent 5'-untranslated mRNAs. *Gene* 349, 97–105. doi: 10.1016/j.gene.2004.11.041
- Ito, K., and Chiba, S. (2013). Arrest peptides: cis-acting modulators of translation. *Annu. Rev. Biochem.* 82, 171–202. doi: 10.1146/annurev-biochem-080211-105026
- Ji, B., Ernest, B., Gooding, J. R., Das, S., Saxton, A. M., Simon, J., et al. (2012). Transcriptomic and metabolomic profiling of chicken adipose tissue in response to insulin neutralization and fasting. *BMC Genomics* 13:441. doi: 10.1186/1471-2164-13-441
- Johansson, M., Jeong, K. W., Trobro, S., Strazewski, P., Aqvist, J., Pavlov, M. Y., et al. (2011). pH-sensitivity of the ribosomal peptidyl transfer reaction dependent on the identity of the A-site aminoacyl-tRNA. *Proc. Natl. Acad. Sci. U.S.A.* 108, 79–84. doi: 10.1073/pnas.1012612107
- Katsura, S., Okumura, T., Ito, R., Sugawara, A., and Yokoyama, A. (2014). Identification of posttranslational modifications in peroxisome proliferator-activated receptor  $\gamma$  using mass spectrometry. *PPAR Res.* 2014:468925. doi: 10.1155/2014/468925
- Koutmou, K. S., Schuller, A. P., Brunelle, J. L., Radhakrishnan, A., Djuranovic, S., and Green, R. (2015). Ribosomes slide on lysine-encoding homopolymeric A stretches. *Elife* 4:e05534. doi: 10.7554/eLife.05534
- Lee, J. E., and Ge, K. (2014). Transcriptional and epigenetic regulation of PPAR $\gamma$  expression during adipogenesis. *Cell Biosci.* 4:29. doi: 10.1186/2045-3701-4-29
- Lefterova, M. I., Haakonsson, A. K., Lazar, M. A., and Mandrup, S. (2014). PPAR $\gamma$  and the global map of adipogenesis and beyond. *Trends Endocrinol. Metab.* 25, 293–302. doi: 10.1016/j.tem.2014.04.001
- Lehrke, M., and Lazar, M. A. (2005). The many faces of PPAR $\gamma$ . *Cell* 123, 993–999. doi: 10.1016/j.cell.2005.11.026
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-</sup>(Delta Delta C(T)) method. *Methods* 25, 402–408. doi: 10.1006/meth.2001.1262
- McClelland, S., Shrivastava, R., and Medh, J. D. (2014). Regulation of translational efficiency by disparate 5' UTRs of PPAR $\gamma$  splice variants. *Ppar Res.* 2009:193413. doi: 10.1155/2009/193413
- Mendell, J. T., Sharifi, N. A., Meyers, J. L., Martinez-Murillo, F., and Dietz, H. C. (2004). Nonsense surveillance regulates expression of diverse classes of mammalian transcripts and mutes genomic noise. *Nat. Genet.* 36, 1073–1078. doi: 10.1038/ng1429
- Mitchell, S. F., Walker, S. E., Algire, M. A., Park, E. H., Hinnebusch, A. G., and Lorsch, J. R. (2010). The 5'-7-methylguanosine cap on eukaryotic mRNAs serves both to stimulate canonical translation initiation and to block an alternative pathway. *Mol. Cell.* 39, 950–962. doi: 10.1016/j.molcel.2010.08.021
- Nasif, S., Contu, L., and Muhlemann, O. (2018). Beyond quality control: the role of nonsense-mediated mRNA decay (n.d.) in regulating gene expression. *Semin. Cell Dev Biol.* 75, 78–87. doi: 10.1016/j.semcdb.2017.08.053

- Normann, K. R., Øystese, K. A. B., Berg, J. P., Lekva, T., Berg-Johnsen, J., Bollerslev, J., et al. (2016). Selection and validation of reliable reference genes for RT-qPCR analysis in a large cohort of pituitary adenomas. *Mol. Cell. Endocrinol.* 437(C), 183–189. doi: 10.1016/j.mce.2016.08.030
- Prigge, W. F., and Grande, F. (1971). Effects of glucagon, epinephrine and insulin on in vitro lipolysis of adipose tissue from mammals and birds. *Compar. Biochem. Physiol. B* 39, 69–82. doi: 10.1016/0305-0491(71)90254-9
- Raney, A., Law, G. L., Mize, G. J., and Morris, D. R. (2002). Regulated translation termination at the upstream open reading frame in s-adenosylmethionine decarboxylase mRNA. *J. Biol. Chem.* 277, 5988–5994. doi: 10.1074/jbc.M1108375200
- Romero, P., Obradovic, Z., Li, X., Garner, E. C., Brown, C. J., and Dunker, A. K. (2001). Sequence complexity of disordered protein. *Proteins* 42, 38–48.
- Roy, B., Vaughn, J. N., Kim, B. H., Zhou, F., Gilchrist, M. A., and Von Arnim, A. G. (2010). The h subunit of eIF3 promotes reinitiation competence during translation of mRNAs harboring upstream open reading frames. *RNA* 16, 748–761. doi: 10.1261/rna.2056010
- Sathirapongsasuti, J. F., Sathira, N., Suzuki, Y., Huttenhower, C., and Sugano, S. (2011). Ultraconserved cDNA segments in the human transcriptome exhibit resistance to folding and implicate function in translation and alternative splicing. *Nucleic Acids Res.* 39, 1967–1979. doi: 10.1093/nar/gkq949
- Seefeldt, A. C., Nguyen, F., Antunes, S., Perebaskine, N., Graf, M., Arenz, S., et al. (2015). The proline-rich antimicrobial peptide Onc112 inhibits translation by blocking and destabilizing the initiation complex. *Nat. Struct. Mol. Biol.* 22, 470–475. doi: 10.1038/nsmb.3034
- Starck, S. R., Tsai, J. C., Chen, K., Shodiya, M., Wang, L., Yahiro, K., et al. (2016). Translation from the 5' untranslated region shapes the integrated stress response. *Science* 351:aad3867. doi: 10.1126/science.aad3867
- Suzuki, Y., Ishihara, D., Sasaki, M., Nakagawa, H., Hata, H., Tsunoda, T., et al. (2000). Statistical analysis of the 5' untranslated region of human mRNA using "oligo-capped" cDNA libraries. *Genomics* 64, 286–297. doi: 10.1006/geno.2000.6076
- Tsai, A., Kornberg, G., Johansson, M., Chen, J., and Puglisi, J. (2014). The dynamics of secm-induced translational stalling. *Cell Rep.* 7, 1521–1533. doi: 10.1016/j.celrep.2014.04.033
- van Beekum, O., Fleskens, V., and Kalkhoven, E. (2009). Posttranslational modifications of PPAR-gamma: fine-tuning the metabolic master regulator. *Obesity (Silver Spring)* 17, 213–219. doi: 10.1038/oby.2008.473
- Vazquez-Laslop, N., Thum, C., and Mankin, A. S. (2008). Molecular mechanism of drug-dependent ribosome stalling. *Mol. Cell* 30, 190–202. doi: 10.1016/j.molcel.2008.02.026
- Wang, W., Zhang, T., Wu, C., Wang, S., Wang, Y., Li, H., et al. (2017). immortalization of chicken preadipocytes by retroviral transduction of chicken TERT and TR. *PLoS ONE* 12:e0177348. doi: 10.1371/journal.pone.0177348
- Wen, Y., Liu, Y., Xu, Y., Zhao, Y., Hua, R., Wang, K., et al. (2009). Loss-of-function mutations of an inhibitory upstream ORF in the human hairless transcript cause Marie Unna hereditary hypotrichosis. *Nat. Genet.* 41, 228–233. doi: 10.1038/ng.276
- Wilson, D. N. (2011). Peptides in the ribosomal tunnel talk back. *Mol. Cell* 41, 247–248. doi: 10.1016/j.molcel.2011.01.017
- Wilson, D. N., Arenz, S., and Beckmann, R. (2016). Translation regulation via nascent polypeptide-mediated ribosome stalling. *Curr. Opin. Struct. Biol.* 37, 123–133. doi: 10.1016/j.sbi.2016.01.008
- Wilson, D. N., and Beckmann, R. (2011). The ribosomal tunnel as a functional environment for nascent polypeptide folding and translational stalling. *Curr. Opin. Struct. Biol.* 21, 274–282. doi: 10.1016/j.sbi.2011.01.007
- Xia, X., and Holcik, M. (2009). Strong eukaryotic IRESs have weak secondary structure. *PLoS ONE* 4:e4136. doi: 10.1371/journal.pone.0004136
- Yamashita, R., Suzuki, Y., Takeuchi, N., Wakaguri, H., Ueda, T., Sugano, S., et al. (2008). Comprehensive detection of human terminal oligo-pyrimidine (TOP) genes and analysis of their characteristics. *Nucleic Acids Res.* 36, 3707–3715. doi: 10.1093/nar/gkn248

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Chu, Huang, Ma, Cui, Yan, Li and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.