



# Sources of Pathogenic Nucleic Acids in Systemic Lupus Erythematosus

Tomas Mustelin\*, Christian Lood and Natalia V. Giltiy

Division of Rheumatology, Department of Medicine, University of Washington, Seattle, WA, United States

## OPEN ACCESS

### Edited by:

Reinhard Edmund Voll,  
Freiburg University Medical Center,  
Germany

### Reviewed by:

Mary K. Crow,  
Hospital for Special Surgery,  
United States  
Kevin M. Nickerson,  
University of Pittsburgh, United States

### \*Correspondence:

Tomas Mustelin  
tomas2@uw.edu

### Specialty section:

This article was submitted to  
Autoimmune and Autoinflammatory  
Disorders,  
a section of the journal  
Frontiers in Immunology

**Received:** 15 December 2018

**Accepted:** 23 April 2019

**Published:** 08 May 2019

### Citation:

Mustelin T, Lood C and Giltiy NV  
(2019) Sources of Pathogenic Nucleic  
Acids in Systemic Lupus  
Erythematosus.  
*Front. Immunol.* 10:1028.  
doi: 10.3389/fimmu.2019.01028

A hallmark of systemic lupus erythematosus (SLE), and several related autoimmune diseases, is the presence of autoantibodies against nucleic acids and nucleic acid-binding proteins, as well as elevated type I interferons (IFNs), which appear to be instrumental in disease pathogenesis. Here we discuss the sources and proposed mechanisms by which a range of cellular RNA and DNA species can become pathogenic and trigger the nucleic acid sensors that drive type I interferon production. Potentially SLE-promoting DNA may originate from pieces of chromatin, from mitochondria, or from reverse-transcribed cellular RNA, while pathogenic RNA may arise from mis-localized, mis-processed, ancient retroviral, or transposable element-derived transcripts. These nucleic acids may leak out from dying cells to be internalized and reacted to by immune cells or they may be generated and remain to be sensed intracellularly in immune or non-immune cells. The presence of aberrant DNA or RNA is normally counteracted by effective counter-mechanisms, the loss of which result in a serious type I IFN-driven disease called Aicardi-Goutières Syndrome. However, in SLE it remains unclear which mechanisms are most critical in precipitating disease: aberrant RNA or DNA, overly sensitive sensor mechanisms, or faulty counter-acting defenses. We propose that the clinical heterogeneity of SLE may be reflected, in part, by heterogeneity in which pathogenic nucleic acid molecules are present and which sensors and pathways they trigger in individual patients. Elucidation of these events may result in the recognition of distinct “endotypes” of SLE, each with its distinct therapeutic choices.

**Keywords:** lupus, interferon, nucleic acid sensors, mitochondria, reverse transcriptase

## INTRODUCTION

Systemic lupus erythematosus (SLE) is a serious autoimmune disease characterized by autoantibodies against nucleic acids and nucleic acid-binding proteins combined with immune complex deposition and inflammatory manifestations in multiple organ systems. The unpredictable course of the disease with its sudden exacerbations, often with new organ manifestations or symptoms, make it particularly difficult to manage (1, 2), not the least because the currently available drugs have limited efficacy and/or serious side-effects. Efforts to develop more selective and more efficacious therapies that address the core pathobiology of SLE, ideally with limited general immune suppression, continue to be hampered by our limited understanding of the underlying molecular drivers and mechanisms (3). To vividly illustrate this, the two newest therapeutics for SLE are hydroxychloroquine (4, 5) and belimumab (6, 7), approved by the FDA in 1966 (sic!) and 2011, respectively. Moreover, the latter had barely significant efficacy, only 9.8% SLE Responder Index improvement over placebo at 52 weeks at the highest 10 mg/kg dose (6).

Furthermore, while the presence of autoantibodies in SLE has been recognized for decades and their role in driving disease is considered well established, B cell depletion by anti-CD20 antibodies have failed to generate statistically significant efficacy in clinical trials in SLE (8). There is, however, a trend toward a benefit for patients in agreement with the ability of belimumab to reduce B cell numbers. It should also be noted that belimumab may affect plasma cells more than the depletion of CD20-positive B cells. T cell-directed therapies, such as calcineurin inhibitors (9) or CD28 blockade with CTLA4-Ig (10), have also yielded limited disease impact. These outcomes suggest that many of the well-documented immune abnormalities in SLE may be consequences, rather than drivers, of this disease.

## AUTOANTIBODIES AGAINST NUCLEIC ACIDS AND NUCLEIC ACID-BINDING PROTEINS

In SLE, the majority of patients develop autoimmunity toward nuclear antigens, conveniently measured as anti-nuclear autoantibodies (ANA). Though not selective for SLE, detecting ANA is a common test used to screen patients, and may, together with clinical presentation and other immunological features, suffice for SLE diagnosis. ANA contains a broad range of autoantibodies targeting among others chromatin, histones, double-stranded (ds) DNA, as well as the RNA-binding proteins Ro, La, Sm, and RNP. Anti-dsDNA antibodies are of particular interest in SLE, given their high diagnostic potential, with about 70–80% of the patients being positive for these antibodies, and titers commonly correlating with disease activity. Indeed, anti-dsDNA antibodies have been included in the classification criteria (11), as well as a serological component of the disease activity index SLEDAI (12). Further, anti-dsDNA antibodies are often associated with severe disease manifestations, including nephritis. Other than the diagnostic value, including associations with distinct disease features, these autoantibodies may be pathogenic through immune complex-mediated inflammation, complement activation and tissue destruction, and antibody-directed cellular cytotoxicity. In this review, we will limit our discussion of autoantibodies to their ability to transport nucleic acids, shielding them from external nucleases, and efficiently mediating their uptake into immune cells through Fc receptors, complement receptors, scavenger receptors, and others.

## THE “IFN SIGNATURE” IN SLE PATIENTS

In 2003, Tim Behrens' group (13), the team of Virginia Pascual and Jacques Banchereau (14), and Mary Crow (15) published their discovery that SLE patient blood contain active type I interferon (IFN) and a high expression level of IFN-stimulated genes (ISGs), now referred to as the “IFN signature.” Although indications that IFN $\alpha$  may be important in the lupus pathogenesis had been published earlier (16–18), this still was a surprising finding because the principal function of type I IFN is in host defense against viral infection, while SLE is not an

infectious disease. Nevertheless, the IFN signature is now a well-established observation in 70–90% of SLE patient populations world-wide (19–22). Individual IFNs are technically difficult to measure (23) due to their very low concentrations and presumed rapid consumption, but it seems that many of the 17 different type I IFNs, which includes 13 IFN $\alpha$  isoforms, IFN $\beta$ , and the three less explored members, IFN $\epsilon$ , IFN $\kappa$ , and IFN $\omega$  are elevated in SLE patients, as well as in patients with Sjögren's syndrome (24, 25), systemic sclerosis (26, 27), polymyositis, dermatomyositis (28, 29), rheumatoid arthritis (30, 31), and other related diseases. Importantly, there seems to be differences between patients in which specific members of the type I IFN family are elevated (see sections Patient heterogeneity with regard to nucleic acids and their sensors? and Can SLE be divided into clinically meaningful subpopulations based on “endotype”?). In addition, patients may have increased type II IFN (IFN $\gamma$ ) (25) and/or type III IFNs (IFN $\lambda$ 1, IFN $\lambda$ 2, and IFN $\lambda$ 3, also known as IL-29, IL-28A, and IL-28B) (32). While the type I and III IFNs are functionally overlapping (all genes induced by type III IFNs are also induced by type I IFNs), IFN $\gamma$  is instrumental in a distinct aspect of the immune system, namely the activation of CD4 Th1 and CD8 T cells, natural killer (NK) cells, and other elements of a general immune response. Nevertheless, over 900 of the 1,300 ISGs induced by IFN $\gamma$  are also induced by type I IFNs, which induces a total of over 1,500 ISGs, suggesting significant overlap in downstream consequences.

Type I IFNs have a spectrum of effects on the immune system and beyond, particularly upregulating numerous mechanisms of on anti-viral defense. They stimulate emergency myelopoiesis (33), monocyte differentiation into myeloid dendritic cells (34, 35), antigen presentation, cytotoxic T cell differentiation (36), and B cell differentiation into plasma cells (37). The 1,500 ISGs encode many immune-modulating as well as direct antiviral proteins (38), including many components of the pathways that lead to type I IFN production in what constitutes a rapid positive feedback loop to augment the response.

While an extensive literature illuminates the close association of type I IFNs with SLE pathogenesis and disease activity (21, 39), perhaps the most conclusive evidence for a causal role in the disease was the statistically significant efficacy in phase 2 clinical trials (40) of an antibody that blocks the type I IFN receptor used by all type I IFNs. In contrast, an antibody that blocks IFN $\alpha$  alone (41) was efficacious only in a small subset of patients. It should also be noted that blocking the type I receptor did not bring clinical improvement to all SLE patients even if the IFN signature declined by over 90% in the treated patient population. Nevertheless, elevated type I IFNs are the closest we have to a smoking gun in SLE and a set of related autoimmune diseases. This, in turn, begs the question: why are type I IFNs elevated in SLE patients?

## NUCLEIC ACID SENSORS COUPLED TO INTERFERON PRODUCTION

Given that the best recognized role of type I IFN is in defense against viral infection (38), it seems that one could find important

clues about the upstream mechanisms of SLE from recent advances in viral immunity. The primary threat that a virus brings is its RNA or DNA genome, which will hijack the cellular biosynthetic machinery for its own replication and virion production, with detrimental consequences for the host cell. Even more alarming, retroviruses will reverse transcribe their RNA genome and insert the resulting DNA into the host genome as a permanent provirus. To combat these ancient foes, evolution has produced several cellular mechanisms for the detection of non-self RNA and DNA (**Figure 1**). Four principal pathways operate in the cytosol and on the surface of intracellular organelles: the DNA-sensor “cyclic GMP, AMP synthase” (cGAS) (42), the RNA sensors “retinoic acid-inducible gene I” (RIG-I) (43), “melanoma differentiation-associated gene 5” (MDA5) (43–45), and “RNA-activated protein kinase” (PKR) (46, 47), while a fifth pathway responds to extracellular DNA or RNA brought into the cell by receptor-mediated endocytosis and is initiated by Toll-like receptors (TLRs) 3, 7, 8, and 9 in the endosomal compartment. A mechanism to blend the extracellular and intracellular sensing pathways was recently reported (48): the transporter protein SIDT2 in the endosomal membrane functions to let dsRNA escape the endosome into the cytosol, where it can trigger MDA5. There are additional, more recently discovered nucleic acid sensors, such as DDX1, 21, 36, and 41, IFI16, and Aim2 (49). All of these pathways lead to type I IFN production through activation of IRF3 and related transcription factors. They also activate other signaling pathways that lead to the production of additional cytokines. The resulting type I IFNs are secreted, bind to the type I IFN receptor, and signal through the JAK/STAT pathways to upregulate ISG-encoded proteins with direct antiviral activity, including nucleases, helicases, chaperones, and many of the sensors and their adapters and signaling proteins (38). Type I IFN can act in both autocrine and paracrine fashion and the response to them may differ between different responding cell types.

Whereas, nucleic acids are the main triggers of type I IFN production, the cell type producing them and the exact nature of the triggering nucleic acid will determine which type I IFNs are produced. For example, plasmacytoid dendritic cells (pDC) have a particularly high capacity to produce several isoforms of IFN $\alpha$  in response to viruses or immune complexes that contain nucleic acids (50–52), including those containing IgE (52), by a TLR7 or 9-dependent mechanism. Non-immune cells, on the other hand, tend to produce predominantly IFN $\beta$  in response to cytosolic RNA or DNA through the sensors MDA5 (dsRNA), RIG-I (RNA), and cGAS (dsDNA), with other sensors participating, particularly in neutrophils that do not express cGAS (53).

## TLRs in SLE

TLRs are central to the immune system’s ability to recognize molecular structures associated with cellular damage or pathogens (54), including nucleic acids by TLRs 3, 7, 8, and 9 (**Figure 1A**). Since their discovery over 20 years ago, much of the early literature assumed that their role in SLE was certain (55–58), particularly since their ligation triggers type I IFN production and circulating immune complexes that contain nucleic acids are present in abundance in most SLE patients


(59), as well as in the patients with related diseases like Sjögren’s syndrome, polymyositis, dermatomyositis, mixed connective tissue disease, and others. It still seems very likely that these immune complexes drive production of IFN $\alpha$  by plasmacytoid dendritic cells (pDC) primarily through activation of TLR7 and maybe 9 (60). However, so far all tested antagonists of TLR 7 and/or 9 (61) have failed to provide any efficacy in placebo-controlled clinical trials in SLE patients. If type I IFNs indeed are important, but TLR7 and 9 inhibition does not produce a therapeutic benefit, then reality must be more complex. Indeed, the more recent discovery of other sensors for nucleic acid, such as cGAS, RIG-I, and MDA5 introduced other options for nucleic acid sensing leading to type I IFN. Nevertheless, it still seems likely that TLR7/9 drive IFN production in response to circulating immune complexes that contain nucleic acid and thereby contribute to the IFN signature seen in SLE patients. Unfortunately, clinical trials with TLR antagonists did not report what effects these drugs had on the IFN signature.

Recent advances in TLR research has revealed intriguing new details about the mechanisms of their ligand interactions, including their ability to bind self-nucleic acids (62–66). While TLR9 was originally proposed to only sense bacterial DNA with CpG sites, it is now clear that it can also recognize chromosomal and mitochondrial DNA (digested into small fragment by DNase II). Similarly, TLR3 responds to self-derived non-coding RNA, such as U1 RNA that might be released upon cellular stress, including exposure to UV radiation, while TLR7 and 8 can also recognize RNA and DNA degradation products (66). Another recent study found that phagocytosis of anti-dsDNA IgE antibodies (found to be increased in some SLE patients) via the high-affinity Fc $\epsilon$ RI receptor for IgE, mediates TLR9-mediated sensing of self-DNA in the phagosomes and potentiates IFN production by plasmacytoid dendritic cells (52).

Another potentially important aspect of the TLR pathways is that the *TLR7* and *TLR8* genes are located on the X-chromosome: there are indications that *TLR7* may escape the normal silencing of one of the two X chromosomes in females (67), resulting in higher levels of TLR7 expression and, hence, stronger responses to TLR7 stimulation in immune cells in women, perhaps contributing to the 9:1 gender bias in SLE. In further support of a role of TLR7 quantity in the disease, copy number variations (68, 69) and single-gene polymorphisms (70) in *TLR7* are associated with SLE susceptibility.

## Activation of cGAS and RNA Sensors in SLE

A recent paper provided the first direct evidence that the cGAS pathway is activated in at least a subset of SLE patients: the second messenger cyclic-guanine, adenosine-2,3-phosphate (cGAMP), which is synthesized exclusively by cGAS upon DNA binding, was detected by mass spectrometry in 7 of 30 SLE patients (71). While it may seem that this represents a small portion of SLE patients, it is important to recognize that the data represent a single snap-shot in time and that cGAMP is a short-lived second messenger present in minute quantities. Thus, it may well be that cGAMP is elevated in more SLE patients.



Direct evidence for activation of RNA sensors in SLE patients was also reported recently (72). Twenty two of sixty-seven examined SLE patients had evidence of polymerization of the mitochondrial antiviral signaling protein MAVS, which is downstream of both RIG-I and MDA5 (Figure 1B) and acts by generating a protein complex that activates the kinases required for IRF3 activation and type I IFN production. This aggregation of MAVS indicates that either RNA sensor was triggered in 32% of the patients.

## AICARDI-GOUTIÈRES SYNDROME—A MONOGENIC DISEASE OF NUCLEIC ACID PROCESSING

Additional insights into the molecular mechanisms that can drive type I interferons and cause interferon-dependent human disease come from a monogenic inherited inflammatory syndrome called Aicardi-Goutières syndrome (AGS) (73–77), which, together with a few related diseases, is included in the concept of the “type I interferonopathies” (78). AGS usually presents neonatally as a suspected serious viral infection with fever, chills, and a failure to thrive, accompanied by high levels of type I IFNs. However, a virus is not detected and the symptoms continue unabated. Over time, AGS patients develop neurological deficits and brain

calcifications, likely due to the neurotoxicity of IFNs, as well as systemic autoimmunity with autoantibodies against nucleic acids and nucleic acid-binding proteins very similar to those in SLE patients. In fact, many AGS patients meet the diagnostic criteria for SLE (73–77).

AGS is caused by mutations in any one of eight genes: *TREX1*, *RNASEH2A*, *RNASEH2B*, *RNASEH2C*, *SAMHD1*, *ADAR1*, *IFIH1*, *TMEM173* (73–77). The first 5 of these genes are primarily involved in the defense against retroviruses and their endogenous remnants in our genome (79). In fact, many of these genes were first discovered as “restriction factors” by researchers studying how HIV replicates in certain cells, but not in others. The revelation that our genome contains many evolutionarily conserved genes that confer resistance to HIV suggested that HIV is not the first exogenous retrovirus to infect us, but, in fact, is just the latest in a very long series of retroviral infections resulting in germline integrations of numerous families of retroviruses that today constitute as much as 8% of our genome (or as much as ~40% if other retroelements of ancient retroviral origin are also counted).

The three other genes that can induce AGS are homozygous loss-of-function mutations in the gene for “adenosine deaminase acting on RNA 1” (*ADAR1*), *IFIH1* which encodes MDA5, and the gain-of-function variant of *TMEM173* which encodes constitutively active STING (the direct effector protein for

cGAS, **Figure 1C**) and results in the constitutive activation of this pathway in the absence of aberrant DNA (80). The two latter genes demonstrate that chronic activation of RNA sensing (MDA5) or DNA (cGAS/STING) leads to an SLE-like condition in humans.

## SOURCES OF TYPE I IFN-TRIGGERING DNA AND RNA

Since type I IFNs are elevated in most SLE patients and appear to play an important role in SLE pathogenesis and since perturbations in nucleic sensing pathways lead to a disease (i.e., AGS) characterized by chronically elevated type I IFN and autoimmunity with many of the same autoantibodies as in SLE, it seems logical to ask if sensor-triggering nucleic acids might be present in SLE patients. Alternatively, the function or regulation of one or several sensors might be faulty. Although evidence exists for the association of genetic variants of DNA and RNA sensors with SLE, and mutations in them can cause type I interferonopathies (80), such mutations are present in a very small subset of SLE patients. Hence, it would be important to elucidate which nucleic acids are aberrantly present in SLE patient. What is their nature and origin?

Although viruses have long been suspected to play some role in triggering several different autoimmune diseases, there is little evidence for a persistent presence of viral RNA or DNA in SLE patients. If aberrant nucleic acids are present in SLE patients to trigger the DNA and/or RNA sensors discussed above, they likely are derived from endogenous sources, such as chromosomal DNA, mitochondrial DNA, DNA made by reverse-transcription from RNA templates, RNA transcripts from normally silent loci of ancient viral origin (that still somehow resemble viral RNA), mis-edited RNA, or otherwise altered or improperly processed RNA molecules (**Figure 1**). We will discuss these potential sources one by one.

### Chromosomal DNA

While chromosomal DNA normally is well protected by myriad binding proteins and a highly ordered packing into nucleosomes and higher order structures, DNA damage or faulty DNA replication can, in principle, dislodge smaller pieces of DNA, for example as nuclear blebs or micronuclei found in cancers (81). The existence of several effective DNA repair mechanisms indicate that DNA damage does occur in cells for a variety of reasons, including during normal aging. It is conceivable that DNA damage could produce pieces of DNA that trigger cGAS and subsequent type I IFN production (82, 83). Loss of DNA degradation by DNase1L3 causes an autosomal recessive form of SLE with early life onset and high prevalence of nephritis (84), and loss of the Trex1 DNase (85, 86) also leads to constitutive type I production and SLE or AGS, indicating that rapid elimination of aberrant DNA is important for the maintenance of health.

Cell death, whether by physiological programmed cell death mechanisms, such as apoptosis, or, more likely, by more pathological, inflammatory, or toxic mechanisms like necrosis, pyroptosis, or necroptosis, can result in the release

of chromosomal DNA (and RNA) into the extracellular milieu (87, 88). Many protective processes have evolved to minimize this exposure to chromatin and the highly toxic histones (89). Apoptotic cells are recognized by specific receptors for phosphatidylserine, annexin V, and other molecules that serve to mark apoptotic cells to facilitate their rapid removal by tissue macrophages (90) and the reticuloendothelial system. When these mechanisms are faulty or overwhelmed by massive numbers of dying cells, anti-nuclear and nucleic acid-directed autoantibodies and autoimmune disease may develop (87). For example, severe viral infections that are accompanied by immune-mediated killing of large numbers of infected cells typically result in measurable titers of anti-nucleic acid autoantibodies in otherwise healthy individuals. However, these titers tend to be relatively modest and they decline after the infections is cleared. The complement system (91), particularly C1q, also participates in the non-inflammatory removal of dying cells, perhaps explaining why complement deficiencies predispose to SLE (92). C1q also influences type I IFN production by a more direct mechanism (93, 94).

### Mitochondrial DNA

Another source of nucleic acids are the mitochondria (95, 96), which serve many functions besides oxidative phosphorylation and production of ATP, such as metabolism, inflammation and cell death. Though mainly found intracellularly, we recently discovered that neutrophils can extrude mitochondria together with chromosomal DNA during the formation of neutrophil extracellular traps (NETs) (95, 97). The externalization of mitochondria depended on the generation of reactive oxygen species (ROS) and the extruded mitochondria contained highly oxidized (8-OHdG) mitochondrial DNA, inducing IFN $\beta$  generation in a process requiring the intracellular DNA sensor adaptor protein, STING (95). Blocking mitochondrial ROS generation *in vivo* ameliorated lupus-like disease in MRL/lpr mice (95). These observations appear to be clinically relevant as *ex vivo* neutrophils from SLE patients displayed ongoing mitochondrial ROS production and spontaneous extrusion of oxidized inflammatory mitochondrial DNA (95). Similar to NET formation, as described above, other forms of cell death, such as TNF-mediated necroptosis, have been shown to involve the release of intact mitochondria into the extracellular environment (98, 99). Though the intracellular source(s) of extruded DNA has yet to be verified in other forms of cell death, e.g., pyroptosis, we find it likely that any form of cell death that includes breakdown of the plasma membrane will result in the release of mitochondria or their components, such as mitochondrial DNA.

Mitochondria and mitochondrial DNA may also be released from live cells upon their activation, as shown in neutrophils, eosinophils, mast cells and platelets (100–103). In neutrophils, this process has been coined “vital” NETosis, as the neutrophil remains alive after the extrusion event. Circulating platelets are thought to be the primary source of cell-free mitochondria given the large abundance of platelets in blood. Work from the laboratory of Eric Boilard has demonstrated that platelets, upon activation, may extrude naked mitochondria able to undergo respiratory burst (103). Unless these mitochondria

are rapidly cleared, secreted phospholipase A2 will hydrolyze and weaken the mitochondrial membrane, causing disruption and the release of inflammatory mitochondrial DNA and other damage-associated molecules (103). Platelet-mediated extrusion of mitochondria can occur in concentrated platelet preparations and is associated with adverse reactions upon transfusions (103, 104). Mitochondria can also be released as part of microparticles from many different cells, including platelets, neuronal and glial cells, as well as hepatocytes (103, 105, 106). The role of platelet-mediated release of mitochondrial DNA is of particular interest in rheumatic disease, including SLE, given the marked platelet activation and subsequent development of cardiovascular morbidity and mortality observed in these patients (107–110).

While the role of mitochondrial extrusion and mitochondrial DNA (oxidized or not) in SLE remains to be clarified, it is clear that they can be derived from many different cells and involve either the activation or death of these cells. It also appears that the DNA sensors TLR9 (105, 111–113) and cGAS (95, 114, 115) can be triggered by mitochondrial DNA, presumably depending on its subcellular location or pathway of receptor-mediated internalization (**Figure 1**). Elevated amounts of free mitochondrial DNA have been observed in several conditions, including chronic inflammatory diseases, trauma, cardiovascular disease and rheumatoid arthritis, perhaps promoting inflammation and even mortality (103, 116–118). Further studies to elucidate the mechanisms by which extruded mitochondria and/or mitochondrial DNA are cleared will be important for our understanding of this biology and for the design of therapeutic regimens to prevent the contribution of mitochondria and/or their DNA to human autoimmunity.

## Reverse-Transcribed RNA

The third source of DNA that may trigger type I IFN synthesis is intracellular DNA made by the reverse transcription of cellular RNA. Our genome encodes three different families of reverse transcriptases (RTs): telomerase (*TERT*), the *pol* genes of many endogenous retroviruses, and the second open-reading frame (ORF2) of the long interspersed nuclear element-1 (LINE1). Of these enzymes, telomerase is highly specialized to synthesize TTAGGG repeats in the 194 telomeres of our diploid chromosomes using the *TERC* RNA template (119, 120), while retroviral RTs only function to convert the RNA genome of an incoming retrovirus to a DNA provirus and to insert it into the genome. Although our genome contains thousands of endogenous retroviral provirus loci, none of them appear to be infectious anymore (with the possible exception of HERV-K113). This leaves only the LINE1 ORF2 enzyme as a candidate RT capable of generating aberrant DNA that could trigger type I IFN production through cGAS activation. It has been demonstrated to have robust RT activity (121–123), which is key for retrotransposition (124) and which is sensitive to some clinically used RT inhibitors (125, 126).

The LINE1 element represents a remnant of an ancient retrovirus that retained, or later acquired, a degree of autonomy through the conservation of a primordial RT, which endows it with the ability to transpose by a “copy and paste” mechanism. The LINE1 RNA transcript is 6 kb long and contains two

open-reading frames (79): ORF1, which encodes a 40-kDa RNA-binding protein that co-localizes with LINE1 mRNA in stress granules together with other RNA-binding proteins (127), such as Ro60, La, and U1 Small nuclear ribonucleoprotein of 70 kDa (127), and ORF2, which encodes a 150-kDa RT and endonuclease. Attesting to the effectiveness of the LINE1 element’s ability to transpose, there are over half a million copies of it throughout our genome. However, due to defense mechanisms (including those encoded by the AGS genes) and mutational rate, the vast majority of these copies are truncated and mutated and no longer have the ability to transpose. It has been estimated that < 180 LINE1 copies are seemingly intact, but that only 5 or 6 of them are active (“hot”) today (128). The LINE1 ORF2-encoded RT is also involved in generating and transposing Alu element copies (129) and was, over evolutionary time, responsible for generating all of our processed pseudogenes (130). In other words, the LINE1-encoded RT has had a profound impact on our genome and our health.

The study of AGS revealed that transcription of retroelement loci is very low in healthy individuals, but that AGS patients have elevated levels of retroelement mRNAs and proteins, including enzymatically active LINE1 RT (79). In fact, LINE1 RT may be the main producer of pathogenic DNA that triggers type I IFN production (131) in AGS patients with *TREX1*, *RNASEH2A*, *RNASEH2B*, *RNASEH2C*, and *SAMHD1* mutations. *TREX1* is the DNase that degrades intracellular DNA made by reverse transcription (86, 132), including DNA in complex with RNA as occurs during reverse transcription, while RNaseH2 preferentially acts on the RNA in such heteroduplexes (77). Finally, *SAMHD1* is a phosphohydrolase specific for the deoxynucleotide triphosphates (dATP, dTTP, dGTP, and dCTP) required for reverse transcription (133). In a mouse model of AGS, the *Trex1*<sup>-/-</sup> mouse (86), the animals develop a systemic inflammation with immune cell infiltrates in many organs and they die early from a severe carditis. These animals can be rescued from death by treatment with the RT inhibitors tenofovir plus nevirapine (134), indicating that reverse transcription is a key step in the pathogenesis of systemic inflammation in this model. However, there is also a published paper refuting these data (135). A human clinical trial with RT inhibitors in AGS is under way.

There is some evidence that LINE1 retroelements are activated also in SLE patients (136–138). This appears to correlate with a global decrease in DNA methylation, which is well documented in SLE (139, 140) and likely relates to the decreased expression of DNA methylases DNMT1 and DNMT3a (141, 142). Demethylating agents like 5-aza-2′-deoxycytidine (143) also cause a dramatic upregulation of LINE1 and Alu element transcription in lymphocytes (144). In addition, transfer of 5-aza-2′-deoxycytidine-treated T cells into healthy mice results in an SLE-like disease (145). The drugs that can induce “drug-induced lupus,” notably hydralazine and procainamide, are demethylating agents (146). Other known triggers of lupus flares, like UV light, oxidative stress, inflammation and exogenous viruses also induce genomic hypomethylation (147, 148).

We are aware of only two papers that report the detection of LINE1-encoded ORF1 and ORF2 proteins in samples from patients with SLE or related diseases. Mavgrani and co-workers

(136) showed by immunoblotting and immunohistochemistry that p40/ORF1 was readily detectable in kidney samples from lupus nephritis patients and in salivary gland biopsies of Sjögren's patients. Staining correlated with IFN $\beta$  in somatic cells and IFN $\alpha$  in infiltrating plasmacytoid dendritic cells. As activation of LINE1 elements in autoimmune patients (136–138) appears to involve demethylation of the LINE1 promoter (136, 138, 140), these authors also analyzed the methylation of CpG sites in the LINE1 promoter and found it to be reduced in patients with elevated LINE1 expression. In the second paper, Kalogirou et al. demonstrated that ORF2 is upregulated in the ductal cells of salivary gland biopsies from patients with Sjögren's syndrome (149).

### Cellular RNAs and RNA Editing by ADAR1

Many viruses have an RNA genome and do not (unlike the retroviruses) generate any DNA. A set of cellular RNA sensors have evolved to detect these viruses (150) (Figure 1B), a challenging task given the abundance of cellular RNA species. It remains incompletely understood how these sensors can distinguish between self and foreign RNA molecules, but the length of double-stranded RNAs (150) and capping modifications of the 5' and 3' ends of RNA molecules (151), as well as the presence or absence of other types of RNA processing, appear to matter. The delicate balance between the recognition of self- vs. foreign RNA is well illustrated by the *IFIH1*-A947T allele, which encodes a variant of MDA5 that enhances anti-viral immunity, but increases the risk of autoimmunity (152, 153).

Extracellular RNA, for example in immune complexes with the Ro protein (154), can also enter immune cells via receptor-mediated endocytosis followed by trafficking to the endosomal compartment where TLR3 will react to dsRNA and TLR7 and 8 with single-stranded RNA (as well as other ligands, see section TLRs in SLE). From this compartment, dsRNA may also exit into the cytosol through the SIDT2 channel (48) to trigger cytosolic MDA5. This pathway likely exists to aid in the detection of RNA viruses that are captured by antibodies, complement, or scavenger receptors.

The role of ADAR1 is also very interesting as this enzyme is involved in the post-transcriptional editing of mRNAs by converting adenosine to inosine, which is read as a guanosine during translation. Interestingly, in humans (unlike other organisms) the majority of the deaminated adenosines are non-coding and located in RNA molecules derived from Alu elements and other transposable sequences (155). The induction of type I IFN by mutated ADAR1 is dependent on MDA5, but not on RIG-I, suggesting that RNA editing is important and that its absence triggers the MDA5 pathway as if viral dsRNA was present.

### Endogenous Retroviral RNA

While exogenous viruses introduce RNA (or DNA) molecules that can be recognized as foreign by cellular sensors, it remains doubtful that RNA transcripts from endogenous proviruses (which constitute as much as 8% of our genome) would be seen as foreign as they are transcribed and processed by the normal cellular machinery. Nevertheless, since these sequences are of viral origin, it is possible that some of them still contain

sequence motifs that allow cellular RNA sensors to recognize them as non-self. If so, one would expect the relevant loci to be among the most recently incorporated ones, which may not yet have accumulated domesticating mutations. Furthermore, since most people do not develop autoimmunity, one would also assume that they normally are effectively silenced in healthy individuals, but perhaps aberrantly expressed in patients with SLE or related diseases.

### Transposable Element RNA

A much more interesting category of RNAs in autoimmunity research are those encoded by Alu elements and other short transposable elements, not perhaps because of their origin, but because they have been experimentally implicated in several settings. An important paper in this respect reported that a large portion of all RNA present in circulating anti-Ro autoantibody immunocomplexes was Alu element RNA (154). In fact, other SLE autoantigens, such as La (156), also bind Alu RNA. Furthermore, Ro<sup>-/-</sup> mice (157) develop autoimmunity resembling SLE, suggesting that the normal function of Ro is important for preventing the Alu element RNA, and perhaps other cellular RNA molecules (158, 159), from triggering RNA sensors. The discovery that the RNA-editing enzyme ADAR1 primarily edits Alu transcripts in humans (155) and that the LINE-1 encoded RT has catalyzed the reverse transcription and genomic insertion of over a million copies of the Alu element in our genome, as well as the co-localization of LINE1 proteins with Ro and La, hints at a central, but still enigmatic, role of this RNA biology in SLE pathogenesis. It also remains unknown how Ro-Alu RNA complexes end up in the extracellular compartment, but one can suspect that cell death by several programmed mechanisms must be involved.

### PATIENT HETEROGENEITY WITH REGARD TO NUCLEIC ACIDS AND THEIR SENSORS?

While it seems likely that type I IFNs play an important role in the pathogenesis of SLE and related diseases, it also becoming clear that their inhibition is not a cure for all patients. For example, 10–30% of SLE patients do not have an IFN signature, suggesting that their disease does not involve elevated type I IFNs and may therefore be molecularly altogether different. Furthermore, within the subpopulation of SLE patients with an IFN signature, therapeutic antibodies that neutralize IFN $\alpha$  (41, 160, 160, 161) or all type I IFNs (by blocking their receptor) (40) have been clinically efficacious in some patients, but not in others. The reasons for this heterogeneity are not understood, but may be related to the complexity of the interferon system, the coverage of different interferons by the therapeutics, the upstream drivers of type I IFN production, which depend not only on the cells that produce them but also the ligands that drive type I IFN. It should also be noted that even if pathogenic nucleic acid species induce much of their downstream effect through type I IFNs, there are also some IFN-independent consequences (e.g., through NF- $\kappa$ B

activation) that may contribute to SLE, but not be blocked by IFN antibodies.

At present, it is not known whether SLE patients with an elevated type I IFN gene signature always have the same nucleic acid sensor(s) activated or if each individual patient has a unique pattern that may include any or all of them. It is also unknown if the same pathogenic nucleic acid species are present in all patients, or if they too are different from patient to patient. Furthermore, we cannot entirely exclude the possibility that the nucleic acid sensors are sufficiently dysfunctional (for any number of reasons) to trigger type I IFN production even in the absence of any aberrant DNA or RNA. However, based on the heterogeneity of SLE and the heterogeneity in response to therapeutic IFN blocking antibodies, we find it most likely that there is also heterogeneity in the presence of pathogenic DNA and RNA species resulting in the activation of a different set of sensors in each patient.

Based on GWAS and other genetic data, it also seems that a great deal of patient heterogeneity is conferred by the presence of disease-predisposing or -protective alleles in many genes, most of which are immune-related. While pathogenic nucleic acids may be instrumental in initiating and perpetuating SLE, the overall sensitivity of the immune system, as determined by all these gene variants in immune-related genes (e.g., *MHC* and *PTPN22*), will affect how readily such nucleic acids tip the balance between transient responses vs. frank autoimmune disease. Interestingly, the SLE-predisposing variant of *PTPN22* not only affects T and B cell signaling, but also type I IFN production (162).

### CAN SLE BE DIVIDED INTO CLINICALLY MEANINGFUL SUBPOPULATIONS BASED ON “ENDOTYPE”?


The unpredictable response of patients with SLE to standard of care medication is a significant challenge in rheumatology. The current paradigm is to treat patients with escalating doses of increasingly potent immunosuppressive drugs until the

clinical response is deemed sufficient and then taper off the strongest immunosuppressants, particularly steroids. Even so, many patients never achieve complete remission but continue to suffer various degrees of symptoms that compromise their health and quality of life, not to mention the threat of sudden exacerbations.

In contrast, the treatment of rheumatoid arthritis made an important advance with the introduction of etanercept in 1999 (163–165). However, this drug also gave rise to the concept of “TNF-non-responders.” While most patients at least initially respond clinically to etanercept and other TNF blockers, 20–30% respond poorly, if at all. Other biologics have typically met with similar outcomes: good efficacy in many patients, but always a number of non-responders or initial responders who lose efficacy over time. It seems that non-responders represent individuals whose disease differs molecularly from the responders, such that the disease process does not involve, or readily circumvents, the specific target for the drug and therefore continues unabated.

We believe that this responder/non-responder dichotomy is also relevant in SLE and related diseases, where new drugs in recent clinical trials have generally yielded poor efficacy or a minority of (partial) responders and a majority of non-responders. As SLE is clinically highly variable, it is easy to believe that it is molecularly heterogenous as well. We propose here that SLE patients could be grouped into molecularly distinct categories (“endotypes”) based on which nucleic acid sensors are active and the IFN species produced in response to them (Figure 2):

- 1) IFN-independent SLE, represented by the 10–30% of patients who do not have a type I IFN gene signature and, therefore, unlikely any nucleic acid sensor activation.
- 2) SLE patients whose elevated and disease-driving type I IFNs are restricted to isoforms of IFN $\alpha$ , which are predominantly made by immune cells via TLR7/9 in response to circulating immune complexes that contain nucleic acids.
- 3) SLE patients with predominantly IFN $\beta$ , which is typically made by epithelial and other cells via activation of cGAS,



**FIGURE 2 |** Proposed endotypes of SLE based on type I IFN subtype and relevant nucleic acid sensors. The first column represents the four proposed endotypes of SLE with double arrows connecting them to the relevant nucleic acid sensors. The predicted (or known) effects on each SLE endotype of an antibody that neutralizes IFN $\alpha$  only (sifalimumab), an antibody that blocks all type I IFNs (anifrolumab), and hypothetical cGAS or RT inhibitors are indicated as “yes” for a substantial clinical benefit, “no” for none, and “partial” if only one of two parallel mechanisms are expected to be inhibited.



RIG-I, or MDA5. These patients may be at an early stage of SLE development (including preclinical disease), and have not yet developed circulating immune complexes with nucleic acids. Alternatively, their disease will never develop such immune complexes.

- 4) SLE patients who have numerous IFN $\alpha$ s and IFN $\beta$  (and perhaps IFN $\epsilon$ , IFN $\kappa$ , or IFN $\omega$ ) and both TLR7/9 and intracellular nucleic acid sensor pathways active. If the third category includes early disease, this fourth endotype may contain severe and late stage disease.

What we propose is the personalized medicine notion that patients suffering from a disease like SLE can be subdivided by patient endotype into subsets that share a specific molecular mechanism that originally initiated and continues to perpetuate their disease, and that the inhibition of this mechanism by a selective therapeutic approach will provide a strong clinical benefit specifically to patients within this subset, but perhaps not to others. Practical examples of this concept exist in medical practice today in oncology and respiratory medicine (166) but are still absent in rheumatology. Key to the utility of this concept is the development of practical tests (“biomarkers”) that can determine which endotype individual patients belong to. In this particular case, such biomarkers would naturally quantitate nucleic acids, the activation of the sensors, and/or assess the spectrum of IFNs in patient blood or tissue.

To fully test our SLE endotype concept, future trials with new and more targeted therapeutics for SLE should include the

relevant biomarkers to ask if therapeutic efficacy falls within one or another SLE endotype. In other examples of the endotype concept (166), this type of patient stratification approach has resulted in astonishing levels of efficacy within the relevant endotype, but marginal impact on patients of other endotypes. Oftentimes, these same clinical trials failed to meet their primary endpoint when the all-comers population was assessed.

## AUTHOR CONTRIBUTIONS

TM, CL, and NG contributed equally to the writing of this review and share accountability for its content.

## FUNDING

Our work is supported by NIH grant R21AR075134 (TM), Lupus Research Alliance grant 519414 (CL), Lupus Research Alliance grant 519408 (NG), and UL1 TR002319 (ITHS Catalyst Award) from the National Center for Advancing Translational Sciences, NIH (NG).

## ACKNOWLEDGMENTS

We apologize to all those colleagues whose many important papers we could not cite due to space constraints. We are grateful to Professor Keith Elkon for providing valuable feedback and suggestions.

## REFERENCES

- Gensous N, Marti A, Barnetche T, Blanco P, Lazaro E, Seneschal J, et al. Predictive biological markers of systemic lupus erythematosus flares: a systematic literature review. *Arthritis Res Ther.* (2017) 19:238. doi: 10.1186/s13075-017-1442-6
- Kramer N, Krey PR. SLE: use of laboratory tests in diagnosis and management. *J Med Soc N J.* (1983) 80:344–9.
- Touma Z, Gladman DD. Current and future therapies for SLE: obstacles and recommendations for the development of novel treatments. *Lupus Sci Med.* (2017) 4:e000239. doi: 10.1136/lupus-2017-000239
- Ponticelli C, Moroni G. Hydroxychloroquine in systemic lupus erythematosus (SLE). *Expert Opin Drug Saf.* (2017) 16:411–9. doi: 10.1080/14740338.2017.1269168
- Rainsford KD, Parke AL, Clifford-Rashotte M, Kean WF. Therapy and pharmacological properties of hydroxychloroquine and chloroquine in treatment of systemic lupus erythematosus, rheumatoid arthritis and related diseases. *Inflammopharmacology.* (2015) 23:231–69. doi: 10.1007/s10787-015-0239-y
- Furie R, Petri M, Zamani O, Cervera R, Wallace DJ, Tegzova D, et al. A phase III, randomized, placebo-controlled study of belimumab, a monoclonal antibody that inhibits B lymphocyte stimulator, in patients with systemic lupus erythematosus. *Arthritis Rheum.* (2011) 63:3918–30. doi: 10.1002/art.30613
- Boyce EG, Fusco BE. Belimumab: review of use in systemic lupus erythematosus. *Clin Ther.* (2012) 34:1006–22. doi: 10.1016/j.clinthera.2012.02.028
- Ramos-Casals M, Sanz I, Bosch X, Stone JH, Khamashta MA. B-cell-depleting therapy in systemic lupus erythematosus. *Am J Med.* (2012) 125:327–36. doi: 10.1016/j.amjmed.2011.09.010
- Mok CC. Calcineurin inhibitors in systemic lupus erythematosus. *Best Pract Res Clin Rheumatol.* (2017) 31:429–38. doi: 10.1016/j.berh.2017.09.010
- Hoi A, Littlejohn G. Is there still a role for abatacept in the treatment of lupus? *Expert Opin Biol Ther.* (2014) 14:1345–50. doi: 10.1517/14712598.2014.935329
- Tan EM, Cohen AS, Fries JP, Masi AT, McShane DJ, Rothfield NF, et al. The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum.* (1982) 25:1271–7. doi: 10.1002/art.1780251101
- Bombardier C, Gladman D, Urowitz M, Caron D, Chang C, SLE tCoPSI. Derivation of the SLEDAI. a disease activity index for lupus patients. *Arthritis Rheum.* (1992) 35:630–40. doi: 10.1002/art.1780350606
- Baechler EC, Batliwalla FM, Karypis G, Gaffney PM, Ortmann WA, Espe KJ, et al. Interferon-inducible gene expression signature in peripheral blood cells of patients with severe lupus. *Proc Natl Acad Sci USA.* (2003) 100:2610–5. doi: 10.1073/pnas.0337679100
- Bennett L, Palucka AK, Arce E, Cantrell V, Borvak J, Banchereau J, et al. Interferon and granulopoiesis signatures in systemic lupus erythematosus blood. *J Exp Med.* (2003) 197:711–23. doi: 10.1084/jem.20021553
- Crow MK, Kirou KA, Wohlgemuth J. Microarray analysis of interferon-regulated genes in SLE. *Autoimmunity.* (2003) 36:481–90. doi: 10.1080/08916930310001625952
- Ronnblom L, Alm GV. A pivotal role for the natural interferon alpha-producing cells (plasmacytoid dendritic cells) in the pathogenesis of lupus. *J Exp Med.* (2001) 194:F59–63. doi: 10.1084/jem.194.12.f59
- Blomberg S, Eloranta ML, Cederblad B, Nordlin K, Alm GV, Ronnblom L. Presence of cutaneous interferon-alpha producing cells in patients with systemic lupus erythematosus. *Lupus.* (2001) 10:484–90. doi: 10.1191/096120301678416042
- Ronnblom LE, Alm GV, Oberg KE. Possible induction of systemic lupus erythematosus by interferon-alpha treatment in a

- patient with a malignant carcinoid tumour. *J Intern Med.* (1990) 227:207–10. doi: 10.1111/j.1365-2796.1990.tb00144.x
19. Ronnblom L, Eloranta ML. The interferon signature in autoimmune diseases. *Curr Opin Rheumatol.* (2013) 25:248–53. doi: 10.1097/BOR.0b013e32835c7e32
  20. Ronnblom L, Eloranta ML, Alm GV. The type I interferon system in systemic lupus erythematosus. *Arthritis Rheum.* (2006) 54:408–20. doi: 10.1002/art.21571
  21. Eloranta ML, Ronnblom L. Cause and consequences of the activated type I interferon system in SLE. *J Mol Med (Berl).* (2016) 94:1103–10. doi: 10.1007/s00109-016-1421-4
  22. Chiche L, Jourde-Chiche N, Whalen E, Presnell S, Gersuk V, Dang K, et al. Modular transcriptional repertoire analyses of adults with systemic lupus erythematosus reveal distinct type I and type II interferon signatures. *Arthritis Rheumatol.* (2014) 66:1583–95. doi: 10.1002/art.38628
  23. Rodero MP, Decalf J, Bondet V, Hunt D, Rice GI, Werneke S, et al. Detection of interferon alpha protein reveals differential levels and cellular sources in disease. *J Exp Med.* (2017) 214:1547–55. doi: 10.1084/jem.20161451
  24. Bodewes ILA, Al-Ali S, van Helden-Meeuwesen CG, Maria NI, Tarn J, Lendrem DW, et al. Systemic interferon type I and type II signatures in primary Sjogren's syndrome reveal differences in biological disease activity. *Rheumatology (Oxford).* (2018) 57:921–30. doi: 10.1093/rheumatology/kex490
  25. Hall JC, Baer AN, Shah AA, Criswell LA, Shiboski CH, Rosen A, et al. Molecular subsetting of interferon pathways in sjogren's syndrome. *Arthritis Rheumatol.* (2015) 67:2437–46. doi: 10.1002/art.39204
  26. Eloranta ML, Franck-Larsson K, Lovgren T, Kalamajski S, Ronnblom A, Rubin K, et al. Type I interferon system activation and association with disease manifestations in systemic sclerosis. *Ann Rheum Dis.* (2010) 69:1396–402. doi: 10.1136/ard.2009.121400
  27. Wuttge DM, Lood C, Tufvesson E, Scheja A, Truedsson L, Bengtsson AA, et al. Increased serum type I interferon activity in early systemic sclerosis patients is associated with antibodies against Sjogren's syndrome antigens and nuclear ribonucleoprotein antigens. *Scand J Rheumatol.* (2013) 42:235–40. doi: 10.3109/03009742.2012.736532
  28. Baechler EC, Bauer JW, Slattery CA, Ortmann WA, Espe KJ, Novitzke J, et al. An interferon signature in the peripheral blood of dermatomyositis patients is associated with disease activity. *Mol Med.* (2007) 13:59–68. doi: 10.2119/2006-00085.Baechler
  29. Niewold TB, Kariuki SN, Morgan GA, Shrestha S, Pachman LM. Elevated serum interferon-alpha activity in juvenile dermatomyositis: associations with disease activity at diagnosis and after thirty-six months of therapy. *Arthritis Rheum.* (2009) 60:1815–24. doi: 10.1002/art.24555
  30. Rodriguez-Carrio J, Alperi-Lopez M, Lopez P, Ballina-Garcia FJ, Suarez A. Heterogeneity of the type I interferon signature in rheumatoid arthritis: a potential limitation for its use as a clinical biomarker. *Front Immunol.* (2017) 8:2007. doi: 10.3389/fimmu.2017.02007
  31. El-Sherbiny YM, Psarras A, Md Yusof MY, Hensor EMA, Tooze R, Doody G, et al. A novel two-score system for interferon status segregates autoimmune diseases and correlates with clinical features. *Sci Rep.* (2018) 8:5793. doi: 10.1038/s41598-018-24198-1
  32. Zahn S, Rehkemper C, Kummerer BM, Ferring-Schmidt S, Bieber T, Tuting T, et al. Evidence for a pathophysiological role of keratinocyte-derived type III interferon (IFNlambda) in cutaneous lupus erythematosus. *J Invest Dermatol.* (2011) 131:133–40. doi: 10.1038/jid.2010.244
  33. Buechler MB, Teal TH, Elkon KB, Hamerman JA. Cutting edge: type I IFN drives emergency myelopoiesis and peripheral myeloid expansion during chronic TLR7 signaling. *J Immunol.* (2013) 190:886–91. doi: 10.4049/jimmunol.1202739
  34. Lee PY, Li Y, Kumagai Y, Xu Y, Weinstein JS, Kellner ES, et al. Type I interferon modulates monocyte recruitment and maturation in chronic inflammation. *Am J Pathol.* (2009) 175:2023–33. doi: 10.2353/ajpath.2009.090328
  35. Blanco P, Palucka AK, Gill M, Pascual V, Banchereau J. Induction of dendritic cell differentiation by IFN-alpha in systemic lupus erythematosus. *Science.* (2001) 294:1540–3. doi: 10.1126/science.1064890
  36. Kolumam GA, Thomas S, Thompson LJ, Sprent J, Murali-Krishna K. Type I interferons act directly on CD8 T cells to allow clonal expansion and memory formation in response to viral infection. *J Exp Med.* (2005) 202:637–50. doi: 10.1084/jem.20050821
  37. Joo H, Coquery C, Xue Y, Gayet I, Dillon SR, Punaro M, et al. Serum from patients with SLE instructs monocytes to promote IgG and IgA plasmablast differentiation. *J Exp Med.* (2012) 209:1335–48. doi: 10.1084/jem.20111644
  38. Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, et al. A diverse range of gene products are effectors of the type I interferon antiviral response. *Nature.* (2011) 472:481–5. doi: 10.1038/nature09907
  39. Ronnblom L. The importance of the type I interferon system in autoimmunity. *Clin Exp Rheumatol.* (2016) 34(4 Suppl 98):21–4.
  40. Furie R, Khamashta M, Merrill JT, Werth VP, Kalunian K, Brohawn P, et al. Anifrolumab, an anti-interferon-alpha receptor monoclonal antibody, in moderate-to-severe systemic lupus erythematosus. *Arthritis Rheumatol.* (2017) 69:376–86. doi: 10.1002/art.39962
  41. Petri M, Wallace DJ, Spindler A, Chindalore V, Kalunian K, Mysler E, et al. Sifalimumab, a human anti-interferon-alpha monoclonal antibody, in systemic lupus erythematosus: a phase I randomized, controlled, dose-escalation study. *Arthritis Rheum.* (2013) 65:1011–21. doi: 10.1002/art.37824
  42. Gao D, Li T, Li XD, Chen X, Li QZ, Wight-Carter M, et al. Activation of cyclic GMP-AMP synthase by self-DNA causes autoimmune diseases. *Proc Natl Acad Sci USA.* (2015) 112(42):E5699–705. doi: 10.1073/pnas.1516465112
  43. Yoneyama M, Kikuchi M, Matsumoto K, Imaizumi T, Miyagishi M, Taira K, et al. Shared and unique functions of the DEXD/H-box helicases RIG-I, MDA5, and LGP2 in antiviral innate immunity. *J Immunol.* (2005) 175:2851–8. doi: 10.4049/jimmunol.175.5.2851
  44. Loo YM, Gale M, Jr. Immune signaling by RIG-I-like receptors. *Immunity.* (2011) 34:680–92. doi: 10.1016/j.immuni.2011.05.003
  45. Keating SE, Baran M, Bowie AG. Cytosolic DNA sensors regulating type I interferon induction. *Trends Immunol.* (2011) 32:574–81. doi: 10.1016/j.it.2011.08.004
  46. Radetsky R, Daher A, Gatignol A. ADAR1 and PKR, interferon stimulated genes with clashing effects on HIV-1 replication. *Cytokine Growth Factor Rev.* (2018) 40:48–58. doi: 10.1016/j.cytogfr.2018.03.007
  47. Wang TT, Li ZG, Wang Q. The RNA-specific adenosine deaminase ADAR1 inhibits human protein kinase R activation. *Viral Immunol.* (2018) 31:537–8. doi: 10.1089/vim.2018.0056
  48. Nguyen TA, Smith BRC, Tate MD, Belz GT, Barrios MH, Elgass KD, et al. SIDT2 Transports extracellular dsRNA into the cytoplasm for innate immune recognition. *Immunity.* (2017) 47:498–509. doi: 10.1016/j.immuni.2017.08.007
  49. Vance RE. Cytosolic DNA sensing: the field narrows. *Immunity.* (2016) 45:227–8. doi: 10.1016/j.immuni.2016.08.006
  50. Bao M, Liu YJ. Regulation of TLR7/9 signaling in plasmacytoid dendritic cells. *Protein Cell.* (2013) 4:40–52. doi: 10.1007/s13238-012-2104-8
  51. Cao W, Bover L, Cho M, Wen X, Hanabuchi S, Bao M, et al. Regulation of TLR7/9 responses in plasmacytoid dendritic cells by BST2 and ILT7 receptor interaction. *J Exp Med.* (2009) 206:1603–14. doi: 10.1084/jem.20090547
  52. Henault J, Riggs JM, Karnell JL, Liarski VM, Li J, Shirinian L, et al. Self-reactive IgE exacerbates interferon responses associated with autoimmunity. *Nat Immunol.* (2016) 17:196–203. doi: 10.1038/ni.3326
  53. Yu Z, Chen T, Cao X. Neutrophil sensing of cytoplasmic, pathogenic DNA in a cGAS-STING-independent manner. *Cell Mol Immunol.* (2016) 13:411–4. doi: 10.1038/cmi2015.34
  54. Kawai T, Akira S. TLR signaling. *Semin Immunol.* (2007) 19:24–32. doi: 10.1016/j.smim.2006.12.004
  55. Barrat FJ, Meeker T, Gregorio J, Chan JH, Uematsu S, Akira S, et al. Nucleic acids of mammalian origin can act as endogenous ligands for Toll-like receptors and may promote systemic lupus erythematosus. *J Exp Med.* (2005) 202:1131–9. doi: 10.1084/jem.20050914
  56. Elshabrawy HA, Essani AE, Szekanez Z, Fox DA, Shahrara S. TLRs, future potential therapeutic targets for RA. *Autoimmun Rev.* (2017) 16:103–13. doi: 10.1016/j.autrev.2016.12.003
  57. Giltiay NV, Chappell CP, Sun X, Kolhatkar N, Teal TH, Wiedeman AE, et al. Overexpression of TLR7 promotes cell-intrinsic expansion and autoantibody production by transitional T1 B cells. *J Exp Med.* (2013). 210:2773–89. doi: 10.1084/jem.20122798
  58. Shen N, Fu Q, Deng Y, Qian X, Zhao J, Kaufman KM, et al. Sex-specific association of X-linked toll-like receptor 7 (TLR7) with male

- systemic lupus erythematosus. *Proc Natl Acad Sci USA*. (2010) 107:15838–43. doi: 10.1073/pnas.1001337107
59. Lloyd W, Schur PH. Immune complexes, complement, and anti-DNA in exacerbations of systemic lupus erythematosus. *Medicine*. (1981) 60:208–17. doi: 10.1097/00005792-198105000-00004
  60. Lovgren T, Eloranta ML, Bave U, Alm GV, Ronnblom L. Induction of interferon-alpha production in plasmacytoid dendritic cells by immune complexes containing nucleic acid released by necrotic or late apoptotic cells and lupus IgG. *Arthritis Rheum*. (2004) 50:1861–72. doi: 10.1002/art.20254
  61. Wang D, Bhagat L, Yu D, Zhu FG, Tang JX, Kandimalla ER, et al. Oligodeoxyribonucleotide-based antagonists for toll-like receptors 7 and 9. *J Med Chem*. (2009) 52:551–8. doi: 10.1021/jm8014316
  62. Miyake K, Kaisho T. Homeostatic inflammation in innate immunity. *Curr Opin Immunol*. (2014) 30:85–90. doi: 10.1016/j.coi.2014.08.003
  63. Ohto U, Ishida H, Shibata T, Sato R, Miyake K, Shimizu T. Toll-like receptor 9 contains two DNA binding sites that function cooperatively to promote receptor dimerization and activation. *Immunity*. (2018) 48:649–58. doi: 10.1016/j.immuni.2018.03.013
  64. Ohto U, Shibata T, Tanji H, Ishida H, Krayukhina E, Uchiyama S, et al. Structural basis of CpG and inhibitory DNA recognition by toll-like receptor 9. *Nature*. (2015) 520:702–5. doi: 10.1038/nature14138
  65. Tanji H, Ohto U, Shibata T, Miyake K, Shimizu T. Structural reorganization of the toll-like receptor 8 dimer induced by agonistic ligands. *Science*. (2013) 339:1426–9. doi: 10.1126/science.1229159
  66. Tanji H, Ohto U, Shibata T, Taoka M, Yamauchi Y, Isobe T, et al. Toll-like receptor 8 senses degradation products of single-stranded RNA. *Nat Struct Mol Biol*. (2015) 22:109–15. doi: 10.1038/nsmb.2943
  67. Souyris M, Mejia JE, Chaumeil J, Guery JC. Female predisposition to TLR7-driven autoimmunity: gene dosage and the escape from X chromosome inactivation. *Semin Immunopathol*. (2018) 41:153–64. doi: 10.1007/s00281-018-0712-y
  68. Kelley J, Johnson MR, Alarcon GS, Kimberly RP, Edberg JC. Variation in the relative copy number of the TLR7 gene in patients with systemic lupus erythematosus and healthy control subjects. *Arthritis Rheum*. (2007) 56:3375–8. doi: 10.1002/art.22916
  69. Garcia-Ortiz H, Velazquez-Cruz R, Espinosa-Rosales F, Jimenez-Morales S, Baca V, Orozco L. Association of TLR7 copy number variation with susceptibility to childhood-onset systemic lupus erythematosus in Mexican population. *Ann Rheum Dis*. (2010) 69:1861–5. doi: 10.1136/ard.2009.124313
  70. Tian J, Ma Y, Li J, Cen H, Wang DG, Feng CC, et al. The TLR7 7926A>G polymorphism is associated with susceptibility to systemic lupus erythematosus. *Mol Med Rep*. (2012) 6:105–10. doi: 10.3892/mmr.2012.865
  71. An J, Durcan L, Woodward JJ, Karr R, Teale T, Elkon KB. cGAMP and cGAS are expressed in a subset of patients with systemic lupus erythematosus and associate with disease activity. *Arthritis Rheum*. (2017) 69:800–7. doi: 10.1136/annrheumdis-2016-eular.3752
  72. Shao WH, Shu DH, Zhen Y, Hilliard B, Priest SO, Cesaroni M, et al. Prion-like aggregation of mitochondrial antiviral signaling protein in lupus patients is associated with increased levels of type I interferon. *Arthritis Rheumatol*. (2016) 68:2697–707. doi: 10.1002/art.39733
  73. Aicardi J, Goutieres F. A progressive familial encephalopathy in infancy with calcifications of the basal ganglia and chronic cerebrospinal fluid lymphocytosis. *Ann Neurol*. (1984) 15:49–54. doi: 10.1002/ana.410150109
  74. Crow YJ, Chase DS, Lowenstein Schmidt J, Szykiewicz M, Forte GM, Gornall HL, et al. Characterization of human disease phenotypes associated with mutations in TREX1, RNASEH2A, RNASEH2B, RNASEH2C, SAMHD1, ADAR, and IFIH1. *Am J Med Genet A*. (2015) 167A:296–312. doi: 10.1055/s-0036-1592307
  75. Crow YJ, Vanderver A, Orcesi S, Kuijpers TW, Rice GI. Therapies in aicardi-goutieres syndrome. *Clin Exp Immunol*. (2014) 175:1–8. doi: 10.1111/cei.12115
  76. Ramantani G, Kohlhase J, Hertzberg C, Innes AM, Engel K, Hunger S, et al. Expanding the phenotypic spectrum of lupus erythematosus in Aicardi-Goutieres syndrome. *Arthritis Rheum*. (2010) 62:1469–77. doi: 10.1002/art.27367
  77. Pokatayev V, Hasin N, Chon H, Cerritelli SM, Sakhuja K, Ward JM, et al. RNase H2 catalytic core Aicardi-Goutieres syndrome-related mutant invokes cGAS-STING innate immune-sensing pathway in mice. *J Exp Med*. (2016) 213:329–36. doi: 10.1084/jem.20151464
  78. Crow YJ, Manel N. Aicardi-Goutieres syndrome and the type I interferonopathies. *Nat Rev Immunol*. (2015) 15:429–40. doi: 10.1038/nri3850
  79. Volkman HE, Stetson DB. The enemy within: endogenous retroelements and autoimmune disease. *Nat Immunol*. (2014) 15:415–22. doi: 10.1038/ni.2872
  80. Melki I, Rose Y, Ugenti C, Van Eyck L, Fremont ML, Kitabayashi N, et al. Disease-associated mutations identify a novel region in human STING necessary for the control of type I interferon signaling. *J Allergy Clin Immunol*. (2017) 140:543–552.e5. doi: 10.1016/j.jaci.2016.10.031
  81. Gekara NO. DNA damage-induced immune response: micronuclei provide key platform. *J Cell Biol*. (2017) 216:2999–3001. doi: 10.1083/jcb.201708069
  82. Ullal AJ, Reich CF III, Clowse M, Criscione-Schreiber LG, Tochacek M, Monestier M, et al. Microparticles as antigenic targets of antibodies to DNA and nucleosomes in systemic lupus erythematosus. *J Autoimmun*. (2011) 36:173–80. doi: 10.1016/j.jaut.2011.02.001
  83. Lande R, Ganguly D, Facchinetti V, Frasca L, Conrad C, Gregorio J, et al. Neutrophils activate plasmacytoid dendritic cells by releasing self-DNA-peptide complexes in systemic lupus erythematosus. *Sci Transl Med*. (2011) 3:73ra19. doi: 10.1126/scitranslmed.3001180
  84. Al-Mayouf SM, Sunker A, Abdwani R, Abrawi SA, Almurshedi F, Alhashmi N, et al. Loss-of-function variant in DNASE1L3 causes a familial form of systemic lupus erythematosus. *Nat Genet*. (2011) 43:1186–8. doi: 10.1038/ng.975
  85. Namjou B, Kothari PH, Kelly JA, Glenn SB, Ojwang JO, Adler A, et al. Evaluation of the TREX1 gene in a large multi-ancestral lupus cohort. *Genes Immun*. (2011) 12:270–9. doi: 10.1038/gene.2010.73
  86. Stetson DB, Ko JS, Heidmann T, Medzhitov R. Trex1 prevents cell-intrinsic initiation of autoimmunity. *Cell*. (2008) 134:587–98. doi: 10.1016/j.cell.2008.06.032
  87. Elkon KB. Review: cell death, nucleic acids, and immunity: inflammation beyond the grave. *Arthritis Rheum*. (2018) 70:805–16. doi: 10.1002/art.40452
  88. Pisetsky DS. DNA as a marker of cell death in systemic lupus erythematosus. *Rheum Dis Clin North Am*. (2004) 30:575–87. doi: 10.1016/j.rdc.2004.04.009
  89. Chen R, Kang R, Fan XG, Tang D. Release and activity of histone in diseases. *Cell Death Dis*. (2014) 5:e1370. doi: 10.1038/cddis.2014.337
  90. Fadok VA, Bratton DL, Konowal A, Freed PW, Westcott JY, Henson PM. Macrophages that have ingested apoptotic cells *in vitro* inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-beta, PGE2, and PAF. *J Clin Invest*. (1998) 101:890–8. doi: 10.1172/JCI1112
  91. Elkon KB, Santer DM. Complement, interferon and lupus. *Curr Opin Immunol*. (2012) 24:665–70. doi: 10.1016/j.coi.2012.08.004
  92. Bowness P, Davies KA, Norsworthy PJ, Athanassiou P, Taylor-Weideman J, Borysiewicz LK, et al. Hereditary C1q deficiency and systemic lupus erythematosus. *Quart J Med*. (1994) 87:455–64.
  93. Lood C, Gullstrand B, Truedsson L, Olin AI, Alm GV, Ronnblom L, et al. C1q inhibits immune complex-induced interferon-alpha production in plasmacytoid dendritic cells: a novel link between C1q deficiency and systemic lupus erythematosus pathogenesis. *Arthritis Rheum*. (2009) 60:3081–90. doi: 10.1002/art.24852
  94. Santer DM, Hall BE, George TC, Tangsombatvisit S, Liu CL, Arkwright PD, et al. C1q deficiency leads to the defective suppression of IFN-alpha in response to nucleoprotein containing immune complexes. *J Immunol*. (2010) 185:4738–49. doi: 10.4049/jimmunol.1001731
  95. Lood C, Blanco LP, Purmalek MM, Carmona-Rivera C, De Ravin SS, Smith CK, et al. Neutrophil extracellular traps enriched in oxidized mitochondrial DNA are interferogenic and contribute to lupus-like disease. *Nat Med*. (2016) 22:146–53. doi: 10.1038/nm.4027
  96. Caielli S, Athale S, Domic B, Murat E, Chandra M, Banchereau R, et al. Oxidized mitochondrial nucleoids released by neutrophils drive type I interferon production in human lupus. *J Exp Med*. (2016) 213:697–713. doi: 10.1084/jem.20151876
  97. Lood C, Hughes GC. Neutrophil extracellular traps as a potential source of autoantigen in cocaine-associated autoimmunity. *Rheumatology (Oxford)*. (2017) 56:638–43. doi: 10.1093/rheumatology/kew256

98. Maeda A, Fadeel B. Mitochondria released by cells undergoing TNF-alpha-induced necroptosis act as danger signals. *Cell Death Dis.* (2014) 5:e1312. doi: 10.1038/cddis.2014.277
99. Nakajima A, Kurihara H, Yagita H, Okumura K, Nakano H. Mitochondrial extrusion through the cytoplasmic vacuoles during cell death. *J Biol Chem.* (2008) 283:24128–35. doi: 10.1074/jbc.M802996200
100. Yousefi S, Gold JA, Andina N, Lee JJ, Kelly AM, Kozlowski E, et al. Catapult-like release of mitochondrial DNA by eosinophils contributes to antibacterial defense. *Nat Med.* (2008) 14:949–53. doi: 10.1038/nm.1855
101. Yousefi S, Mihalache C, Kozlowski E, Schmid I, Simon HU. Viable neutrophils release mitochondrial DNA to form neutrophil extracellular traps. *Cell Death Differ.* (2009) 16:1438–44. doi: 10.1038/cdd.2009.96
102. Zhang B, Asadi S, Weng Z, Sismanopoulos N, Theoharides TC. Stimulated human mast cells secrete mitochondrial components that have autocrine and paracrine inflammatory actions. *PLoS ONE.* (2012) 7:e49767. doi: 10.1371/journal.pone.0049767
103. Boudreau LH, Duchez AC, Cloutier N, Soulet D, Martin N, Bollinger J, et al. Platelets release mitochondria serving as substrate for bactericidal group IIA-secreted phospholipase A2 to promote inflammation. *Blood.* (2014) 124:2173–83. doi: 10.1182/blood-2014-05-573543
104. Marcoux G, Duchez AC, Rousseau M, Levesque T, Boudreau LH, Thibault L, et al. Microparticle and mitochondrial release during extended storage of different types of platelet concentrates. *Platelets.* (2017) 28:272–80. doi: 10.1080/09537104.2016.1218455
105. Garcia-Martinez I, Santoro N, Chen Y, Hoque R, Ouyang X, Caprio S, et al. Hepatocyte mitochondrial DNA drives nonalcoholic steatohepatitis by activation of TLR9. *J Clin Invest.* (2016) 126:859–64. doi: 10.1172/JCI83885
106. Zhao Z, Wang M, Tian Y, Hilton T, Salsbery B, Zhou EZ, et al. Cardiolipin-mediated procoagulant activity of mitochondria contributes to traumatic brain injury-associated coagulopathy in mice. *Blood.* (2016) 127:2763–72. doi: 10.1182/blood-2015-12-688838
107. Linde P, Fortin PR, Lood C, Bengtsson AA, Boilard E. The non-haemostatic role of platelets in systemic lupus erythematosus. *Nat Rev Rheumatol.* (2018) 14:195–213. doi: 10.1038/nrrheum.2018.38
108. Lood C, Amisten S, Gullstrand B, Jonsen A, Allhorn M, Truedsson L, et al. Platelet transcriptional profile and protein expression in patients with systemic lupus erythematosus: up-regulation of the type I interferon system is strongly associated with vascular disease. *Blood.* (2010) 116:1951–7. doi: 10.1182/blood-2010-03-274605
109. Lood C, Tyden H, Gullstrand B, Jonsen A, Kallberg E, Morgelin M, et al. Platelet-derived S100A8/A9 and cardiovascular disease in systemic lupus erythematosus. *Arthritis Rheumatol.* (2016) 68:1970–80. doi: 10.1002/art.39656
110. Manzi S, Meilahn EN, Rairie JE, Conte CG, Medsger TA, Jr, Jansen-McWilliams L, et al. Age-specific incidence rates of myocardial infarction and angina in women with systemic lupus erythematosus: comparison with the Framingham study. *Am J Epidemiol.* (1997) 145:408–15. doi: 10.1093/oxfordjournals.aje.a009122
111. Zhang Q, Raoof M, Chen Y, Sumi Y, Sursal T, Junger W, et al. Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature.* (2010) 464:104–7. doi: 10.1038/nature08780
112. Oka T, Hikoso S, Yamaguchi O, Taneike M, Takeda T, Tamai T, et al. Mitochondrial DNA that escapes from autophagy causes inflammation and heart failure. *Nature.* (2012) 485:251–5. doi: 10.1038/nature10992
113. Bliksoen M, Mariero LH, Torp MK, Baysa A, Ytrehus K, Haugen F, et al. Extracellular mtDNA activates NF-kappaB via toll-like receptor 9 and induces cell death in cardiomyocytes. *Basic Res Cardiol.* (2016) 111:42. doi: 10.1007/s00395-016-0553-6
114. West AP, Khoury-Hanold W, Staron M, Tal MC, Pineda CM, Lang SM, et al. Mitochondrial DNA stress primes the antiviral innate immune response. *Nature.* (2015) 520:553–7. doi: 10.1038/nature14156
115. Rongvaux A, Jackson R, Harman CC, Li T, West AP, de Zoete MR, et al. Apoptotic caspases prevent the induction of type I interferons by mitochondrial DNA. *Cell.* (2014) 159:1563–77. doi: 10.1016/j.cell.2014.11.037
116. Wang L, Xie L, Zhang Q, Cai X, Tang Y, Hang T, et al. Plasma nuclear and mitochondrial DNA levels in acute myocardial infarction patients. *Coron Artery Dis.* (2015) 26:296–300. doi: 10.1097/MCA.0000000000000231
117. Nakahira K, Kyung SY, Rogers AJ, Gazourian L, Youn S, Massaro AF, et al. Circulating mitochondrial DNA in patients in the ICU as a marker of mortality: derivation and validation. *PLoS Med.* (2013) 10:e1001577; discussion e1001577. doi: 10.1371/journal.pmed.1001577
118. Lam NY, Rainer TH, Chiu RW, Joynt GM, Lo YM. Plasma mitochondrial DNA concentrations after trauma. *Clin Chem.* (2004) 50:213–6. doi: 10.1373/clinchem.2003.025783
119. Chan SR, Blackburn EH. Telomeres and telomerase. *Philos Trans R Soc Lond B Biol Sci.* (2004) 359:109–21. doi: 10.1098/rstb.2003.1370
120. Autexier C, Lue NF. The structure and function of telomerase reverse transcriptase. *Annu Rev Biochem.* (2006) 75:493–517. doi: 10.1146/annurev.biochem.75.103004.142412
121. Mathias SL, Scott AF, Kazazian HH, Jr, Boeke JD, Gabriel A. Reverse transcriptase encoded by a human transposable element. *Science.* (1991) 254:1808–10. doi: 10.1126/science.1722352
122. Piskareva O, Denmukhametova S, Schmatchenko V. Functional reverse transcriptase encoded by the human LINE-1 from baculovirus-infected insect cells. *Protein Expr Purif.* (2003) 28:125–30. doi: 10.1016/S1046-5928(02)00655-1
123. Clements AP, Singer MF. The human LINE-1 reverse transcriptase: effect of deletions outside the common reverse transcriptase domain. *Nucleic Acids Res.* (1998) 26:3528–35. doi: 10.1093/nar/26.15.3528
124. Dombroski BA, Feng Q, Mathias SL, Sassaman DM, Scott AF, Kazazian HH, Jr, et al. An *in vivo* assay for the reverse transcriptase of human retrotransposon L1 in *Saccharomyces cerevisiae*. *Mol Cell Biol.* (1994) 14:4485–92. doi: 10.1128/MCB.14.7.4485
125. Dai L, Huang Q, Boeke JD. Effect of reverse transcriptase inhibitors on LINE-1 and Ty1 reverse transcriptase activities and on LINE-1 retrotransposition. *BMC Biochem.* (2011) 12:18. doi: 10.1186/1471-2091-12-18
126. Jones RB, Garrison KE, Wong JC, Duan EH, Nixon DE, Ostrowski MA. Nucleoside analogue reverse transcriptase inhibitors differentially inhibit human LINE-1 retrotransposition. *PLoS ONE.* (2008) 3:e1547. doi: 10.1371/journal.pone.0001547
127. Goodier JL, Cheung LE, Kazazian HH Jr. Mapping the LINE1 ORF1 protein interactome reveals associated inhibitors of human retrotransposition. *Nucleic Acids Res.* (2013) 41:7401–19. doi: 10.1093/nar/gkt512
128. Brouha B, Schustak J, Badge RM, Lutz-Prigge S, Farley AH, Moran JV, et al. Hot L1s account for the bulk of retrotransposition in the human population. *Proc Natl Acad Sci USA.* (2003) 100:5280–5. doi: 10.1073/pnas.0831042100
129. Dewannieux M, Esnault C, Heidmann T. LINE-mediated retrotransposition of marked Alu sequences. *Nat Genet.* (2003) 35:41–8. doi: 10.1038/ng1223
130. Esnault C, Maestre J, Heidmann T. Human LINE retrotransposons generate processed pseudogenes. *Nat Genet.* (2000) 24:363–7. doi: 10.1038/74184
131. Stetson DB, Medzhitov R. Recognition of cytosolic DNA activates an IRF3-dependent innate immune response. *Immunity.* (2006) 24:93–103. doi: 10.1016/j.immuni.2005.12.003
132. Yang YG, Lindahl T, Barnes DE. Trex1 exonuclease degrades ssDNA to prevent chronic checkpoint activation and autoimmune disease. *Cell.* (2007) 131:873–86. doi: 10.1016/j.cell.2007.10.017
133. Rice GI, Bond J, Asipu A, Brunette RL, Manfield IW, Carr IM, et al. Mutations involved in Aicardi-Goutieres syndrome implicate SAMHD1 as regulator of the innate immune response. *Nat Genet.* (2009) 41:829–32. doi: 10.1038/ng.373
134. Beck-Engeser G, Eilat D, Wabl M. An autoimmune disease prevented by anti-retroviral drugs. *Retrovirology.* (2011) 8:91. doi: 10.1186/1742-4690-8-91
135. Achleitner M, Klefisch M, Hennig A, Peschke K, Polikarpova A, Oertel R, et al. Lack of Trex1 causes systemic autoimmunity despite the presence of antiretroviral drugs. *J Immunol.* (2017) 199:2261–9. doi: 10.4049/jimmunol.1700714
136. Mavragani CP, Sagalovskiy I, Guo Q, Nezos A, Kapsogeorgou EK, Lu P, et al. Expression of long interspersed nuclear element 1 retroelements and induction of type I interferon in patients with systemic autoimmune disease. *Arthritis Rheumatol.* (2016) 68:2686–96. doi: 10.1002/art.39795
137. Mavragani CP, Nezos A, Sagalovskiy I, Seshan S, Kirou KA, Crow MK. Defective regulation of L1 endogenous retroelements in primary Sjogren's

- syndrome and systemic lupus erythematosus: role of methylating enzymes. *J Autoimmun.* (2018) 88:75–82. doi: 10.1016/j.jaut.2017.10.004
138. Huang X, Su G, Wang Z, Shangquan S, Cui X, Zhu J, et al. Hypomethylation of long interspersed nucleotide element-1 in peripheral mononuclear cells of juvenile systemic lupus erythematosus patients in China. *Int J Rheum Dis.* (2014) 17:280–90. doi: 10.1111/1756-185X.12239
  139. Lim YW, Sanz LA, Xu X, Hartono SR, Chedin F. Genome-wide DNA hypomethylation and RNA:DNA hybrid accumulation in Aicardi-Goutieres syndrome. *Elife.* (2015) 4: e08007. doi: 10.7554/eLife.08007
  140. Javierre BM, Fernandez AF, Richter J, Al-Shahrour F, Martin-Subero JI, Rodriguez-Ubreva J, et al. Changes in the pattern of DNA methylation associate with twin discordance in systemic lupus erythematosus. *Genome Res.* (2010) 20:170–9. doi: 10.1101/gr.100289.109
  141. Balada E, Ordi-Ros J, Serrano-Acedo S, Martinez-Lastao L, Rosa-Leyva M, Vilardell-Tarres M. Transcript levels of DNA methyltransferases DNMT1, DNMT3A and DNMT3B in CD4+ T cells from patients with systemic lupus erythematosus. *Immunology.* (2008) 124:339–47. doi: 10.1111/j.1365-2567.2007.02771.x
  142. Nawrocki MJ, Majewski D, Puszczewicz M, Jagodzinski PP. Decreased mRNA expression levels of DNA methyltransferases type 1 and 3A in systemic lupus erythematosus. *Rheumatol Int.* (2017) 37:775–83. doi: 10.1007/s00296-017-3711-8
  143. Momparler RL. Molecular, cellular and animal pharmacology of 5-aza-2'-deoxycytidine. *Pharmacol Ther.* (1985) 30:287–99. doi: 10.1016/0163-7258(85)90053-1
  144. Howard G, Eiges R, Gaudet F, Jaenisch R, Eden A. Activation and transposition of endogenous retroviral elements in hypomethylation induced tumors in mice. *Oncogene.* (2008) 27:404–8. doi: 10.1038/sj.onc.1210631
  145. Quddus J, Johnson KJ, Gavalchin J, Amento EP, Chrisp CE, Yung RL, et al. Treating activated CD4+ T cells with either of two distinct DNA methyltransferase inhibitors, 5-azacytidine or procainamide, is sufficient to cause a lupus-like disease in syngeneic mice. *J Clin Invest.* (1993) 92:38–53. doi: 10.1172/JCI116576
  146. Cornacchia E, Golbus J, Maybaum J, Strahler J, Hanash S, Richardson B. Hyalazine and procainamide inhibit T cell DNA methylation and induce autoreactivity. *J Immunol.* (1988) 140:2197–200.
  147. Lieberman MW, Beach LR, Palmiter RD. Ultraviolet radiation-induced metallothionein-I gene activation is associated with extensive DNA demethylation. *Cell.* (1983) 35:207–14. doi: 10.1016/0092-8674(83)90223-4
  148. Li TH, Schmid CW. Differential stress induction of individual Alu loci: implications for transcription and retrotransposition. *Gene.* (2001) 276:135–41. doi: 10.1016/S0378-1119(01)00637-0
  149. Kalogirou EM, Piperi EP, Tosios KI, Tsiambas E, Fanourakis G, Sklavounou A. Ductal cells of minor salivary glands in Sjogren's syndrome express LINE-1 ORF2p and APOBEC3B. *J Oral Pathol Med.* (2018) 47:179–85. doi: 10.1111/jop.12656
  150. Kell AM, Gale M, Jr. RIG-I in RNA virus recognition. *Virology.* (2015) 479–480:110–21. doi: 10.1016/j.virol.2015.02.017
  151. Goubau D, Schlee M, Deddouch S, Pruijssers AJ, Zillinger T, Goldeck M, et al. Antiviral immunity via RIG-I-mediated recognition of RNA bearing 5'-diphosphates. *Nature.* (2014) 514:372–5. doi: 10.1038/nature13590
  152. Robinson T, Kariuki SN, Franek BS, Kumabe M, Kumar AA, Badaracco M, et al. Autoimmune disease risk variant of IFIH1 is associated with increased sensitivity to IFN-alpha and serologic autoimmunity in lupus patients. *J Immunol.* (2011) 187:1298–303. doi: 10.4049/jimmunol.1100857
  153. Gorman JA, Hundhausen C, Errett JS, Stone AE, Allenspach EJ, Ge Y, et al. The A946T variant of the RNA sensor IFIH1 mediates an interferon program that limits viral infection but increases the risk for autoimmunity. *Nat Immunol.* (2017) 18:744–52. doi: 10.1038/ni.3766
  154. Hung T, Pratt GA, Sundaraman B, Townsend MJ, Chaivorapol C, Bhargale T, et al. The Ro60 autoantigen binds endogenous retroelements and regulates inflammatory gene expression. *Science.* (2015) 350:455–9. doi: 10.1126/science.aac7442
  155. Chung H, Calis JJA, Wu X, Sun T, Yu Y, Sarbanes SL, et al. Human ADARI prevents endogenous RNA from triggering translational shutdown. *Cell.* (2018) 172:811–24. doi: 10.1016/j.cell.2017.12.038
  156. Maraia RJ, Intine RV. Recognition of nascent RNA by the human La antigen: conserved and divergent features of structure and function. *Mol Cell Biol.* (2001) 21:367–79. doi: 10.1128/MCB.21.2.367-379.2001
  157. Xue D, Shi H, Smith JD, Chen X, Noe DA, Cedervall T, et al. A lupus-like syndrome develops in mice lacking the Ro 60-kDa protein, a major lupus autoantigen. *Proc Natl Acad Sci USA.* (2003) 100:7503–8. doi: 10.1073/pnas.0832411100
  158. Chen X, Smith JD, Shi H, Yang DD, Flavell RA, Wolin SL. The Ro autoantigen binds misfolded U2 small nuclear RNAs and assists mammalian cell survival after UV irradiation. *Curr Biol.* (2003) 13:2206–11. doi: 10.1016/j.cub.2003.11.028
  159. Stein AJ, Fuchs G, Fu C, Wolin SL, Reinisch KM. Structural insights into RNA quality control: the Ro autoantigen binds misfolded RNAs via its central cavity. *Cell.* (2005) 121:529–39. doi: 10.1016/j.cell.2005.03.009
  160. Merrill JT, Wallace DJ, Petri M, Kirou KA, Yao Y, White WI, et al. Safety profile and clinical activity of sifalimumab, a fully human anti-interferon alpha monoclonal antibody, in systemic lupus erythematosus: a phase I, multicentre, double-blind randomised study. *Ann Rheum Dis.* (2011) 70:1905–13. doi: 10.1136/ard.2010.144485
  161. Rosman Z, Shoenfeld Y, Zandman-Goddard G. Biologic therapy for autoimmune diseases: an update. *BMC Med.* (2013) 11:88. doi: 10.1186/1741-7015-11-88
  162. Kariuki SN, Crow MK, Niewold TB. The PTPN22 C1858T polymorphism is associated with skewing of cytokine profiles toward high interferon-alpha activity and low tumor necrosis factor alpha levels in patients with lupus. *Arthritis Rheum.* (2008) 58:2818–23. doi: 10.1002/art.23728
  163. Goldenberg MM. Etanercept, a novel drug for the treatment of patients with severe, active rheumatoid arthritis. *Clin Ther.* (1999) 21:75–87; discussion 1–2. doi: 10.1016/S0149-2918(00)88269-7
  164. Moreland LW, Schiff MH, Baumgartner SW, Tindall EA, Fleischmann RM, Bulpitt KJ, et al. Etanercept therapy in rheumatoid arthritis. a randomized, controlled trial. *Ann Intern Med.* (1999) 130:478–86. doi: 10.7326/0003-4819-130-6-199903160-00004
  165. Weinblatt ME, Kremer JM, Bankhurst AD, Bulpitt KJ, Fleischmann RM, Fox RI, et al. A trial of etanercept, a recombinant tumor necrosis factor receptor:Fc fusion protein, in patients with rheumatoid arthritis receiving methotrexate. *N Engl J Med.* (1999) 340:253–9. doi: 10.1056/NEJM199901283400401
  166. Lodge KM, Knolle MD, Jha A. Asthma: from diagnosis to endotype to treatment. *Am J Respir Crit Care Med.* (2018) 197:1065–7. doi: 10.1164/rccm.201710-2139RR

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Mustelin, Lood and Giltaiy. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.