



Corrigendum: Spatial Lymphocyte Dynamics in Lymph Nodes Predicts the Cytotoxic T Cell Frequency Needed for HIV Infection Control

Dmitry Grebennikov^{1,2,3*}, Anass Bouchnita⁴, Vitaly Volpert^{3,5,6}, Nikolay Bessonov⁷, Andreas Meyerhans^{8,9} and Gennady Bocharov^{2,10*}

¹ Moscow Institute of Physics and Technology, National Research University, Dolgoprudny, Russia, ² Marchuk Institute of Numerical Mathematics, Russian Academy of Sciences, Moscow, Russia, ³ Peoples' Friendship University of Russia (RUDN University), Moscow, Russia, ⁴ Division of Scientific Computing, Department of Information Technology, Uppsala University, Uppsala, Sweden, ⁵ Institut Camille Jordan, UMR 5208 CNRS, University Lyon 1, Villeurbanne, France, ⁶ INRIA Team Dracula, INRIA Lyon La Doua, Villeurbanne, France, ⁷ Institute of Problems of Mechanical Engineering, Russian Academy of Sciences, Saint Petersburg, Russia, ⁸ Infection Biology Laboratory, Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona, Spain, ⁹ Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain, ¹⁰ Sechenov First Moscow State Medical University, Moscow, Russia

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*Correspondence:

Dmitry Grebennikov
dmitry.ew@gmail.com
Gennady Bocharov
bocharov@m.inm.ras.ru

Specialty section:

This article was submitted to
T Cell Biology,
a section of the journal
Frontiers in Immunology

Received: 18 June 2019

Accepted: 19 June 2019

Published: 03 July 2019

Citation:

Grebennikov D, Bouchnita A,
Volpert V, Bessonov N, Meyerhans A
and Bocharov G (2019) Corrigendum:
Spatial Lymphocyte Dynamics in
Lymph Nodes Predicts the Cytotoxic
T Cell Frequency Needed for HIV
Infection Control.
Front. Immunol. 10:1538.
doi: 10.3389/fimmu.2019.01538

Keywords: lymphoid tissue, cell motility, HIV infection, cytotoxic T cell scanning, multicellular dynamics, dissipative particle dynamics, stochastic differential equation

A Corrigendum on

Spatial Lymphocyte Dynamics in Lymph Nodes Predicts the Cytotoxic T Cell Frequency Needed for HIV Infection Control

by Grebennikov, D., Bouchnita, A., Volpert, V., Bessonov, N., Meyerhans, A., and Bocharov, G. (2019). *Front. Immunol.* 10:1213. doi: 10.3389/fimmu.2019.01213

In the original article, there was a typo in **Figure 2A** color legend as published. The colored circles denoting Ag-specific and non-specific T cells should be swapped. That is, the dark green color should represent Ag-specific CD4⁺ TCs, the light green color—non-specific CD4⁺ TCs; the dark blue color should represent Ag-specific CD8⁺ TCs, the light blue color—non-specific CD8⁺ TCs. The corrected **Figure 2** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2019 Grebennikov, Bouchnita, Volpert, Bessonov, Meyerhans and Bocharov. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

