



Corrigendum: Recirculation and Residency of T Cells and Tregs: Lessons Learnt in Anacapri

Silvia Piconese^{1,2}, Silvia Campello³ and Ambra Natalini^{4,5*}

¹ Dipartimento di Scienze Cliniche Internistiche, Anestesiologiche e Cardiovascolari, Sapienza Università di Roma, Rome, Italy, ² Laboratory Affiliated to Istituto Pasteur Italia – Fondazione Cenci Bolognetti, Rome, Italy, ³ Department of Biology, University of Rome Tor Vergata, Rome, Italy, ⁴ Institute of Molecular Biology and Pathology, National Research Council (CNR), Rome, Italy, ⁵ Dipartimento di Medicina Molecolare (DMM), Sapienza Università di Roma, Rome, Italy

Keywords: T cells, Tregs, cell migration, cell cycle, recirculation

A Corrigendum on

OPEN ACCESS

Approved by:
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

***Correspondence:**
Ambra Natalini
ambra.natalini@uniroma1.it

Specialty section:
This article was submitted to
Molecular Innate Immunity,
a section of the journal
Frontiers in Immunology

Received: 31 May 2020

Accepted: 01 June 2020

Published: 09 July 2020

Citation:
Piconese S, Campello S and
Natalini A (2020) Corrigendum:
Recirculation and Residency of T Cells
and Tregs: Lessons Learnt in
Anacapri. *Front. Immunol.* 11:1401.
doi: 10.3389/fimmu.2020.01401

Recirculation and Residency of T Cells and Tregs: Lessons Learnt in Anacapri

by Piconese, S., Campello, S., and Natalini, A. (2020). *Front. Immunol.* 11:682.
doi: 10.3389/fimmu.2020.00682

In the original article, there was a error in the legend of **Figure 1** as published. In the original manuscript we neglected to cite in the legend of **Figure 1** the article from which the figure was taken and adapted. The correct legend appears below.

Figure 1. Cell cycle analysis of antigen-specific CD8 T cells in the blood after vaccination. Female Balb/c mice were primed and boosted with viral vectors expressing the model antigen gag of HIV-1. At days (*d*) 3, 7, and 44, post-boost blood was collected and blood cells were analyzed with our new method. The figure shows a typical ki67/DNA staining profile of gag-specific CD8 T cells in the blood. Fluorescence Minus One (FMO) controls (*left*) and Ki67 staining (*right*) are shown, as indicated; the *numbers* represent the percentages of cells in the corresponding quadrant. Figure adapted from (30).

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Piconese, Campello and Natalini. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.