



Mucosal Nanoemulsion Allergy Vaccine Suppresses Alarmin Expression and Induces Bystander Suppression of Reactivity to Multiple Food Allergens

Mohammad Farazuddin¹, Jeffrey J. Landers¹, Katarzyna W. Janczak¹, Hayley K. Lindsey¹, Fred D. Finkelman², James R. Baker Jr¹ and Jessica J. O’Konek^{1*}

OPEN ACCESS

Edited by:

Karen Laky,
National Institutes of Health (NIH),
United States

Reviewed by:

Ya-Jen Chang,
Academia Sinica, Taiwan
Giorgio Walter Canonica,
University of Genoa, Italy

*Correspondence:

Jessica J. O’Konek
jjoz@umich.edu

Specialty section:

This article was submitted to
Molecular Innate Immunity,
a section of the journal
Frontiers in Immunology

Received: 26 August 2020

Accepted: 06 January 2021

Published: 25 February 2021

Citation:

Farazuddin M, Landers JJ,
Janczak KW, Lindsey HK,
Finkelman FD, Baker JR and
O’Konek JJ (2021) Mucosal
Nanoemulsion Allergy Vaccine
Suppresses Alarmin Expression and
Induces Bystander Suppression of
Reactivity to Multiple Food Allergens.
Front. Immunol. 12:599296.
doi: 10.3389/fimmu.2021.599296

¹ Mary H. Weiser Food Allergy Center, University of Michigan, Ann Arbor, MI, United States, ² Division of Allergy, Immunology and Rheumatology, University of Cincinnati College of Medicine, Cincinnati, OH, United States

We have demonstrated that intranasal immunotherapy with allergens formulated in a nanoemulsion (NE) mucosal adjuvant suppresses Th2/IgE-mediated allergic responses and protects from allergen challenge in murine food allergy models. Protection conferred by this therapy is associated with strong suppression of allergen specific Th2 cellular immunity and increased Th1 cytokines. Here we extend these studies to examine the effect of NE-allergen immunization in mice sensitized to multiple foods. Mice were sensitized to both egg and peanut and then received NE vaccine formulated with either one or both of these allergens. The animals were then subjected to oral challenges with either egg or peanut to assess reactivity. Immunization with NE formulations containing both egg and peanut markedly reduced reactivity after oral allergen challenge with either allergen. Interestingly, mice that received the vaccine containing only peanut also had reduced reactivity to challenge with egg. Protection from oral allergen challenge was achieved despite the persistence of allergen-specific IgE and was associated with strong suppression of both Th2-polarized immune responses, alarmins and type 2 innate lymphoid cells (ILC2). NE-induced bystander suppression of reactivity required IFN- γ and the presence of an allergen in the NE vaccine. These results demonstrate that anaphylactic reactions to food allergens can be suppressed using allergen-specific immunotherapy without having to eliminate allergen-specific IgE and suggests that modulation of Th2 immunity towards one allergen may induce bystander effects that suppress reactivity to other allergens through the induction of IFN- γ and suppression of alarmins in the intestine. In addition, these data suggest that a NE vaccine for a single food allergen may lead to a global suppression of allergic responses to multiple foods.

Keywords: adjuvant, alarmins, allergy treatment, food allergy, immunotherapy vaccines and mechanisms, vaccine

INTRODUCTION

Food allergy is an emerging epidemic that now affects up to 15 million people in the US, including 8% of children. The economic burden of food allergy in the US alone exceeds \$24.8 billion (1). Allergen-specific immunotherapy for food allergy involves the progressive administration of increasing amounts of a specific allergen by one of several routes and has been the primary approach to suppress allergic reactivity. This approach, however, does not provide long-term protection following cessation of therapy and requires prolonged treatment protocols burdening to patients and their families. Specifically, subcutaneous immunotherapy to food allergens showed promise for protection against IgE-mediated food allergies, however significant adverse reactions limited successful implementation (2–4). Sublingual, oral (OIT) and epicutaneous immunotherapy have demonstrated efficacy in animal models and human trials, however these approaches desensitize only a portion of patients and the protection achieved is rapidly lost after cessation of the therapy (5–11). Thus, there is a need to understand immune mechanisms that modulate Th2-biased immune responses to food and could lead to long-lasting protection from allergic reactions.

The primary immunologic mechanism of allergic hypersensitivity is the induction of Th2-polarized cellular immune responses leading to the production of allergen-specific IgE antibodies critical for mast cell activation. Th2 cytokines also are critical mediators of local allergic inflammation, including IL-4- and IL-13-dependent mucus production and IL-5-mediated eosinophil recruitment (12). Oral or subcutaneous allergen immunotherapy (AIT) appears to achieve desensitization to the allergen by temporarily reducing Th2-biased immunity and allergen-specific IgE. While OIT has been proven clinically useful for treating food allergy, it has not induced a long term redirection of allergen-specific immunity away from a Th2 phenotype (13). Thus, interest has been directed toward new strategies that are able to permanently suppress Th2 cellular immune responses or redirect these cellular Th2 responses towards a Th1 phenotype (14, 15).

We have developed a novel nasal vaccine-based immunotherapy system employing a nanoscale oil-in-water emulsion (nanoemulsion, NE) adjuvant. When administered intranasally (i.n.) with viral and bacterial antigens, this formulation induces robust systemic and mucosal immunity, and cell-mediated immune responses are polarized towards Th1 and Th17 (16–24). We have previously reported that therapeutic immunization with NE and antigen/allergen can suppress established Th2-polarized immunity and protect from allergen challenge in murine models of food allergy (25–28). NE-based allergy vaccines induced sustained unresponsiveness lasting at least 16 weeks, and protection was associated with increased IL-10 and regulatory T cells. Here, we extended our previous work to determine the ability of NE-based allergy vaccines to broadly suppress allergic reactions in mice sensitized to more than one food.


MATERIALS AND METHODS

Antigen and Adjuvants

Nanoemulsion adjuvant (NE) was produced by a high speed emulsification of ultra-pure soybean oil with cetyl pyridinium chloride, Tween 80 and ethanol in water, with resultant NE droplets with average 350–400 nm diameter (17, 29). Aluminum hydroxide (alum, alhydrogel) was purchased from InvivoGen. Peanut extract (Greer) was used for all intraperitoneal (i.p.) and intranasal (i.n.) immunizations. For oral/intragastric (i.g.) challenges, peanut flour (12% fat, light roast, Byrd Mill) was solubilized in PBS. Endotoxin-free ovalbumin (OVA) was purchased from Hyglos. Endotoxin content of all vaccine components was determined by a limulus amoebocyte lysate (LAL) assay (Pierce).

Mice and Immunizations

Specific pathogen-free BALB/c mice (females 3 weeks old) were purchased from Jackson Laboratory. Mice were 4 weeks of age at the onset of the experiment. The experimental design is shown in **Figure 1**. In all experiments, allergic sensitization was induced with i.p. immunizations of 20 μ g OVA and 20 μ g peanut extract



(PN) adsorbed on 1 mg alum at week 0. The experimental design for each experiment is as follows.

Assessment of Combination Vaccine Containing OVA and Peanut

Mice received i.n. immunizations of 12 μ l (6 μ l/nare) of a formulation containing 20 μ g of OVA and 20 μ g PN at weeks 4, 8 and 12. Sensitized control mice received i.n. PBS. Beginning 4 weeks after the final i.n. vaccine dose, anaphylaxis was induced by repeated oral challenge with allergen. Mice were challenged orally every other day for a total of 7 gavages (30). For each challenge, mice were fasted for 5–6 h to ensure gastric emptying and then were challenged by oral gavage (i.g.) with 0.2 ml containing of 10 mg OVA and 10 mg peanut. Reactivity was measured as described below.

Assessment of Bystander Protection

Mice received i.n. immunizations of a formulation containing either 20 μ g of OVA or 20 μ g PN mixed with 20% NE at weeks 4, 8 and 12. Mice were orally challenged with either 10 mg OVA or 10 mg peanut protein and reactivity was determined as described below.

Assessment of Requirement of Specific Antigen for Protection

Following the same schedule as described above, sensitized mice received i.n. immunizations of 20% NE alone (no antigen) or 20 μ g of OVA in 20% NE. Sensitized control mice received i.n. PBS. Mice were challenged orally with OVA and reactivity was determined as described above. In a separate experiment following the same schedule, mice were immunized with 20 μ g hepatitis B surface antigen (HBsAg) mixed with 20% NE.

IFN- γ Depletion Experiments

Mice were sensitized to OVA and peanut at week 0 and received i.n. PN-NE vaccine at weeks 4, 8 and 12. Mice were subjected to our repeated OVA challenge protocol beginning at week 16. Mice were injected i.p. with 0.5 mg anti-IFN- γ (XMG-6) or isotype control rat IgG1 (GL113) [both produced in house (31)] the day before starting oral challenges and every 4 days until the final challenge. All animal procedures were performed according to the National Institutes of Health guide for the care and use of laboratory animals and approved by the University of Michigan Institutional Animal Care and Use Committee (IACUC).

Assessment of Hypersensitivity Reactions

Anaphylactic symptoms were evaluated for one hour following the final (7th) challenge with OVA using the following scoring system [modified from (32, 33)]: 0, no symptoms; 0.5, transient rubbing and scratching; 1, prolonged rubbing and scratching around the nose, eyes or head; 2, puffiness around the eyes or mouth, diarrhea, piloerection, and/or decreased activity with increased respiratory rate; 3, labored respiration, wheezing, stridor, and/or cyanosis around the mouth and tail; 4, tremor, convulsion, no activity after prodding and/or moribund; 5, death. Rectal temperature was monitored prior to and every 15 min for 60 min following challenge, and the maximum temperature change from baseline was reported. Mice were bled 60 min following challenge, and

serum mouse mast cell protease-1 (MCPT-1) was determined by ELISA (eBioscience). To determine hemoconcentration, blood was drawn 60 min following challenge into heparinized capillary tubes and centrifuged for 5 min at 10,000 rpm. Hematocrit values were calculated as the length of packed RBCs as a percentage of the total length of serum and red cells in the capillary tube.

Measurement of Serum IgE

Sera were obtained by cardiac puncture post-euthanasia one day after the final (7th) allergen challenge. Serum was separated from whole blood by centrifugation at 1,500 \times g for 5 min after allowing coagulation for 30–60 min at room temperature. Serum samples were stored at -20°C until analyzed. OVA-specific IgE antibody levels were determined by ELISA. Serially diluted serum samples were incubated on microtiter plates coated with 20 μ g/ml OVA. IgE antibodies were detected with alkaline phosphatase conjugated anti-mouse IgE (Rockland) and Sigma FastTM p-nitrophenyl phosphate substrate and quantified by measuring the optical density (OD) at 405 nm. The antibody concentrations are presented as endpoint titers defined as the reciprocal of the highest serum dilution producing an OD above background of naive sera. The cutoff value is determined as the OD (mean+2 standard deviations) of the corresponding dilution of naive sera (34, 35).

Analysis of Cytokine Production

Mice were sacrificed one day after the final (7th) oral challenge, and mesenteric lymph nodes were harvested. The cellular recall response was evaluated in lymphocytes isolated from mesenteric lymph nodes. Single cell lymphocyte suspensions were cultured *ex vivo* \pm OVA (20 μ g/ml) at 37 $^{\circ}\text{C}$. After 72 h, cytokine secretion was measured in cell culture supernatants using Luminex Multiplex detection system (Millipore). For real-time PCR analysis, RNA was isolated from duodenum homogenates with an RNeasy mini kit (Qiagen), and cDNA was generated with a Superscript II reverse transcription kit (Invitrogen). qPCR was performed with SYBR green master mix and commercially available primer sets (Bio-Rad). Values were normalized to GAPDH and displayed as fold induction over control samples.

Lamina Propria Mononuclear Cells Isolation

Mice were sacrificed one day after the final (7th) oral challenge. Small intestine (SI, 15 cm) was dissected from the mouse and Peyer's patches were trimmed off. SI was cut longitudinally and washed with PBS thoroughly to remove ingested food. SI was incubated in a petri dish with 10 ml PBS with EDTA (5 mM) for 10 min on ice. Intestines were washed with PBS (no EDTA) by vortexing vigorously to remove the epithelial cells. These two steps were repeated 3–4 times until the tissue became clear. Tissue was minced finely and transferred to 8 ml digestion buffer (16 mg collagenase A (Roche) and 1.6mg DNase I (Roche) in RPMI (10% FBS) and incubated at 37 $^{\circ}\text{C}$ for 30 min. After incubation, digested tissue was passed through a 10 ml syringe with 18G needle a few times. Liberated cells were filtered through 70 μ m filter. The cell suspension was washed by adding 20 ml of RPMI with 10% FBS. The cell pellet was suspended in 44% Percoll (4 ml) and loaded on 67% Percoll (3 ml) for

centrifugation. A mononuclear cell gradient was created by spinning the cells down at 1,800 rpm for 20 min at room temperature with centrifuge acceleration set at 5 and deceleration set to 0. The middle interphase of mononuclear cells was collected from the interface and washed again with RPMI (10% FBS). The obtained cells were counted and used for subsequent analysis.

Antibodies

All the antibodies used for flow cytometry were purchased from eBioscience, Biologend and BD biosciences. For cell surface staining, a lineage cocktail consisting anti-mouse CD3 (clone 145-2C11), anti-mouse Ly-6G/Ly-6C (clone RB6-8C5), anti-mouse CD11b (clone M1/70), anti-mouse CD45R/B220 (clone RA3-6B2), and anti-mouse TER-119/Erythroid cells (clone Ter-119) was used. FITC-streptavidin was used to stain biotin labelled primary antibody cocktail. Other antibodies used were rat anti-mouse CD45, anti-mouse CD127 (clone A7R34), anti-mouse CD90.2 (clone 53-2.1), anti-mouse KLRG1 (clone 2F1), and anti-mouse GATA3 (clone TWAJ). Foxp3 fixation and permeabilization kit (eBioscience) was used for intracellular staining.

Flow Cytometry

Cells were stained with lineage antibody cocktail on ice for 20 min followed by washing with flow staining buffer (PBS with 0.1% BSA) two times. Cells were then incubated with FITC-streptavidin antibody on ice for another 20 min. Cells were washed two times with flow staining buffer. Cells were then stained with live-dead eF450, CD45, CD127, CD90.2, and KLRG1 on ice for 20 min followed by washing two times with flow staining buffer. Intracellular staining of GATA3 was done using Foxp3 fixation and permeabilization kit as per the manufacturer's protocol (eBiosciences). Samples were acquired on Novocyt 3000 (Acea biosciences) and data were analyzed using FlowJo v10.1. The gating strategy for identifying ILC2s is shown in **Supplementary Figure 1**.

Statistics

Results presented here are the representatives of at least two independent experiments. Each experiment contained 8–10 mice per group. Statistical comparisons were assessed by the Mann-Whitney test using GraphPad Prism version 8 (GraphPad Software). The p value < 0.05 was considered as significant.

RESULTS

Intranasal Immunization With Allergens in NE Adjuvant Suppresses Allergic Reactions in Polysensitized Mice

BALB/c mice were sensitized to egg and peanut and immunized with OVA and peanut formulated in NE to determine if suppression of reactivity to 2 allergens could be achieved simultaneously. Sensitized control mice had profound physiological reactions to challenge as indicated by severe symptoms of anaphylactic shock, including diarrhea, labored

respiration, wheezing, lack of activity when prodded, core body temperature loss of greater than 2°C, hemoconcentration and increased mast cell degranulation (MCPT-1) (**Figure 1**). The NE vaccine markedly suppressed these responses to allergen challenge. Anaphylaxis symptoms were markedly reduced to mild symptoms such as pruritus or reduced activity (**Figure 1B**), and the incidence of diarrhea was reduced from 100–40% (**Figure 1C**). Mice treated with the NE vaccine also were protected from hypovolemic shock and experienced minimal body temperature loss while hemoconcentration also was prevented (**Figures 1D, E**). MCPT-1 measured in serum following challenge was used to assess mast cell degranulation. Consistent with the clinical symptoms of allergic reaction, immunized mice had significant reductions in MCPT-1, with average levels of 0.6 µg/ml compared with 16 µg/ml in sensitized control mice (**Figure 1F**; $p=0.0079$).

Immunization of Polysensitized Mice With NE Adjuvant Induces Bystander Suppression of Allergic Reactivity

Next, we wanted to assess the effects of immunotherapy with NE and only one allergen on protection in polysensitized animals. Mice were again sensitized to both egg and peanut, and then were nasally treated with either OVA or peanut or both allergens formulated in NE. In general, mice were protected from allergic reaction to whatever allergen was contained in the vaccines, and protection for each allergen was similar if the mice were treated with the vaccine containing either a single allergen or both allergens (**Figure 2**). Surprisingly, mice that were treated with the OVA-NE vaccine were also protected from reactivity to challenge with peanut and mice treated with the PN-NE vaccine were protected from challenge with OVA. While there was a trend that this “bystander protection” was not as complete as protection induced by immunotherapy with NE and both allergens, these differences were not significant, and mice immunized with only one allergen in NE had significantly less severe allergic reactions compared with sensitized control mice that did not receive the i.n. vaccine. Bystander suppression of reactivity persisted for at least 8 weeks after the final vaccine dose (**Supplementary Figure 2**).

We next aimed to determine if the observed bystander suppression of allergic reactivity required immunotherapy with NE and at least one allergen to which the mice were sensitized. As shown in **Figures 3A–C**, in mice that were sensitized to OVA and peanut, i.n. instillation of NE alone (no allergen) did not induce any suppression of allergic reactivity to oral OVA challenge, as reactivity in these mice was equivalent to sensitized control mice and significantly more severe than mice that received the OVA-NE vaccines. Similarly, NE formulated with an unrelated antigen [hepatitis B surface antigen (HBsAg)] did not confer protection from challenge with OVA (**Figures 3D–F**).

Intranasal Immunization With NE Adjuvant Suppresses Allergy Associated Th2 Responses and Alarmin Expression

While food allergic reactions are dependent upon the presence of allergen-specific IgE, many patients with allergen-specific IgE to




FIGURE 2 | Immunization of polysensitized mice with NE and one allergen provides protection against reactivity to another allergen. As described in Figure 1A, mice were sensitized with OVA and peanut-alum and treated i.n. with 3 administrations of PBS (sensitized control), ova and peanut-NE (OVA+PN-NE), OVA-NE or peanut-NE (PN-NE). Mice were challenged orally with (A) OVA or (B) peanut and temperature change and symptoms of anaphylaxis were monitored. Hemoconcentration was determined by hematocrit. Levels of MCPT-1 in the serum 60 min after challenge were determined by ELISA. Statistically significant differences ($p < 0.05$) are indicated by *.

foods do not clinically react to those foods. In the present study, immunization with PN-NE suppressed allergic reactivity to both peanut and OVA without significantly reducing OVA-specific IgE (Figure 4A). This apparent disconnect between the presence of allergen-specific IgE and reactivity to an allergen suggests that other immune changes are behind the suppression of allergic reactions observed here.

Allergen-specific cytokine secretion was measured to characterize changes to the Th2-biased cellular immune responses associated with allergic disease. Upon stimulation with OVA, cells from (OVA and peanut)-alum sensitized mice produced significant levels of Th2-type cytokines IL-4 and IL-13 but not IFN- γ (Figure 4B). Immunotherapy with either OVA or peanut in NE reduced OVA-specific IL-4 and IL-13 and increased IFN- γ . While OVA-specific IL-17 and IL-10 were increased in mice that received the OVA-NE vaccine, the PN-NE vaccine did not affect OVA-specific production of these cytokines. Because of the interplay between Th2 immunity and expression of alarmins, expression of the genes for the alarmins IL-25, IL-33, and TSLP was also evaluated in the small intestine (36–40). Immunization with the NE vaccines significantly reduced the expression of *Il25*, *Il33* and *Tslp* such that the fold change over expression in naïve mice was approximately 1 (Figure 4C). These data indicate that immunization with NE prevents increased alarmin expression normally observed in allergen sensitized mice.

Intranasal Immunization With NE Adjuvant Suppresses ILC2 Populations

We observed a significant decrease in alarmin expression, in animals that received the NE-allergen vaccines compared to sensitized control animals. It has been previously established that IL-25 and IL-33 acts as activation cytokines and regulate ILC2 populations in allergic inflammation (41–43). ILC2s were quantified in the small intestine lamina propria to determine if reduction in reactivity was associated with reduce ILC2 accumulation in the tissue. ILC2s were increased in the intestine of sensitized mice compared to naïve. Conversely, ILC2s were significantly reduced in OVA-NE-and PN-NE immunized animals compared to sensitized controls (Figure 4D). These data suggest that immunization with either OVA-NE-or PN-NE modulates epithelial alarmin production, which in turn prevents the accumulation of ILC2s in the tissues.

Bystander Protection Induced by NE Allergy Vaccines Requires IFN- γ

Because NE immunization increased IFN- γ , which has been shown to suppress Th2 immunity and alarmins, we hypothesized that bystander protection associated with NE was IFN- γ dependent. Mice were sensitized to OVA and peanut and then treated with PN-NE. During the allergen challenge phase in which mice were treated with OVA, IFN- γ was depleted.




FIGURE 3 | Intranasal administration of NE without allergen does not suppress the allergic response. (A–C) Mice were sensitized with OVA and peanut-alum and treated i.n. with 3 administrations of PBS (sensitized control), OVA-NE (OVA-NE) or NE only (no antigen) or Mice were challenged orally with OVA and temperature change and symptoms of anaphylaxis were monitored. Serum MCPT-1 levels were determined by ELISA. (D–F) In a separate experiment, mice were similarly sensitized with OVA and peanut with alum and treated i.n. with 3 administrations of PBS (sensitized control) or hepatitis B surface antigen-NE (HBsAg-NE). Statistically significant differences ($p < 0.05$) are indicated by *.

Depletion of IFN- γ during the challenge phase completely abrogated the protection induced by the PN-NE vaccine, as mice treated with anti-IFN- γ antibody had severe reactions to challenge, including decreases in core body temperature and increased clinical symptoms, diarrhea, hematocrit and MCPT-1 similar to animals not treated with NE (Figure 5). Suppression of alarmin expression by the PN-NE vaccine was also reversed following IFN- γ depletion, as mice that were depleted of IFN- γ had similar expression of alarmins in the small intestine as sensitize mice that did not receive the vaccine (Figure 6).


DISCUSSION

Thirty to forty percent of patients with food allergies are sensitized to multiple foods (44, 45). While allergen-specific immunotherapy has the potential to relieve the burden of fear of reactivity to specific foods, this approach is more difficult for polysensitized individuals. Approved, allergen-specific immunotherapy for food allergy involves a single food, and regulatory issues may preclude the development of therapies containing multiple foods. While some studies have demonstrated the ability to desensitize patients with OIT for up to 5 foods simultaneously (multi-OIT) (46, 47), the amount of each food required to be consumed daily is a burden for some children as the food required for multi-OIT can be a significant proportion of the daily caloric intake for a child. Given this, there

is interest in the development of therapies for food allergy that work more broadly and are not specific to one allergen.


The present studies demonstrate that NE can be formulated with multiple allergens and lead to reduction of allergic reactivity to all allergens included in the vaccine. NE was formulated with two allergens and maintained therapeutic efficacy for both foods. Based on the protein loading capacity of NE, formulation with more than 2 allergens would also be possible and the therapy only has to be administered three to four times to achieve sustained unresponsiveness of at least 4 months (25–27). Given that long-term protection can be achieved with only a few doses administered at monthly intervals, this approach would significantly reduce the burden on patients with multiple food allergies over daily, allergen-specific immunotherapies.

Importantly, these studies also demonstrate a “bystander” effect in that NE-induced reduction of allergic reactivity with one allergen also reducing reactions to unrelated allergens not included in the vaccine. The bystander protection did require immunization with an allergen to which the animal was previously sensitized, suggesting the non-specific reduction of food allergic reactions required prior sensitization and immune recognition of the vaccine. This bystander suppression of reactivity was maintained for at least 8 weeks, supporting our previous work that demonstrated the long-term sustained unresponsiveness induced by this approach (27). The mechanism of this effect appeared to be redirection of the underlying immune polarization from a Th2 to a Th1



phenotype, especially since no suppressive activity was demonstrated with treatment using either the adjuvant alone or with an immunogen for which there was no pre-existing allergy (HBsAg). This demonstrates that while NE allergy vaccines provide bystander protection, it requires the recognition and induction of an antigen-specific immune response.

There are very few reports of bystander activation of antigen non-specific immune responses, but there is some evidence these do exist. Immunization with aluminum adjuvants (alum) has been shown to induce bystander polarization of Th2 immune responses to unrelated antigens. Specifically, if mice were immunized with alum and “antigen 1”, and then later exposed to antigens 1 and 2 together in the absence of alum, mice developed Th2-polarized immune responses to antigen 2 (48, 49). These studies demonstrated the need for co-administration of antigens to drive the bystander effects. Eisenbarth et al.



described this phenomenon as “collateral priming” and demonstrated that it was based on an adaptive T cell-derived, cytokine dependent mechanism that did not require innate toll-like receptor 4 signaling (49).

Other studies also support the concept of protection to proteins not included in the immunotherapy. For example, after sensitization with whole peanut extract, immunotherapy with individual peanut components such as Ara h 1 or Ara h 2 conferred protection against challenge with the whole allergen, despite the fact that mice were sensitized to multiple protein components of the allergen that the immunotherapy did not target (50–52). It also has been reported that epicutaneous immunotherapy with peanut in mice generates a Treg population that can prevent the subsequent sensitization to either peanut or house dust mite (53, 54). These Tregs were broadly effective at suppressing inflammation, including T cell-mediated intestinal inflammation in a mouse model of colitis (55). Along with our data, these reports suggest that antigen-specific T cell responses to newly introduced immunogens can be




FIGURE 6 | IFN- γ is required for suppression of alarmins by NE allergy vaccines. Mice were sensitized with OVA and peanut-alum and treated i.n. with 3 administrations of PBS (sensitized control) or peanut-NE (PN-NE). Mice were challenged orally with OVA and IFN- γ was depleted during the challenge phase. Duodenum samples were homogenized and mRNA was extracted to determine relative gene expression compared to GAPDH. Statistically significant differences ($p < 0.05$) are indicated by *.

influenced by non-specific elements (mainly regulatory cells and cytokines) from ongoing immune responses to other antigens.

While our previous work has shown that induction of Tregs and IL-10 were important to NE mediated immune modulation (25, 28), we now include the production of IFN- γ as critical for allergic protection. Interestingly, while the NE vaccines induced antigen-specific production of IL-17 and IL-10, there was no bystander modulation of these cytokines, as immunization with PN-NE did not induce production of OVA-specific IL-17 or IL-10. The significant increase in OVA-specific IFN- γ production in PN-NE-immunized mice suggested a key role of IFN- γ in driving the bystander modulation of allergen-specific immune responses and reduction of allergic reactivity. The induction of Th1 cytokines, including IFN- γ , has been associated with resolution of food allergy and favorable outcomes for immunotherapy in humans (56). It has been suggested that IFN- γ can reduce allergic disease through suppression of Th2 cells as well as effects on the innate cells and alarmins which are required for both the induction and maintenance of allergic disease. Innate cytokines including the alarmins IL-25, IL-33, and TSLP are produced by epithelial cells and are key mediators of allergic disease (57). The NE vaccines in our studies suppressed alarmin expression in an IFN- γ dependent mechanism, and this correlated with reduction of allergic reactivity to bystander allergens. This suggests local NE-induced IFN- γ modulates the small intestine environment to suppress the allergic response. Since these components of the innate immune system function in an antigen-independent manner, this may be responsible for the non-antigen specific bystander effects that confer protection against allergens not included in the NE vaccine. Additional studies are required to determine if modulation of a specific alarmin is required for suppression of reactivity, and further dissection of the

mechanism of IFN- γ induction on suppression of reactivity is currently in progress.

A role for IFN- γ -mediated suppression of ILC2s in the small intestine cannot be ruled out, as mice that received the NE vaccines also had reduced ILC2s in the small intestine. Previous work has demonstrated that in the lung, IFN- γ prevents the accumulation of these lymphoid cells in mucosal tissues by limiting the recruitment and maintenance of these cells (58–60). The reduction in ILC2s in the mice that received the NE vaccines may also be a critical factor for the observed reduction in allergic reactivity, as IL-4 and IL-13 producing ILC2s have also been shown to promote experimental food allergy (36, 61).

Overall, these results show that modulation of Th2 immunity towards one food can induce bystander effects that suppress allergic reactivity to unrelated foods. This may lead to a global reduction of allergic reactivity to multiple foods. This allergen-non-specific protection may also be induced by other therapies that increase IFN- γ and decreased expression of alarmins in the gut mucosa. Taken together, these results suggest new targets for the suppression of allergic disease.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The animal studies were reviewed by the University of Michigan Institutional Animal Care and Use Committee.

AUTHOR CONTRIBUTIONS

MF, FF, JB, and JO designed the studies and prepared the manuscript. MF, JL, KJ, HL, and JO performed experiments and analyzed data. All authors contributed to the article and approved the submitted version.

FUNDING

This project has been funded by a Food Allergy Research and Education New Investigator Award, the National Institute for

Allergy and Infectious Disease, National Institutes of Health under Grant R01AI145991, the Michigan Food Allergy Research Accelerator (M-FARA), and a generous gift from Robert and Caren Vondell.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2021.599296/full#supplementary-material>

REFERENCES

- Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. *JAMA Pediatr* (2013) 167:1026–31. doi: 10.1001/jamapediatrics.2013.2376
- Shamji MH, Durham SR. Mechanisms of immunotherapy to aeroallergens. *Clin Exp Allergy* (2011) 41:1235–46. doi: 10.1111/j.1365-2222.2011.03804.x
- Oppenheimer JJ, Nelson HS, Bock SA, Christensen F, Leung DY. Treatment of peanut allergy with rush immunotherapy. *J Allergy Clin Immunol* (1992) 90:256–62. doi: 10.1016/0091-6749(92)90080-L
- Nelson HS, Lahr J, Rule R, Bock A, Leung D. Treatment of anaphylactic sensitivity to peanuts by immunotherapy with injections of aqueous peanut extract. *J Allergy Clin Immunol* (1997) 99:744–51. doi: 10.1016/S0091-6749(97)80006-1
- Palisade Group of Clinical Investigators, Vickery BP, Vereda A, Casale TB, Beyer K, du Toit G, et al. AR101 Oral Immunotherapy for Peanut Allergy. *N Engl J Med* (2018) 379:1991–2001. doi: 10.1056/NEJMoa1812856
- Jones SM, Sicherer SH, Burks AW, Leung DY, Lindblad RW, Dawson P, et al. Epicutaneous immunotherapy for the treatment of peanut allergy in children and young adults. *J Allergy Clin Immunol* (2017) 139:1242–52.e9. doi: 10.1016/j.jaci.2016.08.017
- Mondoulet L, Dioszeghy V, Vanoirbeek JA, Nemery B, Dupont C, Benhamou PH. Epicutaneous immunotherapy using a new epicutaneous delivery system in mice sensitized to peanuts. *Int Arch Allergy Immunol* (2011) 154:299–309. doi: 10.1159/000321822
- Jones SM, Burks AW, Dupont C. State of the art on food allergen immunotherapy: oral, sublingual, and epicutaneous. *J Allergy Clin Immunol* (2014) 133:318–23. doi: 10.1016/j.jaci.2013.12.1040
- Keet CA, Frischmeyer-Guerrero PA, Thyagarajan A, Schroeder JT, Hamilton RG, Boden S, et al. The safety and efficacy of sublingual and oral immunotherapy for milk allergy. *J Allergy Clin Immunol* (2012) 129:448–55.e1-5. doi: 10.1016/j.jaci.2011.10.023
- Burks AW, Jones SM, Wood RA, Fleischer DM, Sicherer SH, Lindblad RW, et al. Oral immunotherapy for treatment of egg allergy in children. *N Engl J Med* (2012) 367:233–43. doi: 10.1056/NEJMoa1200435
- Bauer RN, Manohar M, Singh AM, Jay DC, Nadeau KC. The future of biologics: applications for food allergy. *J Allergy Clin Immunol* (2015) 135:312–23. doi: 10.1016/j.jaci.2014.12.1908
- Galli SJ, Tsai M. IgE and mast cells in allergic disease. *Nat Med* (2012) 18:693–704. doi: 10.1038/nm.2755
- Akdis M, Akdis CA. Mechanisms of allergen-specific immunotherapy: multiple suppressor factors at work in immune tolerance to allergens. *J Allergy Clin Immunol* (2014) 133:621–31. doi: 10.1016/j.jaci.2013.12.1088
- Broide DH. Immunomodulation of Allergic Disease. *Annu Rev Med* (2009) 60:279–91. doi: 10.1146/annurev.med.60.041807.123524
- Casale T, Stokes J. Immunomodulators for allergic respiratory disorders. *J Allergy Clin Immunol* (2008) 121:288–96. doi: 10.1016/j.jaci.2007.11.040
- Bielinska AU, Janczak KW, Landers JJ, Makidon P, Sower LE, Peterson JW, et al. Mucosal immunization with a novel nanoemulsion-based recombinant anthrax protective antigen vaccine protects against *Bacillus anthracis* spore challenge. *Infect Immun* (2007) 75:4020–9. doi: 10.1128/IAI.00070-07
- Makidon PE, Bielinska AU, Nigavekar SS, Janczak KW, Knowlton J, Scott AJ, et al. Pre-clinical evaluation of a novel nanoemulsion-based hepatitis B mucosal vaccine. *PLoS One* (2008) 3:e2954. doi: 10.1371/journal.pone.0002954
- Passmore C, Makidon PE, O'Konek JJ, Zahn JA, Pannu J, Hamouda T, et al. Intranasal immunization with W 5EC adjuvanted recombinant RSV rF-ptn enhances clearance of respiratory syncytial virus in a mouse model. *Hum Vaccin Immunother* (2013) 10:615–22. doi: 10.4161/hv.27383
- Lindell DM, Morris SB, White MP, Kallal LE, Lundy PK, Hamouda T, et al. A novel inactivated intranasal respiratory syncytial virus vaccine promotes viral clearance without Th2 associated vaccine-enhanced disease. *PLoS One* (2011) 6:e21823. doi: 10.1371/journal.pone.0021823
- Hamouda T, Chepurinov A, Mank N, Knowlton J, Chepurnova T, Myc A, et al. Efficacy, immunogenicity and stability of a novel intranasal nanoemulsion-adjuvanted influenza vaccine in a murine model. *Hum Vaccin* (2010) 6:585–94. doi: 10.4161/hv.6.7.11818
- Bielinska AU, Chepurinov AA, Landers JJ, Janczak KW, Chepurnova TS, Luker GD, et al. A novel, killed-virus nasal vaccinia virus vaccine. *Clin Vaccine Immunol* (2008) 15:348–58. doi: 10.1128/CVI.00440-07
- Bielinska AU, Gerber M, Blanco LP, Makidon PE, Janczak KW, Beer M, et al. Induction of Th17 cellular immunity with a novel nanoemulsion adjuvant. *Crit Rev Immunol* (2010) 30:189–99. doi: 10.1615/CritRevImmunol.v30.i2.60
- Bielinska AU, Makidon PE, Janczak KW, Blanco LP, Swanson B, Smith DM, et al. Distinct pathways of humoral and cellular immunity induced with the mucosal administration of a nanoemulsion adjuvant. *J Immunol* (2014) 192:2722–33. doi: 10.4049/jimmunol.1301424
- Makidon PE, Belyakov IM, Blanco LP, Janczak KW, Landers J, Bielinska AU, et al. Nanoemulsion mucosal adjuvant uniquely activates cytokine production by nasal ciliated epithelium and induces dendritic cell trafficking. *Eur J Immunol* (2012) 42:2073–86. doi: 10.1002/eji.201142346
- O'Konek JJ, Landers JJ, Janczak KW, Goel RR, Mondrusov AM, Wong PT, et al. Nanoemulsion adjuvant-driven redirection of TH2 immunity inhibits allergic reactions in murine models of peanut allergy. *J Allergy Clin Immunol* (2018) 141:2121–31. doi: 10.1016/j.jaci.2018.01.042
- O'Konek JJ, Baker JR Jr. Treatment of allergic disease with nanoemulsion adjuvant vaccines. *Allergy* (2019) 74:246–9. doi: 10.1111/all.13977
- O'Konek JJ, Landers JJ, Janczak KW, Lindsey HK, Mondrusov AM, Totten TD, et al. Intranasal nanoemulsion vaccine confers long-lasting immunomodulation and sustained unresponsiveness in a murine model of milk allergy. *Allergy* (2020) 75:872–81. doi: 10.1111/all.14064
- Bielinska AU, O'Konek JJ, Janczak KW, Baker JR Jr. Immunomodulation of TH2 biased immunity with mucosal administration of nanoemulsion adjuvant. *Vaccine* (2016) 34:4017–24. doi: 10.1016/j.vaccine.2016.06.043
- Myc A, Kukowska-Latallo JF, Bielinska AU, Cao P, Myc PP, Janczak K, et al. Development of immune response that protects mice from viral pneumonitis after a single intranasal immunization with influenza A virus and nanoemulsion. *Vaccine* (2003) 21:3801–14. doi: 10.1016/S0264-410X(03)00381-5
- Ahrens R, Osterfeld H, Wu D, Chen CY, Arumugam M, Groschwitz K, et al. Intestinal mast cell levels control severity of oral antigen-induced anaphylaxis in mice. *Am J Pathol* (2012) 180:1535–46. doi: 10.1016/j.ajpath.2011.12.036
- Finkelman FD, Katona IM, Mosmann TR, Coffman RL. IFN-gamma regulates the isotypes of Ig secreted during in vivo humoral immune responses. *J Immunol* (1988) 140:1022–7.

32. Li XM, Serebrisky D, Lee SY, Huang CK, Bardina L, Schofield BH, et al. A murine model of peanut anaphylaxis: T- and B-cell responses to a major peanut allergen mimic human responses. *J Allergy Clin Immunol* (2000) 106:150–8. doi: 10.1067/mai.2000.107395
33. Rodriguez B, Prioult G, Hacini-Rachinel F, Moine D, Bruttin A, Ngom-Bru C, et al. Infant gut microbiota is protective against cow's milk allergy in mice despite immature ileal T-cell response. *FEMS Microbiol Ecol* (2012) 79:192–202. doi: 10.1111/j.1574-6941.2011.01207.x
34. Classen DC, Morningstar JM, Shanley JD. Detection of antibody to murine cytomegalovirus by enzyme-linked immunosorbent and indirect immunofluorescence assays. *J Clin Microbiol* (1987) 25:600–4. doi: 10.1128/JCM.25.4.600-604.1987
35. Frey A, Di Canzio J, Zurakowski D. A statistically defined endpoint titer determination method for immunoassays. *J Immunol Methods* (1998) 221:35–41. doi: 10.1016/S0022-1759(98)00170-7
36. Lee JB, Chen CY, Liu B, Mugge L, Angkasekwinai P, Facchinetti V, et al. IL-25 and CD4(+) TH2 cells enhance type 2 innate lymphoid cell-derived IL-13 production, which promotes IgE-mediated experimental food allergy. *J Allergy Clin Immunol* (2016) 137:1216–1225 e5. doi: 10.1016/j.jaci.2015.09.019
37. Oyoshi MK, Larson RP, Ziegler SF, Geha RS. Mechanical injury polarizes skin dendritic cells to elicit a T(H)2 response by inducing cutaneous thymic stromal lymphopoietin expression. *J Allergy Clin Immunol* (2010) 126:976–84. doi: 10.1016/j.jaci.2010.08.041
38. Allinne J, Scott G, Lim WK, Birchard D, Erjefalt JS, Sanden C, et al. IL-33 blockade affects mediators of persistence and exacerbation in a model of chronic airway inflammation. *J Allergy Clin Immunol* (2019) 144:1624–1637 e10. doi: 10.1016/j.jaci.2019.08.039
39. Muto T, Fukuoka A, Kabashima K, Ziegler SF, Nakanishi K, Matsushita K, et al. The role of basophils and proallergic cytokines, TSLP and IL-33, in cutaneously sensitized food allergy. *Int Immunol* (2014) 26:539–49. doi: 10.1093/intimm/dxu058
40. Hussain M, Borcard L, Walsh KP, Pena Rodriguez M, Mueller C, Kim BS, et al. Basophil-derived IL-4 promotes epicutaneous antigen sensitization concomitant with the development of food allergy. *J Allergy Clin Immunol* (2018) 141:223–234 e5. doi: 10.1016/j.jaci.2017.02.035
41. Stier MT, Zhang J, Goleniewska K, Cephus JY, Rusznak M, Wu L, et al. IL-33 promotes the egress of group 2 innate lymphoid cells from the bone marrow. *J Exp Med* (2018) 215:263–81. doi: 10.1084/jem.20170449
42. von Moltke J, Ji M, Liang HE, Locksley RM. Tuft-cell-derived IL-25 regulates an intestinal ILC2-epithelial response circuit. *Nature* (2016) 529:221–5. doi: 10.1038/nature16161
43. Huang Y, Guo L, Qiu J, Chen X, Hu-Li J, Siebenlist U, et al. IL-25-responsive, lineage-negative KLRG1(hi) cells are multipotential 'inflammatory' type 2 innate lymphoid cells. *Nat Immunol* (2015) 16:161–9. doi: 10.1038/ni.3078
44. Gupta RS, Springston EE, Warrier MR, Smith B, Kumar R, Pongracic J, et al. The prevalence, severity, and distribution of childhood food allergy in the United States. *Pediatrics* (2011) 128:e9–17. doi: 10.1542/peds.2011-0204
45. Gupta RS, Warren CM, Smith BM, Jiang J, Blumenstock JA, Davis MM, et al. Prevalence and Severity of Food Allergies Among US Adults. *JAMA Netw Open* (2019) 2:e185630. doi: 10.1001/jamanetworkopen.2018.5630
46. Begin P, Winterroth LC, Dominguez T, Wilson SP, Bacal L, Mehrotra A, et al. Safety and feasibility of oral immunotherapy to multiple allergens for food allergy. *Allergy Asthma Clin Immunol* (2014) 10:1. doi: 10.1186/1710-1492-10-1
47. Andorf S, Manohar M, Dominguez T, Block W, Tupa D, Kshirsagar RA, et al. Feasibility of sustained response through long-term dosing in food allergy immunotherapy. *Allergy Asthma Clin Immunol* (2017) 13:52. doi: 10.1186/s13223-017-0224-7
48. Wang LF, Chen JS, Hsu CJ, Liu CY, Yu JS, Miaw SC. Antigen-driven bystander effect accelerates epicutaneous sensitization with a new protein allergen. *J BioMed Sci* (2009) 16:28. doi: 10.1186/1423-0127-16-28
49. Eisenbarth SC, Zhadkevich A, Ranney P, Herrick CA, Bottomly K. IL-4-dependent Th2 collateral priming to inhaled antigens independent of Toll-like receptor 4 and myeloid differentiation factor 88. *J Immunol* (2004) 172:4527–34. doi: 10.4049/jimmunol.172.7.4527
50. Gregory JA, Shepley-McTaggart A, Umpierrez M, Hurlburt BK, Maleki SJ, Sampson HA, et al. Immunotherapy using algal-produced Ara h 1 core domain suppresses peanut allergy in mice. *Plant Biotechnol J* (2016) 14:1541–50. doi: 10.1111/pbi.12515
51. Storni F, Zeltins A, Balke I, Heath MD, Kramer MF, Skinner MA, et al. Vaccine against peanut allergy based on engineered virus-like particles displaying single major peanut allergens. *J Allergy Clin Immunol* (2020) 145:1240–53. doi: 10.1016/j.jaci.2019.12.007
52. Bannon GA, Cockrell G, Connaughton C, West CM, Helm R, Stanley JS, et al. Engineering, characterization and in vitro efficacy of the major peanut allergens for use in immunotherapy. *Int Arch Allergy Immunol* (2001) 124:70–2. doi: 10.1159/000053672
53. Mondoulet L, Dioszeghy V, Puteaux E, Ligouis M, Dhelft V, Plaquet C, et al. Specific epicutaneous immunotherapy prevents sensitization to new allergens in a murine model. *J Allergy Clin Immunol* (2015) 135:1546–57.e4. doi: 10.1016/j.jaci.2014.11.028
54. Mondoulet L, Dioszeghy V, Busato F, Plaquet C, Dhelft V, Bethune K, et al. Gata3 hypermethylation and Foxp3 hypomethylation are associated with sustained protection and bystander effect following epicutaneous immunotherapy in peanut-sensitized mice. *Allergy* (2019) 74:152–64. doi: 10.1111/all.13479
55. Dunkin D, Berin MC, Mondoulet L, Tobar S, Yeretssian G, Tordesillas L, et al. Epicutaneous Tolerance Induction to a Bystander Antigen Abrogates Colitis and Ileitis in Mice. *Inflammation Bowel Dis* (2017) 23:1972–82. doi: 10.1097/MIB.0000000000001273
56. Turcanu V, Maleki SJ, Lack G. Characterization of lymphocyte responses to peanuts in normal children, peanut-allergic children, and allergic children who acquired tolerance to peanuts. *J Clin Invest* (2003) 111:1065–72. doi: 10.1172/JCI200316142
57. Hammad H, Lambrecht BN. Barrier Epithelial Cells and the Control of Type 2 Immunity. *Immunity* (2015) 43:29–40. doi: 10.1016/j.immuni.2015.07.007
58. Molofsky AB, Van Gool F, Liang HE, Van Dyken SJ, Nussbaum JC, Lee J, et al. Interleukin-33 and Interferon-gamma Counter-Regulate Group 2 Innate Lymphoid Cell Activation during Immune Perturbation. *Immunity* (2015) 43:161–74. doi: 10.1016/j.immuni.2015.05.019
59. Li BWS, de Bruijn MJW, Lukkes M, van Nimwegen M, Bergen IM, Kleinjan A, et al. T cells and ILC2s are major effector cells in influenza-induced exacerbation of allergic airway inflammation in mice. *Eur J Immunol* (2019) 49:144–56. doi: 10.1002/eji.201747421
60. Starkey MR, McKenzie AN, Belz GT, Hansbro PM. Pulmonary group 2 innate lymphoid cells: surprises and challenges. *Mucosal Immunol* (2019) 12:299–311. doi: 10.1038/s41385-018-0130-4
61. Noval Rivas M, Burton OT, Oettgen HC, Chatila T. IL-4 production by group 2 innate lymphoid cells promotes food allergy by blocking regulatory T-cell function. *J Allergy Clin Immunol* (2016) 138:801–11.e9. doi: 10.1016/j.jaci.2016.02.030

Conflict of Interest: JB holds stock in Blue Willow Biologics, a company that has licensed the adjuvant technology from the University of Michigan. JB and JO'K are inventors of the adjuvant technology involved in this research and patent applications have been submitted for this technology.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Farazuddin, Landers, Janczak, Lindsey, Finkelman, Baker and O'Konek. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.