



Case Report: Interferon- γ Restores Monocytic Human Leukocyte Antigen Receptor (mHLA-DR) in Severe COVID-19 With Acquired Immunosuppression Syndrome

Steffen Dickel^{1*}, Clemens Grimm¹, Katharina Amschler², Sebastian Uwe Schnitzler¹, Julie Schanz³, Onnen Moerer¹, Didier Payen⁴, Bjoern Tampe⁵ and Martin Sebastian Winkler¹

OPEN ACCESS

Edited by:

Fabienne Venet,
Hospices Civils de Lyon, France

Reviewed by:

Julie Demaret,
Queen Mary University of London,
United Kingdom
Vijayakumar Velu,
Emory University, United States

*Correspondence:

Steffen Dickel
steffen.dickel@med.uni-goettingen.de

Specialty section:

This article was submitted to
Viral Immunology,
a section of the journal
Frontiers in Immunology

Received: 22 December 2020

Accepted: 18 March 2021

Published: 07 April 2021

Citation:

Dickel S, Grimm C, Amschler K,
Schnitzler SU, Schanz J, Moerer O,
Payen D, Tampe B and Winkler MS
(2021) Case Report: Interferon- γ
Restores Monocytic Human
Leukocyte Antigen Receptor
(mHLA-DR) in Severe
COVID-19 With Acquired
Immunosuppression Syndrome.
Front. Immunol. 12:645124.
doi: 10.3389/fimmu.2021.645124

¹ Department of Anesthesiology and Intensive Care Medicine, University Medical Center Goettingen, Goettingen, Germany, ² Department of Dermatology, University Medical Center Goettingen, Goettingen, Germany, ³ Institute of Clinical Chemistry and Laboratory Medicine, University Medical Center, Goettingen, Goettingen, Germany, ⁴ Universite Paris 7 Cite Sorbonne, UMR INSERM 1160, Paris, France, ⁵ Department of Nephrology and Rheumatology, University Medical Center, Goettingen, Goettingen, Germany

Background: The major histocompatibility complex (MHC) class II characterized by monocytes CD14+ expression of human leukocyte antigen receptors (HLA-DR), is essential for the synapse between innate and adaptive immune response in infectious disease. Its reduced expression is associated with a high risk of secondary infections in septic patients and can be safely corrected by Interferon- γ (IFN γ) injection. Coronavirus disease (COVID-19) induces an alteration of Interferon (IFN) genes expression potentially responsible for the observed low HLA-DR expression in circulating monocytes (mHLA-DR).

Methods: We report a case of one-time IFN γ injection (100 mcg s.c.) in a superinfected 61-year-old man with COVID-19-associated acute respiratory distress syndrome (ARDS), with monitoring of mHLA-DR expression and clinical tolerance.

Observations: Low mHLA-DR pretreatment expression (26.7%) was observed. IFN γ therapy leading to a rapid increase in mHLA-DR expression (83.1%).

Conclusions: Severe ARDS in a COVID-19 patient has a deep reduction in mHLA-DR expression concomitantly with secondary infections. The unique IFN γ injection was safe and led to a sharp increase in the expression of mHLA-DR. Based on immune and infection monitoring, more cases of severe COVID-19 patients with low mHLA-DR should be treated by IFN γ to test the clinical effectiveness.

Keywords: SARS-CoV-2, COVID-19, monocytic human leukocyte antigen receptor, interferon- γ , intensive care, acute respiratory distress syndrome, acquired immunosuppression syndrome, case report

INTRODUCTION

Among the reported clinical phenotypes after severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection, acute respiratory distress syndrome (ARDS) is the most frequent life-threatening context requiring treatment in intensive care unit (ICU) (1). The underlying molecular mechanisms of the SARS-CoV-2 induced host response remain partially unknown. However, it is clear that inflammation patterns can rapidly change over time (2, 3). We have recently shown that an early and sharp decrease in monocytic human leukocyte antigen receptors (HLA-DR) expression in peripheral blood was associated with changes in the proportion of monocyte subpopulations (2). Since the impairment of interferon (IFN) gene expression have been demonstrated, a deficit in interferons has been suspected with a potential treatment option with IFN molecules to reinforce immune defenses (4). This is the first case report of a patient with severe ARDS from COVID-19 suffering from multiple secondary infections, associated with a dramatic reduction in mHLA-DR expression on circulating monocytes. Even after several days of adequate pharmacologic antimicrobial treatment the patient displayed ongoing clinical deterioration, repetitive positive blood cultures and low expression of mHLA-DR. This led to the compassionate use of one dose of IFN γ to boost innate immunity and antigen presentation.

MEDICAL HISTORY

A 61-year-old man presented himself to his family physician two days after the onset of an irritable and dry cough. Preexisting medical conditions included cardiovascular risk factors, arterial hypertension, nicotine abuse and obesity. SARS-CoV-2 RNA was confirmed on day three after onset of symptoms. On this day he was admitted to a regional hospital ICU due to a continuous decline of his physical status. Severe hypoxia was first treated by

non-invasive, then invasive ventilation and subsequently by insertion of veno-venous extracorporeal membrane oxygenation (ECMO) on the twelfth day of the illness. The patient was then transferred to our ARDS center with the diagnosis of superinfected COVID-19 ARDS.

CLINICAL COURSE

Immediately after ICU admission the patient's condition worsened and tolerable blood gases (therapeutic goals: $\text{paO}_2 > 60$ mm Hg, $\text{SpO}_2 > 90\%$ and $\text{paCO}_2 40\text{--}50$ mm Hg) were only achieved under high-flow ECMO with 5.5 L/min blood- and 10 L/min of gas-flow ($\text{FiO}_2 1.0$). Several ventilation regimes were applied to optimize gas exchange under ECMO such as two-level ventilation with neuromuscular blockade and nitric oxide. To overcome this critical hypoxemia a secondary venous cannula was implanted on day 18 to increase ECMO blood-flow up to 7 L/min (Figure 1). The main clinical parameters are presented in Table 1. According to our standard operating procedure (SOP), the patient received intravenously (i.v.) dexamethasone for 10 days (6 mg per day). The diagnosis of a superinfected COVID-19 ARDS was made by the detection of Haemophilus influenzae, which was first treated with piperacillin/tazobactam for 6 days. Antibiotic treatment was switched to meropenem and ciprofloxacin after detection of a resistant Pseudomonas aeruginosa (3-MRGN) and a non-resistant Klebsiella oxytoca (Figure 1).

DIAGNOSTICS

Sequential chest X-ray examinations showed infiltration, congestion and severe pneumonia (Figure 2). Routine inflammatory markers such as leukocytes, C-reactive protein (CRP), procalcitonin (PCT) and interleukin-6 (IL-6) are presented in Table 1. We extended our immune monitoring to





TABLE 1 | Clinical findings and laboratory markers of the COVID-19 patient after ICU admission.

	After onset of first COVID-19 symptoms*									
	Normal range	Day 12*	Day 14	Day 16	Day 18	Day 20	Day 22	Day 25	Day 26	Day 29
Laboratory markers										
Leukocytes ($\times 10^3/\mu\text{l}$)	4.0–11.0	15.40	17.50	16.30	11.87	13.10	10.60	8.54	6.99	7.10
CRP (mg/L)	≤ 5	316.8	351.3	432.4	324.0	N/D	313.9	242.8	239.4	190.7
PCT ($\mu\text{g/L}$)	< 0.07	0.39	0.37	0.47	1.04	0.69	1.15	0.69	0.72	1.43
IL-6 (pg/ml)	< 7.0	164.3	146.8	472.6	238.6	262.5	285.9	187.2	72.9	120.4
Ventilation										
Horowitz-Quotient	> 300	127.5	97.8	116.9	56.0	65.0	69.0	80.0	57.0	74.6
ECMO										
Flow (l/min)		4.2	4.3	5.0	8.0	7.0	7.7	7.0	7.3	7.3
Clinical Finding										
SOFA-Score		10	11	11	12	16	15	15	16	15
Medication										
Norepinephrine in ($\mu\text{g/kg/min}$)		0.01	0.04	0.25	0.28	0.25	0.22	0.16	0.11	0.08

*Days after first COVID-19 symptoms, Admission to ICU (ARDS center) on day 12. SOFA, sepsis-related organ failure assessment score; N/D, not determined.



flow-cytometry analysis (FACS; measured on a Navios[®] system) of CD14+ monocytes and CD3+ lymphocytes, as shown in **Table 2**. Above all, we noticed a profound reduction in absolute number of monocytes that poorly expressed HLA-DR (CD14+ mHLA-DR+) within 22.4% to 28.4% on repetitive measurements, strongly suggesting a severe acquired immunodepression.

INTERVENTION

The extreme severity, the repetitive secondary infection, the dramatic reduction in mHLA-DR expression and the stagnant

clinical status after 16 days on our ICU prompted the off-label use of subcutaneously (s.c.) IFN γ in order to attenuate the acquired immunosuppression. We administered 100 mcg IFN γ on day 28 after written consent by the patient’s legal representative. Monocytic HLA-DR expression levels were measured the day after.

OUTCOMES/RESULTS

After a single dose of IFN γ the proportion of HLA-DR+/CD 14+ cells increased by 56.4% (**Table 2**). There was no impact on mHLA-DR expression in lymphocytes (CD 3+). Unfortunately,

TABLE 2 | Flow-cytometry (FACS) results.

Parameters	Day 16*	Day 18	Day 26	Day 29
Monocytes (cells/ μl)	378	203	125	111
CD14 + (cells/ μl)	325 (85.3%)	160 (78.9%)	97 (78.0%)	91 (82.0%)
CD14 + HLA-DR + (cells/ μl)	73 (22.4%)	45 (28.4)	26 (26.3%)	75 (82.2%)
Lymphocytes (cells/ μl)	1699	1210	1507	1181
CD3 + (cells/ μl)	875 (51.5%)	734 (60.7)	1014 (67.3%)	858 (72.6%)
HLA-DR + (cells/ μl)	126 (14.4%)	67 (9.1%)	87 (8.6%)	76 (8.9%)

*Days after first COVID-19 symptoms, admission to ICU (ARDS center) on day 12. SOFA, sepsis-related organ failure assessment score; N/D not determined; CD, Cluster of differentiation, HLA-DR, human leukocyte antigen receptors.

the patient died from refractory ARDS 29 days after the first symptoms of COVID-19.

DISCUSSION

The reported case of a patient with severe COVID-19 ARDS treated for 29 days in ICU matches the “typical” severe cases of COVID-19 pneumonia. The nonspecific inflammatory markers such as leukocytes or IL-6 are often moderately increased and do not necessarily correlate with the clinical severity, as observed in our case with low IL-6 levels and normal leukocytes (5, 6) (**Table 1**). The reported “hyperinflammation and cytokine storms” have also been discussed as a promoter of severe COVID-19 (7, 8), despite the big differences when compared to severe sepsis or septic shock (6). The consecutive overactivation of the adaptive immune system has been associated with lung failure and death (7–9). This background is one rationale for initial dexamethasone treatment, which had been also applied in our patient (10, 11). However, during the course of COVID-19 some patients develop signs of severe immunosuppression (8). This has been shown in longitudinal monitoring, when patients elicited markers of severe immunosuppression as observed in sepsis (2, 12, 13). Although not routinely incorporated in current sepsis care guidelines despite existence of functional cellular markers (13–15), the detection of an acquired immunosuppression syndrome (AIS) seems to be essential to tailor the immune therapy (16). Reduced mHLA-DR expression impairs the T-cells activation, which increases the likelihood of secondary infections by five times and is per-se associated with endotoxin tolerance (17, 18). Measurements beginning on day 17 of the illness revealed that merely 22% to 28% of CD14+ monocytes expressed HLA-DR. This supported our diagnosis of a severe AIS with high risk for secondary infection. The documented presence of opportunistic pathogens (*Haemophilus influenzae*, *Pseudomonas aeruginosa* and *Klebsiella oxytoca*) fits well with this concept. The timeline of our case corresponds to a recent longitudinal study of fifteen COVID-19 patients showing reduced mHLA-DR expression between days 11 and 14 (2). Taken together, these data support the hypothesis of a critical role of antigen presentation in COVID-19 and its potential impact on secondary infections and survival (2, 16, 19). In general, reduced HLA receptor status is an important risk factor for secondary infections in ICU (3, 19). Therefore mHLA-DR might present a comprehensive cellular and functional marker to detect the phase of highest vulnerability for secondary complications, as reported in COVID-19 (2). Interestingly, the bacterial superinfection occurred within this period in our patient. The unique dose of IFN γ was able to restore mHLA-DR expression by 56.4%. To the best of our knowledge this is the first case report testing the efficacy and tolerance of IFN γ therapy to overcome the COVID-19-induced AIS. Although this salvage therapy did not allow the patient to survive, it is important to note that circulating monocytes responded well after only one dose. IFN γ treatment was justified by compassionate use and based on several case reports. IFN γ treatment was done in

close consultation with the patient’s relative and was concurrent with the patient’s probable will. The patient’s further clinical deterioration most likely did not result from the administration of IFN γ . The safety of the drug has been demonstrated in various studies to date (20, 21). Known side effects include flu-like symptoms, fever, headaches, fatigue, diarrhea and rarely neutropenia and thrombocytopenia, which were not observed in this case after a single dose of IFN γ (20, 21). Consequently, it is probably reasonable to suspect that a single dose was insufficient and the delay to treatment too long to reverse the disease. Our case report describes one patient with an acquired immunosuppressive syndrome potentially induced by SARS-CoV-2, which was first identified at day 12 after COVID-19 symptoms. Reduced mHLA-DR is a well-described and reliable marker for sepsis induced immune paralysis and a risk factor for secondary complications (19, 22). Our single observation is in line with reports that COVID-19 patients present significant reduced mHLA-DR levels between days 11 and 14 of their critical illness, when compared to earlier stages of the disease (2). We have learned, that for ICU supportive care phenotyping individuals and monitoring the immune status to identify a “period at risk” for secondary complications seems essential. Treatment of COVID-19 within this period with an old molecules such as IFN γ might be one option to restore the COVID-19 sepsis-induced immune paralysis. Furthermore, immune monitoring is required since therapies, which have a broad impact on the immunity such as dexamethasone, are standard of care. Additionally, the consequences on other cells and cytokine levels will have to be investigated to ensure that IFN γ does not impair tissue recovery or induce specific side effects such as fever or leukopenia (23, 24).

CONCLUSION

Even though incompletely understood, the COVID-19 immune response moves along time evolution. After a few days, an AIS seems to be very frequent, which may facilitate the occurrence of secondary infections. This may prolong the ICU length of stay and may worsen the prognosis. Among the potential drugs shown to improve AIS, IFN γ has specific targets on innate cells, with modest or no effect on adaptive immunity. Although more information is needed, it seems reasonable to monitor the mHLA-DR in association with microbiological screening in order to optimize the timely application and the duration of treatment with IFN γ . Further studies will provide important information whether the application of IFN γ should play an extended role in the safe and efficient treatment of AIS in COVID-19.

PATIENT CONSENT

The case described does not contain any information on the identity of the patient.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are available from the corresponding author on reasonable request. Further inquiries can be directed to the corresponding author.

AUTHOR CONTRIBUTIONS

SD, CG, SS, OM, and MW treated the patient in ICU. SS and MW initiated the interferon therapy. JS performed the FACS analysis. SD, CG, DP, KA, BT, and MW analyzed the case and

the laboratory findings. SD, CG, DP, and MW wrote the manuscript. All authors read and accepted the final version of the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

The publication of the case report was supported by the open access publication fund of the Goettingen University Medical Center.

REFERENCES

- Hasan SS, Capstick T, Ahmed R, Kow CS, Mazhar F, Merchant H, et al. Mortality in covid-19 patients with acute respiratory distress syndrome and corticosteroids use: a systematic review and meta-analysis. *Expert Rev Res Med* (2020) 14:1149–63. doi: 10.1080/17476348.2020.1804365
- Payen D, Cravat M, Maadadi H, Didelot C, Prosic L, Dupuis C, et al. A longitudinal study of immune cells in severe covid-19 patients. *Front Immunol* (2020) 11:580250. doi: 10.3389/fimmu.2020.580250
- Pfortmueller CA, Meisel C, Fux M, Schefold JC. Assessment of immune organ dysfunction in critical illness: utility of innate immune response markers. *Intens Care Med* (2017) 5:49. doi: 10.1186/s40635-017-0163-0
- Payen D, Faivre V, Miatello J, Leentjens J, Brumpt C, Tissières P, et al. Multicentric experience with interferon gamma therapy in sepsis induced immunosuppression. a case series. *BMC Infect Dis* (2019) 19:931. doi: 10.1186/s12879-019-4526-x
- Ghahramani S, Tabrizi R, Lankarani KB, Kashani SMA, Rezaei S, Zeidi N, et al. Laboratory features of severe vs. non-severe covid-19 patients in asian populations: a systematic review and meta-analysis. *Eur J Med Res* (2020) 25:30. doi: 10.1186/s40001-020-00432-3
- Leisman DE, Ronner L, Pinotti R, Taylor MD, Sinha P, Calfee CS, et al. Cytokine elevation in severe and critical covid-19: a rapid systematic review, meta-analysis, and comparison with other inflammatory syndromes. *Lancet Respir Med* (2020) 8:1233–44. doi: 10.1016/s2213-2600(20)30404-5
- Ragab D, Salah Eldin H, Taimow M, Khattab R, Salem R. The covid-19 cytokine storm; what we know so far. *Front Immunol* (2020) 11:1446. doi: 10.3389/fimmu.2020.01446
- Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ. Covid-19: consider cytokine storm syndromes and immunosuppression. *Lancet* (2020) 395:1033–4. doi: 10.1016/s0140-6736(20)30628-0
- Pearce L, Davidson SM, Yellon DM. The cytokine storm of covid-19: a spotlight on prevention and protection. *Expert Opin Ther Tar* (2020) 24:723–30. doi: 10.1080/14728222.2020.1783243
- Sharun K, Tiwari R, Dhama J, Dhama K. Dexamethasone to combat cytokine storm in covid-19: Clinical trials and preliminary evidence. *Int J Surg* (2020) 82:179–81. doi: 10.1016/j.ijssu.2020.08.038
- Lammers T, Sofias AM, van der Meel R, Schifflers R, Storm G, Tacke F, et al. Dexamethasone nanomedicines for covid-19. *Nat Nanotechnol* (2020) 15:622–4. doi: 10.1038/s41565-020-0752-z
- Venet F, Lukaszewicz AC, Payen D, Hotchkiss R, Monneret G. Monitoring the immune response in sepsis: a rational approach to administration of immunoadjuvant therapies. *Curr Opin Immunol* (2013) 25:477–83. doi: 10.1016/j.coi.2013.05.006
- Long B, Koyfman A. Ready for prime time? biomarkers in sepsis. *Emerg Med Clin N Am* (2017) 35:109–22. doi: 10.1016/j.emc.2016.09.004
- Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving sepsis campaign: International guidelines for management of sepsis and septic shock: 2016. *Intens Care Med* (2017) 43:304–77. doi: 10.1007/s00134-017-4683-6
- Pierrakos C, Vincent JL. Sepsis biomarkers: a review. *Crit Care* (2010) 14:R15. doi: 10.1186/cc8872
- Spinetti T, Hirzel C, Fux M, Walti LN, Schober P, Stueber F, et al. Reduced monocytic human leukocyte antigen-dr expression indicates immunosuppression in critically ill covid-19 patients. *Anesth Analg* (2020) 131:993–9. doi: 10.1213/ANE.0000000000005044
- Lachmann G, von Haefen C, Kurth J, Yuerek F, Wernecke KD, Spies C. Smoking, gender, and overweight are important influencing factors on monocytic hla-dr before and after major cancer surgery. *BioMed Res Int* (2017) 2017:5216562. doi: 10.1155/2017/5216562
- Winkler MS, Rissiek A, Prießler M, Schwedhelm E, Robbe L, Bauer A, et al. Human leucocyte antigen (hla-dr) gene expression is reduced in sepsis and correlates with impaired tnf response: A diagnostic tool for immunosuppression? *PLoS One* (2017) 12:e0182427. doi: 10.1371/journal.pone.0182427
- Lukaszewicz AC, Grienay M, Resche-Rigon M, Pirracchio R, Faivre V, Boval B, et al. Monocytic hla-dr expression in intensive care patients: Interest for prognosis and secondary infection prediction *. *Crit Care Med* (2009) 37:2746–52. doi: 10.1097/00003246-200910000-00011
- Baldo BA. Side effects of cytokines approved for therapy. *Drug Safety* (2014) 37:921–43. doi: 10.1007/s40264-014-0226-z
- Razaghi A, Owens L, Heimann K. Review of the recombinant human interferon gamma as an immunotherapeutic: Impacts of production platforms and glycosylation. *J Biotechnol* (2016) 240:48–60. doi: 10.1016/j.jbiotec.2016.10.022
- Landelle C, Lepape A, Voirin N, Tognet E, Venet F, Bohé J, et al. Low monocyte human leukocyte antigen-dr is independently associated with nosocomial infections after septic shock. *Intensive Care Med* (2010) 36:1859–66. doi: 10.1007/s00134-010-1962-x
- Shen H, Yao P, Lee E, Greenhalgh D, Soulika AM. Interferon-gamma inhibits healing post scald burn injury. *Wound Repair Regen* (2012) 20:580–91. doi: 10.1111/j.1524-475X.2012.00812.x
- Miller CHT, Maher SG, Young HA. Clinical use of interferon-gamma. *Ann NY Acad Sci* (2009) 1182:69–79. doi: 10.1111/j.1749-6632.2009.05069.x

Conflict of Interest: MW received unrestricted funding from SARTORIUS Ag-Lung research.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Dickel, Grimm, Amschler, Schnitzler, Schanz, Moerer, Payen, Tampe and Winkler. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.