



# Corrigendum: Generation of Powerful Human Tolerogenic Dendritic Cells by Lentiviral-Mediated IL-10 Gene Transfer

## OPEN ACCESS

### Edited and reviewed by:

Julia Szekeres-Bartho,  
University of Pécs, Hungary

### \*Correspondence:

Silvia Gregori  
gregori.silvia@hsr.it

### Specialty section:

This article was submitted to  
Immunological Tolerance  
and Regulation,  
a section of the journal  
Frontiers in Immunology

**Received:** 26 February 2021

**Accepted:** 08 March 2021

**Published:** 22 March 2021

### Citation:

Comi M, Amodio G, Passeri L,  
Fortunato M, Santoni de Sio FR,  
Andolfi G, Kajaste-Rudnitski A,  
Russo F, Cesana L and Gregori S  
(2021) Corrigendum: Generation  
of Powerful Human Tolerogenic  
Dendritic Cells by Lentiviral-  
Mediated IL-10 Gene Transfer.  
*Front. Immunol.* 12:672701.  
doi: 10.3389/fimmu.2021.672701

Michela Comi, Giada Amodio, Laura Passeri, Marta Fortunato,  
Francesca Romana Santoni de Sio, Grazia Andolfi, Anna Kajaste-Rudnitski,  
Fabio Russo, Luca Cesana and Silvia Gregori\*

San Raffaele Telethon Institute for Gene Therapy (SR-TIGET), San Raffaele Scientific Institute (IRCCS), Milan, Italy

**Keywords:** dendritic cells, IL-10, cell therapy, immune tolerance, allogeneic transplantation

## A Corrigendum on


### Generation of Powerful Human Tolerogenic Dendritic Cells by Lentiviral-Mediated IL-10 Gene Transfer

By Comi M, Amodio G, Passeri L, Fortunato M, Santoni de Sio FR, Andolfi G, Kajaste-Rudnitski A, Russo F, Cesana L and Gregori S (2020). *Front. Immunol.* 11:1260. doi: 10.3389/fimmu.2020.01260.

In the original article, there was a mistake: a duplication of one dot plot in **Figure 8B** as published. The corrected **Figure 8** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Comi, Amodio, Passeri, Fortunato, Santoni de Sio, Andolfi, Kajaste-Rudnitski, Russo, Cesana and Gregori. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



**FIGURE 8** | *In vitro* characterization of murine DC<sup>IL-10</sup>. Female Balb/c BM cells were differentiated into DC, transduced at day 2 with LV-GFP (DC<sup>GFP</sup>) or LV-IL-10 (DC<sup>IL-10</sup>), and activated with LPS (200 ng/ml) during the last 2 days of differentiation. **(A)** Transduction efficiency was quantified based on ΔNGFR expression and the expression of the indicated markers was analyzed at day 9 of differentiation by flow cytometry. Each dot represents a single experiment (n = 8-9), lines indicate median, while whiskers are minimum and maximum levels. **(B)** DC were plated and left unstimulated or stimulated with LPS (200 ng/ml) for 24 h, with the addition of brefeldin A at 6 h. The expression of human IL-10 was quantified by intracytoplasmic staining. One representative donor out of two is depicted, and percentages of positive cells are indicated. **(C)** Spleen cells from female C57Bl/6 mice were stained with a proliferation dye and stimulated with Balb/c DC<sup>GFP</sup> and DC<sup>IL-10</sup> at 1:10 ratio. At day 5, proliferation of CD4<sup>+</sup> and CD8<sup>+</sup> T cells was measured by flow cytometry. Each dot represents a single donor (n = 6), lines indicate median, while whiskers are minimum and maximum levels. \*P ≤ 0.05 (Wilcoxon matched pairs test, two-tailed).