



CD47 Deficiency Ameliorates Ocular Autoimmune Inflammation

Yoko Okunuki, Steven J. Tabor, May Y. Lee and Kip M. Connor*

Angiogenesis Laboratory, Department of Ophthalmology, Massachusetts Eye and Ear, Harvard Medical School, Boston, MA, United States

OPEN ACCESS

Edited by:

Kutty Selva Nandakumar,
Southern Medical University, China

Reviewed by:

Andrew W. Taylor,
Boston University, United States

Wenru Su,
Sun Yat-Sen University, China

Thomas Langmann,
Universität zu Köln, Germany

John Vincent Forrester,
University of Aberdeen,
United Kingdom

*Correspondence:

Kip M. Connor
kip_connor@meei.harvard.edu

Specialty section:

This article was submitted to
Autoimmune and
Autoinflammatory Disorders,
a section of the journal
Frontiers in Immunology

Received: 14 March 2021

Accepted: 29 April 2021

Published: 20 May 2021

Citation:

Okunuki Y, Tabor SJ, Lee MY and
Connor KM (2021) CD47
Deficiency Ameliorates Ocular
Autoimmune Inflammation.
Front. Immunol. 12:680568.
doi: 10.3389/fimmu.2021.680568

Autoimmune uveitis is a sight-threatening ocular inflammatory condition in which the retina and uveal tissues become a target of autoreactive immune cells. The CD47 is a ubiquitously expressed transmembrane protein which plays multiple roles in fundamental cellular functions including phagocytosis, proliferation, and adhesion. Signal regulatory protein alpha (SIRP α), one of the CD47 ligands, is predominantly expressed in myeloid lineage cells such as dendritic cells (DCs) or macrophages, and CD47-SIRP α signaling pathway is implicated in the development of autoimmune diseases. Our current study demonstrates how CD47 depletion is effective in the prevention of experimental autoimmune uveitis (EAU), an animal model of human autoimmune uveitis, in animals deficient of CD47 (CD47^{-/-}). Systemic suppression of SIRP α ⁺ DCs in animals deficient in CD47 resulted in the inability of autoreactive CD4⁺ T cells to develop, which is crucial to induction of EAU. Of interest, retinal microglia, the resident immune cell of the retina, express SIRP α , however these cells were not operative in EAU suppression in response to CD47 depletion. These results identify CD47 as a significant regulator in the development of SIRP α ⁺ DCs that is vital to disease induction in EAU.

Keywords: uveitis, retina, autoimmune diseases, CD47 antigen, antigen-presenting cells

INTRODUCTION

Autoimmune uveitis, which occurs in a variety of diseases, including Behcet's disease, sarcoidosis, and Vogt-Koyanagi-Harada disease, among many others, is a sight-threatening ocular inflammatory disease (1–4). Although autoimmune uveitis covers a range of different clinical entities, autoimmunity against the retina and the uveal tissues is thought to be fundamental to its pathogenesis (5).

Experimental autoimmune uveitis (EAU) is an animal model of human autoimmune uveitis, and is widely used to delineate the pathophysiological processes of ocular autoimmunity and develop new approaches to treat patients (6). EAU is induced by immunization against retinal antigens such as interphotoreceptor retinoid-binding protein (IRBP), which is a major component of photoreceptor outer segments. IRBP immunization thus targets the immune response to photoreceptors (7, 8).

Uveitogenic antigen-specific CD4⁺ T-cells are crucial effector cells that drive inflammation and tissue damage after being activated by dendritic cells (DCs) (9). Systemic immunization with IRBP

and additional adjuvants leads to the antigen uptake by systemic DCs which in response activate antigen specific CD4⁺ T cells by presenting the antigen *via* MHC class II (MHC-II) in the peripheral lymphoid tissues.

The activated antigen specific CD4⁺ T cells proliferate and invade the retina following the breakdown of blood retinal barrier. Our group have shown that microglia facilitate autoreactive CD4⁺ T cell extravasation into the retina concurrent with other activated immune cell populations such as antigen presenting cells (APCs) (10). These immune cells are re-activated by cross-reactive retinal autoantigens and subsequently substantial inflammation ensues (11). There are multiple receptors, chemokines, and cytokines suggested to be involved in the development of autoimmune uveitis. Recently CD47-signal regulatory protein alpha (SIRP α) interactions have been implicated in induction of T cell-mediated inflammation (12, 13), and thus we sought to assess the role of this signaling axis in a pre-clinical model of uveitis that is largely driven by T-cell activation.

CD47, also known as integrin-associated protein (IAP), is an immunoglobulin-like protein ubiquitously expressed on the surface of virtually all cell types including erythrocytes and immune cells (14, 15). CD47 exerts pleiotropic functions by interacting with its ligands including SIRP α , thrombospondin-1 (TSP-1), and various integrins (16). Unlike the ubiquitous expression of CD47, SIRP α is expressed on limited number of cell populations including myeloid cells (monocytes, macrophages, DCs, and granulocytes) and neurons (17). Engagement of CD47 with SIRP α ⁺ phagocytes, such as DCs and macrophages, provides a down regulatory signal that inhibits host cell phagocytosis by SIRP α ⁺ phagocytes (18). Therefore, CD47 functions as a marker of self and is called the “don’t-eat-me” signal (19). This regulatory mechanism of phagocytosis initiated by CD47-SIRP α ⁺ interaction is well studied in red blood cell elimination by splenic macrophages (18) and in cancer immunity (19).

However, evidence from recent studies suggests that CD47-SIRP α interactions are key for efficient induction of T cell-mediated inflammation (20–22). One mechanism by which this is thought to occur is mediated through the interaction between SIRP α on DCs and CD47 on T cells to promote antigen presentation and T cell proliferation (23). The CD47-SIRP α interaction appears to activate and direct the migration of SIRP α ⁺ cells including neutrophils, melanoma cells and monocytes (24–26). Interestingly, it has been reported that SIRP α ⁺ monocytes interact with CD47 on brain endothelial cells to facilitate their transmigration across the blood brain barrier (26). Similarly, in an *in vivo* study of inflammatory bowel disease, it was demonstrated that interaction between SIRP α ⁺ DCs and CD47⁺ T cells is required for DCs to exert their migration and antigen presentation abilities (12). Therefore, it is likely that CD47-SIRP α interaction contributes to antigen presentation and T cell activation in the induction of a T cell mediated inflammatory diseases. Thus, this CD47-SIRP α interaction between DCs and CD4⁺ T cells has the potential to be an important mechanism in the pathogenesis of autoimmune diseases. Here we demonstrate that systemic expression of CD47

is essential in induction of autoimmune uveitis through promotion of development of SIRP α ⁺ DCs which has a crucial role in the priming of autoreactive CD4⁺ T cells.

MATERIALS AND METHODS

Animals and Reagents

Female animals were used for all of the experiments otherwise mentioned. All animal experiments followed the guidelines of the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research and were approved by the Animal Care Committee of the Massachusetts Eye and Ear Infirmary. C57BL/6J mice (stock no. 00664) were purchased from Jackson Laboratories (Bar Harbor, ME). Standard laboratory chow was fed. Mice were allowed free access to food and water in a climate-controlled room with a 12-hour light/12-hour dark cycle. All mice used for experiments were 7–10 weeks old. For anesthesia, intraperitoneal injection of 250 mg/kg of 2,2,2-tribromoethanol (Sigma-Aldrich Corp., St. Louis, MO) was used for survival procedures, and 400 mg/kg was used for non-survival procedures. High performance liquid chromatography-purified human interphotoreceptor retinoid binding protein peptide 1-20 (IRBP-p) was purchased from Biomatik (Wilmington, DE). Complete Freund’s Adjuvant and *Mycobacterium tuberculosis* H37Ra were purchased from Difco (Detroit, MI).

Induction of EAU

For active induction of EAU, 200 μ g of IRBP-p 1-20 was emulsified in Complete Freund’s Adjuvant (1:1 w/v) containing additional 5 mg/ml *M. tuberculosis* H37Ra. On day 0, 200 μ l of the emulsion was injected subcutaneously in the neck (100 μ l), one footpad (50 μ l) and the contralateral inguinal region (50 μ l). Concurrent with immunization, 1 μ g of purified Bordetella pertussis toxin (Sigma-Aldrich) was injected intraperitoneally.

Adoptive transfer EAU was induced as previously described (10). Briefly, donor mice were immunized as described above, and SPs and draining lymph nodes (LNs) were collected at 14 days after immunization. Lymphocytes from spleens (SPs) and draining LNs were cultured in the presence of 10 μ g/ml IRBP-p and 10 ng/ml IL-23 (R&D systems, Minneapolis, MN) for 72 hours in RPMI 1640 supplemented with 10% FBS, 10 mM HEPES, 1 x nonessential amino acid, and 1 mM sodium pyruvate (ThermoFisher, Waltham, MA), and 50 μ M 2-mercaptoethanol, 100 U/ml penicillin and 100 μ g/ml streptomycin (Sigma-Aldrich). IL-23 was added in the culture in order to induce effector T cells efficiently (27). The non-adherent cells in suspension were transferred to new dishes on day one and two of culture. After 3 days, activated lymphocytes were purified by gradient centrifugation on Histopaque 1083 (Sigma-Aldrich) and counted. The cells were injected intraperitoneally in 0.2 ml of PBS into recipient mice (4×10^7 cells/mouse).

Assessment of EAU

Fundus images were observed using a Micron IV retinal imaging microscope (Phoenix, Pleasanton, CA) and the clinical score of active inflammation was graded in a blinded manner on a scale

between 0-4 in half-point increments as described previously (28) with brief modification. Briefly, trace chorioretinal lesions and minimal vasculitis were score as 0.5. Mild vasculitis with small focal chorioretinal lesions (≤ 5) were scored as 1. Severe vasculitis with multiple chorioretinal lesions (>5) were scored as 2. Pattern of linear chorioretinal lesion, subretinal neovascularization and hemorrhages were scored as 3. Inflammation with large retinal detachment and severe hemorrhages were score as 4. For histological assessment, enucleated eyes were fixed in a buffer of 70% methanol and 30% acetic acid. The fixed tissues were embedded in paraffin and processed. Sections of 5 μm were cut and stained with hematoxylin and eosin (H&E). The severity of EAU in each eye was scored on a scale between 0-4 in half-point increments in a blinded manner as described previously (29). Briefly, focal non-granulomatous, monocytic infiltration in the choroid, ciliary body and retina were scored as 0.5. Retinal perivascular infiltration and monocytic infiltration in the vitreous were scored as 1. Granuloma formation in the uvea and retina, the presence of occluded retinal vasculitis, along with photoreceptor folds, serous retinal detachment and loss of photoreceptor were scored as 2. In addition, the formation of granulomas at the level of retinal pigment epithelium and the development of subretinal neovascularization were scored as 3 and 4 according to the number and the size of the lesions (30). The average of scores from both eyes was determined as the score of the animal.

Delayed Hypersensitivity Measurement

Antigen-specific delayed hypersensitivity was measured as previously described (31). On day 19 after immunization, mice were injected intradermally with 10 μg of IRBP-p suspended in 10 μl PBS into the pinna of one ear. Ear swelling was measured after 48 hours using a micrometer (Mitutoyo, Tokyo, Japan). Delayed hypersensitivity was measured as the difference in ear thickness before and after challenge. Results were expressed as: specific ear swelling = (48-hour measurement - 0-hour measurement) for test ear - (48-hour measurement - 0-hour measurement) for control ear.

Flow Cytometric Analysis of Lymph Nodes and Spleens

Cervical, axillary, and inguinal LNs and SPs were harvested from naive mice and EAU mice. Single cell suspensions were blocked with anti-mouse CD16/32 mAb (ThermoFisher). Dead cells were stained with LIVE/DEADTM fixable dead cell stain kit (blue or violet) (ThermoFisher) or Zombie Fixable Viability Kit (UV or NIR) (Biolegend). The following anti-mouse antibodies were used for staining: CD4 (clone: GK1.5), CD8 (53-6.7), CD19 (6D5), CD11c (N418), CD11b (M1/70), CD45 (30-F11), CD3 (17A2), CD44 (IM7), CD62L (MEL-14), CD25 (PC61.5), CD80 (16-10A1), CD86 (GL-1), MHC-II (M5/114.15.2), NK1.1 (PK136), and Ly6G (1A8) were purchase from BioLegend (San Diego, CA). CD11c-BB515 (N418) and SIRP α (P84) was purchased from BD Bioscience. Flow cytometric data were acquired on a CytoFlex S (Beckman Coulter, Brea, CA). Single

color staining and fluorescence minus one (FMO) controls were used for accurate compensation and gating. Acquired data was analyzed using FlowJo 10.5.3 (Ashland, OR).

Flow Cytometric Analysis of Retinal Microglia

For flowcytometric analysis of retinal microglia, mice were first perfused with PBS, and then the eyes were enucleated and the retinas were collected. Both retinas of each animal were combined and incubated for 45 min at 37°C in HBSS containing 10% FBS, 10 mM HEPES, 0.7 mg/ml calcium chloride (Sigma-Aldrich), 1 mg/ml collagenase D, and 0.1 mg/ml DNase I (Roche). After the incubation, retinas were homogenized by gentle trituration by pipetting and transferred to a FACS tube by passing through a 40 μm cell strainer. The samples were centrifuged at 350 rcf for 5 min, re-suspended in FACS buffer, and further centrifuged at 50 \times g for 1 min to precipitate debris (32). Supernatant containing cells was collected and used for subsequent staining after washing with FACS buffer a couple of times. The cell suspensions were blocked with anti-mouse CD16/32 mAb. Dead cells were stained with DAPI and excluded from the analysis. Microglia were identified as a CD11b⁺CD45^{low} population and SIRP α positive cells were evaluated.

Lymphocyte Proliferation Assay

Cells were isolated from draining LNs and SPs, and suspended at 2×10^5 per 200 μL medium in 96-well flat-bottom plates. The cells were cultured in triplicate in the presence of various concentration of IRBP-p or medium alone. The last four hours of 72 hours culture, 100 μl of supernatant in the culture medium was removed and 10 μl of Cell Counting Kit-8 (Sigma-Aldrich) was added each well and the cells were incubated for 4 hours. At the end of culture, the viable cell numbers in each well were measured as the absorbance (450 nm) of reduced WST-8 (33).

For mixed lymphocyte proliferation assay of CD11c⁺ DC priming of the CD4⁺ T cell, first CD4⁺ T cells were isolated from LNs by MACS system (Miltenyi Biotec, Bergisch Gladbach, Germany) using CD4⁺ T cell isolation kit. CD11c⁺ DCs were isolated from SPs using CD11c MicroBeads UltraPure. Isolated CD4⁺ T cells were added to a solution containing CellTraceTM Far Red (CTFR)(ThermoFisher) in PBS at 37°C for 20 min following the manufacture instruction (34). Aliquots of CTFR-labeled CD4⁺ T cells (2×10^5) in U-bottom 96-well plates were co-cultured with isolated CD11c⁺ DC (1×10^5) in 200 μl /well in the presence of 10 $\mu\text{g}/\text{ml}$ IRBP-p for 3 days. The frequency and number of proliferated CD4⁺ T cell in each sample was determined by diluted CTFR gated on live CD3⁺CD4⁺ cells.

Immunohistochemistry of Whole-Mount Retinas

Anesthetized mice were perfused with 20 mL of 1X PBS and eyes were enucleated. Whole eyes were fixed in 4% paraformaldehyde in 2X PBS for 45 minutes and then transferred to 2X PBS on ice for 15 minutes. Following the removal of the lens, retinas were isolated and prepared with four radical incisions. Retinas were

subsequently transferred to ice cold methanol and stored for further use at -80°C . For immunohistochemistry, retinas were washed in 1X PBS three times and blocked overnight in blocking buffer (5% goat serum, 0.03% triton-X100, 0.2% BSA in 1X PBS). Retinas were briefly washed in 1X PBS and then incubated with primary antibodies: rabbit anti-P2ry12 (1:500; a gift from H. Weiner, Brigham and Women's Hospital) and rat anti-SIRP α (1:500 Clone: P84 from Biolegend) in blocking buffer overnight at 4°C . Retinas were then washed four times in 1X PBS. Retinas were incubated with secondary antibodies goat anti-rabbit 594 (1:500), and goat anti-rat 647 (1:500) from ThermoFisher in blocking buffer overnight at 4°C . Whole-mounts were then washed four times in 1X PBS and mounted using anti-fade medium (Permaflour, Thermo Fisher Scientific, Waltham, MA). Images were taken on a Lecia Confocal Sp8.

Quantification and Statistical Analysis

Data are presented as mean \pm standard error of the mean (SEM). Differences between two groups were analyzed using an unpaired Student's *t*-test or Mann-Whitney test. Multiple-group comparison was performed by one-way ANOVA followed by Tukey's multiple comparison test. All statistical analyses were performed using graphing software (Prism 6, GraphPad Software, Inc., La Jolla, CA). Significance levels are marked * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$ in figures.

RESULTS

Loss of CD47 Suppresses Autoimmune Uveitis

To evaluate the overall contribution of CD47 in development of autoimmune uveitis, we assessed disease severity in mice with a systemic deficiency of CD47. CD47 knockout ($CD47^{-/-}$) mice were utilized and EAU was induced by immunizing the animals with IRBP-peptide 1–20 (IRBP-p) in both knockout animals and littermate wild type (WT) controls. Ocular inflammation associated with EAU was observed from day 7 to day 28 after immunization. Clinical assessment of these animals by fundoscopic examination found that EAU was significantly suppressed in female $CD47^{-/-}$ mice throughout the course of disease, whereas the control animals developed significant retinal inflammation (Figures 1A, B). Similarly, $CD47^{-/-}$ male mice were assessed and found to exhibit an equivalent degree of EAU suppression as their female counterparts (Supplemental Figure S1). Female animals had been conventionally used in studies of EAU (7, 30, 35–40), however there did not seem to be an obvious difference in susceptibility to EAU between males and females (29, 41). Since male $CD47^{-/-}$ mice exhibited the same degree of disease inhibition as females, female mice were utilized for the remainder of the study. Histological evaluation of EAU disease severity at the peak of inflammation in this model, day 21, further confirmed that CD47 deficiency significantly suppressed infiltrating inflammatory cells and retinal damage (Figures 1C, D). Although autoimmune uveitis is a condition observed locally in the eye, systemic immune activation against the

retinal auto-antigen leads to ocular inflammation. Therefore, we next evaluated the systemic immune response against the immunized peptide by (i) IRBP-specific delayed hyper-sensitivity *in vivo*, and (ii) IRBP-specific and nonspecific (Concanavalin A-induced) lymphocyte proliferation *in vitro*. We found that the systemic IRBP-specific immune response was suppressed in $CD47^{-/-}$ mice both *in vivo* and *in vitro*. The induction of delayed hypersensitivity in the ear against IRBP-p was significantly reduced in $CD47^{-/-}$ mice (Figure 1E) compared to control animals. Moreover, IRBP-specific and non-specific *in vitro* lymphocyte proliferation using lymph node (LN) cells and spleen (SP) cells, isolated from mice with EAU at 14 days (Supplemental Figure S2) and 21 days (Figures 1F, G) post immunization was significantly reduced in $CD47^{-/-}$ mice. These results indicate that the systemic immune response against the immunized peptide was suppressed in $CD47^{-/-}$ mice and that the suppressive effects of EAU observed in $CD47^{-/-}$ mice involve the suppression of systemic immune cell activation in response to IRBP-p autoantigen, not locally in the retina.

Suppression of EAU in CD47 Deficient Mice Was Not Regulated Within the Retinal Microenvironment

EAU was suppressed in animals that lack systemic expression of CD47 and our data assessing the delayed hypersensitivity response and immune cell proliferation against the antigen strongly implicated the suppression of systemic immunity as the mechanism of EAU inhibition. However, CD47 as well as its ligands, especially TSP-1, are expressed within the retina (42). In addition, we identified that microglia, the resident immune cells in the retina, express SIRP α under normal conditions (Supplemental Figure S3) and that microglial expression of SIRP α in $CD47^{-/-}$ mice was higher than in WT mice, although total microglial numbers in the retina was not largely affected by the loss of CD47 (Figure 2). It has been demonstrated that microglia are actively involved in the development of EAU especially in the early stage of inflammation (10). Therefore, it was possible that locally expressed CD47 and its ligands play a direct role in the disease course of EAU. To examine this possibility, we utilized the adoptive transfer model of EAU.

In this model, IRBP-reactive immune cells were transferred from donor immunized mice into naive recipient mice, subsequently donor cells induced EAU in recipient animals without additional stimulation. Here, primed cells from IRBP-immunized WT donor mice were adoptively transferred into naive WT or naive $CD47^{-/-}$ recipient mice. This approach excluded any potential effects of systemic immune suppression in $CD47^{-/-}$ recipient mice on the systemic cell priming stage and will allow us to elucidate the involvement of the retinal specific immune response in recipient mice. We found that significant inflammation in both recipient groups (naive WT and naive $CD47^{-/-}$) with a comparable severity of disease induction (Figures 3A, B), indicating that expression of CD47 on primed circulating immune cells significantly affects the disease course of EAU. Strengthening this finding was the converse experiment, when primed cells from $CD47^{-/-}$ mice were transferred to naive





FIGURE 1 | CD47 deficient mice are protected from autoimmune uveitis. Female WT and *CD47*^{-/-} mice were immunized with IRBP-p and the animals were evaluated for (A–D) severity of retinal inflammation and (E–G) systemic immune responses against the IRBP-p. (A) Time course of EAU clinical scores were evaluated by fundus observation (n = 10–11 mice per group). (B) Representative retinal fundus images on day 21. (C) Histopathological EAU score was evaluated on day 21 (n = 6 mice per group). (D) Representative histopathological (H&E staining) images (Scale bars: 200 μm). (A, C) Data were analyzed by Mann–Whitney *U* test. (E) Delayed hypersensitivity, as determined by ear swelling, was evaluated on day 21 (n = 10–11). (F, G) Cell proliferation was evaluated by using the cells isolated from lymph nodes and spleens on day 21. The cells were cultured in triplicate for 3 days in the presence of IRBP-p (10 μg/ml), Con A (1 μg/ml), or medium only (n = 5 mice per group). (E–G) Data were analyzed by Student's *t* test. Data are expressed as mean ± SEM. **P* < 0.05; ***P* < 0.01; ****P* < 0.001.

WT mice, the recipient animals did not develop EAU (Figures 3C, D). In this approach, the systemic immune response of donor mice (WT or *CD47*^{-/-} mice) defined the severity of EAU in the recipient mice. Therefore, these results further indicate that systemic immune suppression is the fundamental mechanism of decreased EAU severity in *CD47*^{-/-} mice.

Loss of CD47 Decreases Systemic CD11b⁺ Populations

To begin to identify the key mechanism by which EAU is suppressed from the loss of CD47, we examined how specific

immune cell populations in peripheral lymphoid organs were affected due to the loss of CD47. We first examined major immune cell populations known to be involved in the disease pathogenesis of EAU (CD4⁺ T cells, CD8⁺ T cells, CD19⁺ B cells, CD11b⁺ macrophages, CD11c⁺ DCs) in SPs and LNs by flow cytometry, before (naïve) and after EAU induction (days 14 and 21). *CD47*^{-/-} mice had fewer cell numbers in SPs and LNs than WT mice before and after EAU (Figure 4A). Further analysis of each specific population revealed that CD4⁺ and CD8⁺ T cells were mildly decreased in *CD47*^{-/-} mice, whereas the number of CD19⁺ B cells remained unchanged. On the contrary, CD11b⁺ cells in both SPs



and LNs and CD11c⁺ cells in LNs were significantly decreased during EAU in $CD47^{-/-}$ mice (**Figure 4B**, **Supplemental Figure S4**). This indicated that macrophages (CD11b⁺) and DCs (CD11c⁺) are directly regulated by CD47 during EAU and that T cells are regulated to a lesser extent.

CD47 is expressed on homeostatic cells and activates the SIRP α receptor on phagocytes (macrophages and DCs) to suppress engulfment. It has been reported that the lack of CD47 caused diminished expression of its ligand SIRP α and subsequently affects the development of SIRP α ⁺ DC cell populations (22, 43). CD11c^{high} DCs are conventional DCs (cDCs), which can be further classified into two subpopulations according to the expression profile of CD8a and SIRP α ; CD8a⁺SIRP α ⁻ cDCs and CD8a⁻SIRP α ⁺ cDCs (44, 45). Consequently, we sought to examine SIRP α expression on phagocytes in our EAU model. Macrophage and DC populations (CD11b⁺ and/or CD11c⁺) in SP cells from 14 days post immunization were analyzed by flow cytometry. We found that the cells with high SIRP α expression were largely found within the CD11b⁺ population regardless of CD11c expression, whereas CD8a⁺ cells were mainly found within CD11c⁺ populations with lower CD11b expression (**Figures 5A, B**). Thus, CD11b⁺ DCs (**Figure 5A**, gate a) mainly consist of CD8a⁻SIRP α ⁺ cDCs, while CD11b^{low} DCs (**Figure 5A**, gate c) were mainly CD8a⁺SIRP α ⁻ cDCs.

Looking more specifically at SIRP α ⁺ populations, SIRP α expression levels in CD11b⁺ cells were decreased in $CD47^{-/-}$ mice compared to WT mice regardless of CD11c expression: SIRP α

expression in CD11b⁺ DCs and CD11b⁺ macrophages (**Figure 5A**, gate b), was lower in $CD47^{-/-}$ mice (**Figure 5C**). In addition, the cell number and frequency of CD11b⁺ DCs and CD11b⁺ macrophages were decreased in $CD47^{-/-}$ mice (**Figures 5D, E**). On the other hand, SIRP α expression as well as the frequency and the number of CD11b^{low} DCs (**Figure 5A**, gate c) and CD11b⁻ DCs (**Figure 5A**, gate d), was maintained or rather increased in $CD47^{-/-}$ mice (**Figures 5C, D**). These results indicate that lack of CD47 leads to diminished SIRP α expression in systemic CD11b⁺ phagocyte populations and consequently leads to impairment in the development of CD11b⁺ DCs and macrophages.

It is known that CD8a⁻SIRP α ⁺ cDCs mainly induce CD4⁺ T cell responses, which is an important effector T cell population in autoimmune diseases, whereas CD8a⁺SIRP α ⁻ cDCs has greater ability to induce a CD8⁺ T cell response (9, 44, 46). We had identified that CD11b⁺ DCs (corresponding to CD8a⁻SIRP α ⁺ cDCs) were decreased in number (**Figure 5E**) and their SIRP α expression profile was also decreased in $CD47^{-/-}$ mice with EAU (**Figure 5C**). These results indicate that DCs in $CD47^{-/-}$ mice have decreased ability to induce development of autoreactive CD4⁺ T cell in disease development in response to EAU.

Macrophage and Dendritic Cell Activation Were Suppressed in CD47 Deficient EAU

Given that CD47 has explicit roles in the expression of SIRP α in CD11b⁺ DC/macrophage populations and that the development of




FIGURE 3 | CD47 deficiency in systemic immune cells but not locally in the eye is protective for autoimmune uveitis. Time course of clinical score and representative fundus images using adoptive transfer EAU model. **(A, B)** WT donor mice were immunized and the primed cells were transferred to naive WT or $CD47^{-/-}$ recipient mice. **(A)** Time course of EAU clinical scores were evaluated by fundus observation ($n = 4$ mice per group) **(B)** Representative retinal fundus images on day 13. **(C, D)** WT and $CD47^{-/-}$ donor mice were immunized and the primed cells were transferred to naive WT recipient mice. **(C)** Time course of EAU clinical scores were evaluated by fundus observation ($n = 5-6$ mice per group) **(D)** Representative retinal fundus images on day 17. Data were analyzed by Mann-Whitney U test and expressed as mean \pm SEM. ** $P < 0.01$; n.s., not significant.

$CD11b^{+}$ DC/macrophage populations are impaired in EAU in $CD47^{-/-}$ mice, we next sought to examine activation markers on systemic immune cells that are crucial in the development of EAU. $CD4^{+}$ T cells in LNs, and DCs ($CD11c^{+}$) and macrophages ($CD11b^{+}$) in SPs were examined for activation markers 14 days after immunization. $CD25$, $CD44$ and $CD62L$ were examined in $CD4^{+}$ T cells. $CD25$ is the interleukin-2 receptor alpha chain present on activated T cells. $CD44$ is the molecule involved in leukocyte trafficking required for entry into the retina (47). $CD62L$ is a homing receptor expressed on naive T cells to enter secondary lymphoid tissues and its expression is reduced after activation (48). Effector T cells ($CD62L^{-}CD44^{+}$) and naive T cells ($CD62L^{+}CD44^{-}$) along with $CD25$ collectively represent the activation level of $CD4^{+}$ T cells in EAU (49–52) and various autoimmune diseases (53–55). We found that the frequency between effector and naive $CD4^{+}$ T cells that were positive for these markers were unchanged in $CD47^{-/-}$ mice compared to WT mice (Figures 6A, B), suggesting that T cell activation does not seem to be significantly affected by loss of CD47.

We next examined macrophage and DC expression of MHC-II, $CD80$, and $CD86$; important surface molecules for antigen presentation and co-stimulation required for efficient stimulation of antigen specific $CD4^{+}$ cells. Activated DCs and macrophages

increase expression of these molecules and acquire strong T cell priming capabilities (56, 57). Macrophages increase these molecules after activation (58), but to a lesser extent than DCs. We found that all of the macrophage/DC activation markers were decreased in $CD47^{-/-}$ mice, more strikingly in $CD11c^{+}$ cells (Figures 6C, D). These results indicate that DC/macrophage activity related to antigen presentation are impaired in CD47 deficient EAU. Therefore, this functional defect of antigen presentation in $CD47^{-/-}$ APCs likely caused the suppression of development of $CD4^{+}$ T cells observed in LNs and SPs (Figure 4B). However, lack of CD47 didn't appear to affect the function of $CD4^{+}$ T cells once they were activated given that activation markers in T cells were maintained in $CD47^{-/-}$ mice (Figures 6A, B).

DC Stimulation Capability of T Cells Was Impaired in CD47 Deficient Mice During EAU; Whereas T Cell Proliferation Was Maintained

Our cell proliferation assay in Figure 1 demonstrated that the overall immune cell proliferative response against the autoantigen (IRBP) was suppressed in $CD47^{-/-}$ mice. However, this result represents the proliferative response from total




FIGURE 4 | The number of T cells, macrophages, and DCs are decreased in CD47 deficient mice with autoimmune uveitis. Spleen and lymph node cells from WT and CD47^{-/-} mice were collected before (naive), and 14 days and 21 days post immunization. **(A)** Total cell number in spleens and lymph nodes. **(B)** CD4⁺, CD8⁺, CD19⁺, CD11b⁺, CD11c⁺ cells in spleens and lymph nodes were detected by flowcytometry and cell number of each population was calculated. The cells were gated on live CD45⁺CD3⁺ cells to detect CD4⁺ T cells and CD8⁺ T cells. CD45⁺CD3⁺ population was further gated on CD19, CD11b, and CD11c. Data were analyzed by Student's *t* test and expressed as mean ± SEM. **P*<0.05; ***P*<0.01; ****P*<0.001. n.s., not significant. n=4 mice per group.

immune cells in the culture, including DCs and macrophages as well as CD4⁺ T cells. As we demonstrated earlier, our flow cytometry results suggested that antigen presenting ability of DCs (CD11c⁺) were significantly affected in CD47^{-/-} mice, whereas CD4⁺ T cell functions were maintained. Therefore, we next sought to identify which immune cell population, CD4⁺ T cells or DCs, are responsible in suppressing the immune response and subsequent disease severity of EAU in CD47^{-/-} mice. CD4⁺ T cells in LNs and DCs in SPs isolated by the MACS system were used for mixed lymphocyte proliferation assay. WT and CD47^{-/-} EAU mice 14 days post immunization was utilized. Isolated CD4⁺ T cells were labeled with CellTrace and cultured with DCs for 72 hours in the presence of IRBP-p. The proliferated CD4⁺ T cell was measured by flowcytometry.

The purity of the cells after MACS isolation was more than 94% for CD4⁺ T cells and 99% for DCs (**Supplemental Figure S5**). WT-CD4⁺ T cells and CD47^{-/-} T cells cultured with WT-DCs (WT-T: WT-DC and CD47^{-/-} T: WT-DC, respectively) exhibited comparable proliferation, indicating that CD47^{-/-} CD4⁺ T cells were capable of proliferation with a proper stimulation by DCs. WT-T cells cultured with CD47^{-/-} DC (WT-T: CD47^{-/-} DC) had diminished proliferation compared to WT-T: WT-DC, indicating that DCs from CD47^{-/-} mice were impaired in antigen presentation and T cell stimulation. The proliferation of CD47^{-/-} T cells cultured with CD47^{-/-} DC (CD47^{-/-} T: CD47^{-/-} DC) was impaired compared to CD47^{-/-} T: WT-DC but

comparable to WT-T: CD47^{-/-} DC. These results again highlight that antigen presentation and T cell stimulation ability of CD47^{-/-} DCs are impaired but CD47^{-/-} T cell proliferative ability is maintained (**Figure 7**).

Together, these results demonstrate that CD47^{-/-} CD4⁺ T cells maintain their ability to proliferate, which is consistent with our results in **Figure 6**, where we observed that activation markers in CD47^{-/-} CD4⁺ T were conserved; however, CD47^{-/-} DCs failed to stimulate both WT and CD47^{-/-} T cells. This indicates that antigen presenting functions in DCs are impaired in CD47^{-/-} mice, which is also consistent with our observation in **Figure 6**, in which we demonstrated reduced DC/macrophage activation markers in CD47^{-/-} mice.

In conclusion, we demonstrated that CD47^{-/-} mice have reduced ocular inflammation in EAU. It is likely that the lack of CD47 expression in CD47^{-/-} mice caused reduced expression of its receptor SIRPα specifically on CD11b⁺ APC populations (CD11b⁺ DCs and CD11b⁺ macrophages), resulting in impaired development of these APC populations. In addition, CD47^{-/-} DC/macrophages express reduced amount of activation markers and CD47^{-/-} DCs failed to induce CD4⁺ T cell proliferation, resulting in development of reduced number of autoreactive CD4⁺ cell and consequent reduced immune cell infiltration in the retina. Overall our results demonstrated that CD47 plays an important role in development of autoimmune uveitis by promoting SIRPα⁺ DC maturation.




FIGURE 5 | SIRP α expression in CD11b⁺ macrophages/DCs is decreased in CD47 deficient mice with autoimmune uveitis. WT and CD47^{-/-} mice were immunized and spleen cells were collected 14 days post immunization and examined by flow cytometry. **(A)** Live cells (Zombie UV⁻) excluding a dump gate (T cells; CD3, B cells; CD19, NK cells; NK1.1, and granulocytes; Ly6G) were gated on CD45⁺ cells and further separated by CD11b and CD11c. The gates were defined as follows: Gates a and a'; CD11b⁺CD11c^{high} (CD11b⁺ DCs), gates b and b'; CD11b⁺CD11c⁻ (macrophages), gates c and c'; CD11b^{low}CD11c⁺ (CD11b^{low} DCs), and gates d and d'; CD11b⁺CD11c^{low} (CD11b⁺ DCs). **(B)** SIRP α ^{high} cells and CD8a⁺ cells in **(A)** are shown in green and blue, respectively. **(C)** SIRP α expression on each macrophage and DC population gated in **(A)** was examined. **(D)** Frequency of each macrophage/DC population in spleens was determined by using frequency of each population in ZombieUV⁻ dump⁺CD45⁺ cells. **(E)** Cell number of each macrophage/DC population was obtained by using the number of cells in each SP and the frequency of each population in ZombieUV⁻ cells. Data were analyzed by Student's *t* test and expressed as mean \pm SEM. **P*<0.05; ***P*<0.01; ****P*<0.001. n.s., not significant. *n*=4 mice per group.

DISCUSSION

CD47 plays pleiotropic roles depending on the cell type expressing CD47 and its ligands. In this study, we examined the role of CD47 in development of autoimmune uveitis. Our

results demonstrated that a systemic loss of CD47 is protective in autoimmune uveitis, whereas local retinal loss of CD47 does not affect disease severity. CD47^{-/-} CD4⁺ T cells maintained their proliferative ability and activation markers, however, CD47^{-/-} DCs were hindered in their ability to induce an antigen specific




FIGURE 6 | Activation of macrophages and DCs is suppressed in EAU in CD47 deficient mice. WT and CD47^{-/-} mice were immunized and SP and LN cells were collected 14 days post immunization. Expression of cell activation makers on CD4⁺ T cells, and CD11b⁺ (DCs/macrophages) cells and CD11c⁺ (DCs) cells were evaluated by flow cytometry. **(A, B)** For CD4⁺ T cells, live cells excluding B cell (CD19), NK cells (NK1.1), and granulocytes (Ly6G) were gated on CD45⁺CD3⁺CD4⁺ cells. Frequency of CD62⁺CD44⁺ (effector) cells, CD62⁺CD44⁻ (naïve) cells, and CD25⁺ activated CD4⁺ T cells were examined in spleens **(A)** and lymph nodes **(B)**. **(C, D)** For DCs/macrophages, live cells excluding T cells (CD3), B cells (CD19), NK cells (NK1.1), and granulocytes (Ly6G) were gated on CD45⁺CD11b⁺ and CD45⁺CD11c⁺ cells. Expression of activation markers CD80, CD86, and MHC-II was examined in spleens **(C)** and lymph nodes **(D)**. Data were analyzed by Student's *t* test and expressed as mean ± SEM. **P*<0.05; ***P*<0.01; ****P*<0.001. n.s., not significant. n=4 mice per group.

CD4⁺ T cell response. In CD47^{-/-} mice, DCs and macrophages were decreased in overall numbers as well as key activation markers (CD80, CD86, and MHC-II). The failure in the development of mature CD11b⁺DCs and macrophages, which highly express SIRPα but little CD8α, appears to be a vital mechanism required for EAU induction which is significantly impaired in CD47^{-/-} mice.

Several studies have demonstrated that the interaction of CD47 with its receptor SIRPα is required for development of SIRPα⁺ DCs (22, 43). It has been reported that SIRPα expression

on CD8α⁻ DCs is much greater than that on the corresponding CD8α⁺ DCs and that the number of CD8α⁻ DCs in SIRPα mutant mice was lower than that in wild-type mice (59, 60). Therefore, lack of CD47 can cause insufficient development of CD8α⁻ DCs and decreased expression of SIRPα on these cells, which is likely the reason that CD47^{-/-} mice had decreased number of CD11b⁺ DCs in EAU.

Although the number of CD4⁺ T cells was decreased in CD47^{-/-} mice, frequency of CD62⁺CD44⁺ effector T cells and CD25 expression among CD4⁺ T cells were not affected. This




FIGURE 7 | DCs from CD47 deficient mice failed to stimulate T cells in antigen specific response. WT and *CD47*^{-/-} mice (n=3-4) were immunized and SP and LN cells were collected 14 days post immunization. CD4⁺ T cells and DCs (CD11c⁺) purified from pooled LN cells and SP cells, respectively, were incubated at a ratio of T cells: DCs =2:1 (2×10^5 : 1×10^5 /well) in the presence of 10 μ g/ml IRBP-p in a 96 well U-bottom plate for 3 days. **(A)** Proliferated cells with reduced expression of CellTrace™ Far Red in live CD3⁺ CD4⁺ cells were analyzed by flowcytometry. **(B)** Representative flow cytometry plots. The samples were run triplicates. Data were analyzed by one-way ANOVA followed by Tukey's multiple comparison. Data are expressed as mean \pm SEM. **P<0.01; ***P<0.001.

implies that CD4⁺ T cell activation by antigen stimulation is maintained once these cells are mature, however, it is likely that the decrease in DC and macrophage number and their insufficient antigen presentation ability resulted in a subsequent decrease in the number of mature T cells. This hypothesis is supported by the result from our mixed lymphocyte proliferation assay in which CD4⁺ T cells from *CD47*^{-/-} mice were able to proliferate when stimulated by DCs from WT mice. These results imply that CD4⁺ T cells from *CD47*^{-/-} are able to proliferate with a proper stimulation. An autoimmune disease model in the brain recently reported the similar findings that *CD47*^{-/-} CD4⁺ T cells were activated and produced even larger amounts of cytokines compared to WT mice (61). Moreover, CD47-SIRP α interaction has bidirectional negative effect on regulation of human T cells and DCs (62) and it has been reported that T cell immune response was enhanced by interruption of CD47 in viral infection (63) and tumor immunity (64). Thus, absence of negative regulation due to lack of CD47 could be the reason that CD4⁺ cells maintain their activation level even with insufficient antigen presenting stimulation by DCs.

In the retina, it has been reported that CD47 is expressed in almost all retinal cells including endothelial cells and a small population of cone and rods photoreceptors also express CD47, however, their expression decreases in EAU (65). Although decreased CD47 expression after EAU induction could be a result of inflammation, endothelial CD47 could play an important role in the mechanism of EAU development. Evidence suggests that CD47-SIRP α interactions also regulate

leukocyte adhesion and transmigration to vascular endothelial cells. CD47 expressed on both sides (vascular endothelial cells and leukocytes) could contribute to this mechanism. CD47 on vascular endothelial cells interact with SIRP γ on various leukocytes (66) as well as SIRP α on monocytes (26). However, in our EAU model, when primed WT donor cells were transferred to naïve *CD47*^{-/-} recipient mice, which lacks CD47 on retinal vascular endothelial cells, the recipient mice developed comparable inflammation to control WT recipient mice. This implies that local CD47 expression in retinal endothelial cells does not play a major role in transmigration of leukocytes. Indeed, Van et al. demonstrated that CD47 expression is dispensable in the endothelium (67). On the contrary, CD47 on DCs could be a critical factor in controlling migration and efficient initiation of the immune response (12, 59, 67). It has also been reported that CD47 on T cells is an important regulator of LFA-1 and VLA-4 integrins required for their adhesion to endothelial cell ligands, ICAM-1 and VCAM-1 (61). Therefore, it is possible that in addition to the defect in SIRP α ⁺ DC and macrophage maturation, defect in adhesion and transmigration of T cells and DCs due to their lack of CD47 might play a role in suppressive mechanism of EAU in *CD47*^{-/-} mice.

Complete failure of development of ocular inflammation in an adoptive transfer experiment using *CD47*^{-/-} donor cells and WT recipient cells supported our hypothesis that systemic immune response in *CD47*^{-/-} EAU is impaired. However, when we interpret the data of adoptive transfer of *CD47*^{-/-} mice, we need to be aware that the cells isolated from *CD47*^{-/-} mice are at higher risk of being phagocytized. Indeed, it has been reported that LN cells

from *CD47*^{-/-} mice injected into WT recipient animals are more efficiently phagocytized by macrophages and DCs in recipient mice which express intact SIRP α (68). This could explain why adoptive transfer of *CD47*^{-/-} donor cells to WT recipient mice resulted in complete inhibition of inflammation, whereas *CD47*^{-/-} mice induced with regular EAU was capable to develop mild inflammation. Similar phenomenon is described in an adoptive transfer model of experimental encephalomyelitis (69).

Microglia are resident immune cells in the retina that can be defined as CD11b⁺ cells with low CD45 expression and share some common function with systemic macrophages and DCs. It is noteworthy that microglia expressed SIRP α at a higher level in *CD47*^{-/-} mice compared to WT mice and microglial number were not affected in *CD47*^{-/-} mice. In contrast, SIRP α in the systemic CD11b⁺ cells in *CD47*^{-/-} mice was decreased and the number of CD11b⁺ cells was also fewer. Although the detailed mechanism needs further investigation, these results imply that retinal microglial SIRP α have a distinct regulation mechanism that does not depend on CD47 expression.

In summary, our data demonstrate that systemic CD47 deficiency is protective for autoimmune uveitis. *CD47*^{-/-} mice had reduced inflammation both locally in the eye and systemically. Failure to produce mature CD11b⁺ DCs and macrophages due to their reduced expression of SIRP α appears to be the major mechanism in suppression of EAU in *CD47*^{-/-} mice. Since CD47 interacts with multiple receptors and are involved in complex mechanism of immune regulation, various CD47 receptors could be a treatment option for autoimmune uveitis. Of interest, blocking CD47 with CD47-Fc fusion protein has been shown to effectively impair infiltration of Th17 cells into the central nervous system, preventing and resolving EAE (69). Conversely, an additional study that utilized a CD47-Ab failed to suppress the inflammation associated with EAE (70). Given these conflicting studies the route and agent utilized to inhibit the CD47-SIRP α interaction is likely key when considering a therapeutic approach targeting CD47. Thus, further investigation is needed to elucidate the function of CD47 and its receptors and find a possible treatment option in autoimmune uveitis.

REFERENCES

- Goto H, Mochizuki M, Yamaki K, Kotake S, Usui M, Ohno S. Epidemiological Survey of Intraocular Inflammation in Japan. *Jpn J Ophthalmol* (2007) 51:41–4. doi: 10.1007/s10384-006-0383-4
- Durrani OM, Tehrani NN, Marr JE, Moradi P, Stavrou P, Murray PI. Degree, Duration, and Causes of Visual Loss in Uveitis. *Br J Ophthalmol* (2004) 88:1159–62. doi: 10.1136/bjo.2003.037226
- Rosenbaum JT, Bodaghi B, Couto C, Zierhut M, Acharya N, Pavesio C, et al. New Observations and Emerging Ideas in Diagnosis and Management of non-Infectious Uveitis: A Review. *Semin Arthritis Rheum* (2019) 49:438–45. doi: 10.1016/j.semarthrit.2019.06.004
- Kunimi K, Usui Y, Tsubota K, Mitsuhashi R, Umazume A, Kezuka T, et al. Changes in Etiology of Uveitis in a Single Center in Japan. *Ocul Immunol Inflamm* (2020) 1–6. doi: 10.1080/09273948.2019.1709649
- Lee RW, Nicholson LB, Sen HN, Chan CC, Wei L, Nussenblatt RB, et al. Autoimmune and Autoinflammatory Mechanisms in Uveitis. *Semin Immunopathol* (2014) 36:581–94. doi: 10.1007/s00281-014-0433-9
- Caspi RR, Silver PB, Luger D, Tang J, Cortes LM, Pennesi G, et al. Mouse Models of Experimental Autoimmune Uveitis. *Ophthalmic Res* (2008) 40:169–74. doi: 10.1159/000119871
- Rizzo LV, Silver P, Wiggert B, Hakim F, Gazzinelli RT, Chan CC, et al. Establishment and Characterization of a Murine CD4⁺ T Cell Line and Clone That Induce Experimental Autoimmune Uveoretinitis in B10.A mice. *J Immunol* (1996) 156:1654–60.
- Sanui H, Redmond TM, Kotake S, Wiggert B, Hu LH, Margalit H, et al. Identification of an Immunodominant and Highly Immunopathogenic Determinant in the Retinal Interphotoreceptor Retinoid-Binding Protein (IRBP). *J Exp Med* (1989) 169:1947–60. doi: 10.1084/jem.169.6.1947
- Chen P, Denniston AK, Hirani S, Hannes S, Nussenblatt RB. Role of Dendritic Cell Subsets in Immunity and Their Contribution to Noninfectious Uveitis. *Surv Ophthalmol* (2015) 60:242–9. doi: 10.1016/j.survophthal.2015.01.003
- Okunuki Y, Mukai R, Nakao T, Tabor SJ, Butovsky O, Dana R, et al. Retinal Microglia Initiate Neuroinflammation in Ocular Autoimmunity. *Proc Natl Acad Sci U S A* (2019) 116:9989–98. doi: 10.1073/pnas.1820387116
- Bose T, Diedrichs-Mohring M, Wildner G. Dry Eye Disease and Uveitis: A Closer Look at Immune Mechanisms in Animal Models of Two Ocular

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The animal study was reviewed and approved by the Animal Care Committee of the Massachusetts Eye and Ear.

AUTHOR CONTRIBUTIONS

KC and YO contributed to the study conception and design. YO, ST, and ML performed EAU and quantification. YO and ST performed flow cytometry. KC oversaw all research related experiments. YO, ST, and KC contributed to the analysis and interpretation of the data. YO, ST, and KC wrote the paper. All authors contributed to the article and approved the submitted version.

FUNDING

This study was supported by the National Institute of Health/National Eye Institute: Grants: R01EY031291, R01EY027303 (KC), the Massachusetts Lions Eye Research Fund (KC), and the American Macular Degeneration Foundation Prevention Award (KC).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2021.680568/full#supplementary-material>

- Autoimmune Diseases. *Autoimmun Rev* (2016) 15:1181–92. doi: 10.1016/j.autrev.2016.09.001
12. Fortin G, Raymond M, Van VQ, Rubio M, Gautier P, Sarfati M, et al. A Role for CD47 in the Development of Experimental Colitis Mediated by SIRPalpha +CD103- Dendritic Cells. *J Exp Med* (2009) 206:1995–2011. doi: 10.1084/jem.20082805
 13. Nishimura T, Saito Y, Washio K, Komori S, Respatika D, Kotani T, et al. SIRPalpha on CD11c(+) Cells Induces Th17 Cell Differentiation and Subsequent Inflammation in the CNS in Experimental Autoimmune Encephalomyelitis. *Eur J Immunol* (2020) 50:1560–70. doi: 10.1002/eji.201948410
 14. Lindberg FP, Lublin DM, Telen MJ, Veile RA, Miller YE, Donis-Keller H, et al. Rh-Related Antigen CD47 is the Signal-Transducer Integrin-Associated Protein. *J Biol Chem* (1994) 269:1567–70. doi: 10.1016/S0021-9258(17)42058-8
 15. Reinhold MI, Lindberg FP, Plas D, Reynolds S, Peters MG, Brown EJ. In Vivo Expression of Alternatively Spliced Forms of Integrin-Associated Protein (CD47). *J Cell Sci* (1995) 108(Pt 11):3419–25. doi: 10.1242/jcs.108.11.3419
 16. Hayat SMG, Bianconi V, Pirro M, Jaafari MR, Hatamipour M, Sahebkar A. CD47: Role in the Immune System and Application to Cancer Therapy. *Cell Oncol (Dordr)* (2020) 43:19–30. doi: 10.1007/s13402-019-00469-5
 17. Barclay AN, Brown MH. The SIRP Family of Receptors and Immune Regulation. *Nat Rev Immunol* (2006) 6:457–64. doi: 10.1038/nri1859
 18. Ishikawa-Sekigami T, Kaneko Y, Okazawa H, Tomizawa T, Okajo J, Saito Y, et al. SHPS-1 Promotes the Survival of Circulating Erythrocytes Through Inhibition of Phagocytosis by Splenic Macrophages. *Blood* (2006) 107:341–8. doi: 10.1182/blood-2005-05-1896
 19. Jaiswal S, Jamieson CH, Pang WW, Park CY, Chao MP, Majeti R, et al. CD47 is Upregulated on Circulating Hematopoietic Stem Cells and Leukemia Cells to Avoid Phagocytosis. *Cell* (2009) 138:271–85. doi: 10.1016/j.cell.2009.05.046
 20. Myers LM, Tal MC, Torrez Dulgeroff LB, Carmody AB, Messer RJ, Gulati G, et al. A Functional Subset of CD8(+) T Cells During Chronic Exhaustion is Defined by SIRPalpha Expression. *Nat Commun* (2019) 10:794. doi: 10.1038/s41467-019-08637-9
 21. Tomizawa T, Kaneko Y, Kaneko Y, Saito Y, Ohnishi H, Okajo J, et al. Resistance to Experimental Autoimmune Encephalomyelitis and Impaired T Cell Priming by Dendritic Cells in Src Homology 2 Domain-Containing Protein Tyrosine Phosphatase Substrate-1 Mutant Mice. *J Immunol* (2007) 179:869–77. doi: 10.4049/jimmunol.179.2.869
 22. Matozaki T, Murata Y, Okazawa H, Ohnishi H. Functions and Molecular Mechanisms of the CD47-SIRPalpha Signalling Pathway. *Trends Cell Biol* (2009) 19:72–80. doi: 10.1016/j.tcb.2008.12.001
 23. Seiffert M, Brossart P, Cant C, Cella M, Colonna M, Brugger W, et al. Signal-Regulatory Protein Alpha (SIRPalpha) But Not SIRPbeta is Involved in T-cell Activation, Binds to CD47 With High Affinity, and is Expressed on Immature CD34(+)CD38(-) Hematopoietic Cells. *Blood* (2001) 97:2741–9. doi: 10.1182/blood.V97.9.2741
 24. Liu Y, Buhning HJ, Zen K, Burst SL, Schnell FJ, Williams IR, et al. Signal Regulatory Protein (SIRPalpha), a Cellular Ligand for CD47, Regulates Neutrophil Transmigration. *J Biol Chem* (2002) 277:10028–36. doi: 10.1074/jbc.M109720200
 25. Motegi S, Okazawa H, Ohnishi H, Sato R, Kaneko Y, Kobayashi H, et al. Role of the CD47-SHPS-1 System in Regulation of Cell Migration. *EMBO J* (2003) 22:2634–44. doi: 10.1093/emboj/cdg278
 26. de Vries HE, Hendriks JJ, Honing H, De Lavalette CR, van der Pol SM, Hooijberg E, et al. Signal-Regulatory Protein alpha-CD47 Interactions are Required for the Transmigration of Monocytes Across Cerebral Endothelium. *J Immunol* (2002) 168:5832–9. doi: 10.4049/jimmunol.168.11.5832
 27. Zhang L, Bell BA, Yu M, Chan CC, Peachey NS, Fung J, et al. Complement Anaphylatoxin Receptors C3aR and C5aR are Required in the Pathogenesis of Experimental Autoimmune Uveitis. *J Leukoc Biol* (2016) 99:447–54. doi: 10.1189/jlb.3A0415-157R
 28. Thurau SR, Chan CC, Nussenblatt RB, Caspi RR. Oral Tolerance in a Murine Model of Relapsing Experimental Autoimmune Uveoretinitis (EAU): Induction of Protective Tolerance in Primed Animals. *Clin Exp Immunol* (1997) 109:370–6. doi: 10.1046/j.1365-2249.1997.4571356.x
 29. Caspi RR, Roberge FG, Chan CC, Wiggert B, Chader GJ, Rozenszajn LA, et al. A New Model of Autoimmune Disease. Experimental Autoimmune Uveoretinitis Induced in Mice With Two Different Retinal Antigens. *J Immunol* (1988) 140:1490–5.
 30. Chan CC, Caspi RR, Ni M, Leake WC, Wiggert B, Chader GJ, et al. Pathology of Experimental Autoimmune Uveoretinitis in Mice. *J Autoimmun* (1990) 3:247–55. doi: 10.1016/0896-8411(90)90144-H
 31. Kezuka T, Takeuchi M, Keino H, Usui Y, Takeuchi A, Yamakawa N, et al. Peritoneal Exudate Cells Treated With Calcitonin Gene-Related Peptide Suppress Murine Experimental Autoimmune Uveoretinitis Via IL-10. *J Immunol* (2004) 173:1454–62. doi: 10.4049/jimmunol.173.2.1454
 32. Abcouwer SF, Lin CM, Shanmugam S, Muthusamy A, Barber AJ, Antonetti DA. Minocycline Prevents Retinal Inflammation and Vascular Permeability Following Ischemia-Reperfusion Injury. *J Neuroinflamm* (2013) 10:149. doi: 10.1186/1742-2094-10-149
 33. Itano N, Atsumi F, Sawai T, Yamada Y, Miyaishi O, Senga T, et al. Abnormal Accumulation of Hyaluronan Matrix Diminishes Contact Inhibition of Cell Growth and Promotes Cell Migration. *Proc Natl Acad Sci U S A* (2002) 99:3609–14. doi: 10.1073/pnas.052026799
 34. Tempny JC, Zhou JH, Hodgkin PD, Bryant VL. Superior Properties of CellTrace Yellow as a Division Tracking Dye for Human and Murine Lymphocytes. *Immunol Cell Biol* (2018) 96:149–59. doi: 10.1111/imcb.1020
 35. Lipski DA, Dewispelaere R, Foucart V, Caspers LE, Defrance M, Bruyns C, et al. MHC Class II Expression and Potential Antigen-Presenting Cells in the Retina During Experimental Autoimmune Uveitis. *J Neuroinflamm* (2017) 14:136. doi: 10.1186/s12974-017-0915-5
 36. Nakamura YK, Metea C, Karstens L, Asquith M, Gruner H, Moscirocki C, et al. Gut Microbial Alterations Associated With Protection From Autoimmune Uveitis. *Invest Ophthalmol Vis Sci* (2016) 57:3747–58. doi: 10.1167/iovs.16-19733
 37. Barbour M, Allan D, Xu H, Pei C, Chen M, Niedbala W, et al. IL-33 Attenuates the Development of Experimental Autoimmune Uveitis. *Eur J Immunol* (2014) 44:3320–9. doi: 10.1002/eji.201444671
 38. Usui Y, Takeuchi M, Hattori T, Okunuki Y, Nagasawa K, Kezuka T, et al. Suppression of Experimental Autoimmune Uveoretinitis by Regulatory Dendritic Cells in Mice. *Arch Ophthalmol* (2009) 127:514–9. doi: 10.1001/archophthol.2009.34
 39. Crane IJ, Xu H, Wallace C, Manivannan A, Mack M, Liversidge J, et al. Involvement of CCR5 in the Passage of Th1-type Cells Across the Blood-Retina Barrier in Experimental Autoimmune Uveitis. *J Leukoc Biol* (2006) 79:435–43. doi: 10.1189/jlb.0305130
 40. Chan CC, Caspi RR, Roberge FG, Nussenblatt RB. Dynamics of Experimental Autoimmune Uveoretinitis Induced by Adoptive Transfer of S-Antigen-Specific T Cell Line. *Invest Ophthalmol Vis Sci* (1988) 29:411–8.
 41. Agarwal RK, Caspi RR. Rodent Models of Experimental Autoimmune Uveitis. *Methods Mol Med* (2004) 102:395–419. doi: 10.1385/1-59259-805-6:395
 42. Huang J, Zhou L, Wang H, Luo J, Zeng L, Xiong K, et al. Distribution of Thrombospondins and Their Neuronal Receptor alpha2delta1 in the Rat Retina. *Exp Eye Res* (2013) 111:36–49. doi: 10.1016/j.exer.2013.03.012
 43. Saito Y, Iwamura H, Kaneko T, Ohnishi H, Murata Y, Okazawa H, et al. Regulation by SIRPalpha of Dendritic Cell Homeostasis in Lymphoid Tissues. *Blood* (2010) 116:3517–25. doi: 10.1182/blood-2010-03-277244
 44. Pakalniskyte D, Schraml BU. Tissue-Specific Diversity and Functions of Conventional Dendritic Cells. *Adv Immunol* (2017) 134:89–135. doi: 10.1016/bs.ai.2017.01.003
 45. Williams M, Dutertre CA, Scott CL, McGovern N, Sichien D, Chakarov S, et al. Unsupervised High-Dimensional Analysis Aligns Dendritic Cells Across Tissues and Species. *Immunity* (2016) 45:669–84. doi: 10.1016/j.immuni.2016.08.015
 46. Dudziak D, Kamphorst AO, Heidkamp GF, Buchholz VR, Trumppfheller C, Yamazaki S, et al. Differential Antigen Processing by Dendritic Cell Subsets In Vivo. *Science* (2007) 315:107–11. doi: 10.1126/science.1136080
 47. Xu H, Manivannan A, Liversidge J, Sharp PF, Forrester JV, Crane IJ. Involvement of CD44 in Leukocyte Trafficking at the Blood-Retinal Barrier. *J Leukoc Biol* (2002) 72:1133–41. doi: 10.1189/jlb.72.6.1133
 48. Ivetic A, Ridley AJ. The Telling Tail of L-Selectin. *Biochem Soc Trans* (2004) 32:1118–21. doi: 10.1042/BST0321118
 49. Takeda A, Yamada H, Hasegawa E, Arima M, Notomi S, Myojin S, et al. Crucial Role of P2X7 Receptor for Effector T Cell Activation in Experimental Autoimmune Uveitis. *Jpn J Ophthalmol* (2018) 62:398–406. doi: 10.1007/s10384-018-0587-4

50. Usui Y, Akiba H, Takeuchi M, Kezuka T, Takeuchi A, Hattori T, et al. The Role of the ICOS/B7RP-1 T Cell Costimulatory Pathway in Murine Experimental Autoimmune Uveoretinitis. *Eur J Immunol* (2006) 36:3071–81. doi: 10.1002/eji.200636138
51. Liang D, Zuo A, Zhao R, Shao H, Kaplan HJ, Sun D. Regulation of Adenosine Deaminase on Induced Mouse Experimental Autoimmune Uveitis. *J Immunol* (2016) 196:2646–54. doi: 10.4049/jimmunol.1502294
52. Okunuki Y, Usui Y, Nakagawa H, Tajima K, Matsuda R, Ueda S, et al. Peroxisome Proliferator-Activated Receptor-Gamma Agonist Pioglitazone Suppresses Experimental Autoimmune Uveitis. *Exp Eye Res* (2013) 116:291–7. doi: 10.1016/j.exer.2013.09.017
53. Quast I, Keller CW, Weber P, Schneider C, von Gunten S, Lunemann JD. Protection From Experimental Autoimmune Encephalomyelitis by Polyclonal IgG Requires Adjuvant-Induced Inflammation. *J Neuroinflamm* (2016) 13:42. doi: 10.1186/s12974-016-0506-x
54. Moen LV, Sener Z, Volchenkov R, Svarstad AC, Eriksen AM, Holen HL, et al. Ablation of the Cbeta2 Subunit of PKA in Immune Cells Leads to Increased Susceptibility to Systemic Inflammation in Mice. *Eur J Immunol* (2017) 47:1880–9. doi: 10.1002/eji.201646809
55. Liu Z, Wu L, Zhu J, Zhu X, Zhu J, Liu JQ, et al. Interleukin-27 Signalling Induces Stem Cell Antigen-1 Expression in T Lymphocytes In Vivo. *Immunology* (2017) 152:638–47. doi: 10.1111/imm.12805
56. Lin W, Liu T, Wang B, Bi H. The Role of Ocular Dendritic Cells in Uveitis. *Immunol Lett* (2019) 209:4–10. doi: 10.1016/j.imlet.2019.03.016
57. Steinman RM, Hawiger D, Nussenzweig MC. Tolerogenic Dendritic Cells. *Annu Rev Immunol* (2003) 21:685–711. doi: 10.1146/annurev.immunol.21.120601.141040
58. Huang X, Li Y, Fu M, Xin HB. Polarizing Macrophages In Vitro. *Methods Mol Biol* (2018) 1784:119–26. doi: 10.1007/978-1-4939-7837-3_12
59. Hagnerud S, Manna PP, Cella M, Stenberg A, Frazier WA, Colonna M, et al. Deficit of CD47 Results in a Defect of Marginal Zone Dendritic Cells, Blunted Immune Response to Particulate Antigen and Impairment of Skin Dendritic Cell Migration. *J Immunol* (2006) 176:5772–8. doi: 10.4049/jimmunol.176.10.5772
60. Okajo J, Kaneko Y, Murata Y, Tomizawa T, Okuzawa C, Saito Y, et al. Regulation by Src Homology 2 Domain-Containing Protein Tyrosine Phosphatase Substrate-1 of Alpha-Galactosylceramide-Induced Antimetastatic Activity and Th1 and Th2 Responses of NKT Cells. *J Immunol* (2007) 178:6164–72. doi: 10.4049/jimmunol.178.10.6164
61. Azcutia V, Bassil R, Herter JM, Engelbertsen D, Newton G, Autio A, et al. Defects in CD4+ T Cell LFA-1 Integrin-Dependent Adhesion and Proliferation Protect Cd47-/- Mice From EAE. *J Leukoc Biol* (2017) 101:493–505. doi: 10.1189/jlb.3A1215-546RR
62. Latour S, Tanaka H, Demeure C, Mateo V, Rubio M, Brown EJ, et al. Bidirectional Negative Regulation of Human T and Dendritic Cells by CD47 and its Cognate Receptor Signal-Regulator Protein-Alpha: Down-Regulation of IL-12 Responsiveness and Inhibition of Dendritic Cell Activation. *J Immunol* (2001) 167:2547–54. doi: 10.4049/jimmunol.167.5.2547
63. Cham LB, Torrez Dulgeroff LB, Tal MC, Adomati T, Li F, Bhat H, et al. Immunotherapeutic Blockade of CD47 Inhibitory Signaling Enhances Innate and Adaptive Immune Responses to Viral Infection. *Cell Rep* (2020) 31:107494. doi: 10.1016/j.celrep.2020.03.058
64. Tseng D, Volkmer JP, Willingham SB, Contreras-Trujillo H, Fathman JW, Fernhoff NB, et al. Anti-CD47 Antibody-Mediated Phagocytosis of Cancer by Macrophages Primes an Effective Antitumor T-cell Response. *Proc Natl Acad Sci U S A* (2013) 110:11103–8. doi: 10.1073/pnas.1305569110
65. Liu J, Tang M, Harkin K, Du X, Luo C, Chen M, et al. Single-Cell RNA Sequencing Study of Retinal Immune Regulators Identified CD47 and CD59a Expression in photoreceptors-Implications in Subretinal Immune Regulation. *J Neurosci Res* (2020) 98:1498–513. doi: 10.1002/jnr.24618
66. Azcutia V, Stefanidakis M, Tsuboi N, Mayadas T, Croce KJ, Fukuda D, et al. Endothelial CD47 Promotes Vascular Endothelial-Cadherin Tyrosine Phosphorylation and Participates in T Cell Recruitment at Sites of Inflammation In Vivo. *J Immunol* (2012) 189:2553–62. doi: 10.4049/jimmunol.1103606
67. Van VQ, Lesage S, Bouguermouh S, Gautier P, Rubio M, Levesque M, et al. Expression of the Self-Marker CD47 on Dendritic Cells Governs Their Trafficking to Secondary Lymphoid Organs. *EMBO J* (2006) 25:5560–8. doi: 10.1038/sj.emboj.7601415
68. Blazar BR, Lindberg FP, Ingulli E, Panoskaltis-Mortari A, Oldenborg PA, Iizuka K, et al. CD47 (Integrin-Associated Protein) Engagement of Dendritic Cell and Macrophage Counterreceptors is Required to Prevent the Clearance of Donor Lymphohematopoietic Cells. *J Exp Med* (2001) 194:541–9. doi: 10.1084/jem.194.4.541
69. Gao Q, Zhang Y, Han C, Hu X, Zhang H, Xu X, et al. Blockade of CD47 Ameliorates Autoimmune Inflammation in CNS by Suppressing IL-1-triggered Infiltration of Pathogenic Th17 Cells. *J Autoimmun* (2016) 69:74–85. doi: 10.1016/j.jaut.2016.03.002
70. Han MH, Lundgren DH, Jaiswal S, Chao M, Graham KL, Garris CS, et al. Janus-Like Opposing Roles of CD47 in Autoimmune Brain Inflammation in Humans and Mice. *J Exp Med* (2012) 209:1325–34. doi: 10.1084/jem.20101974

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Okunuki, Tabor, Lee and Connor. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.