



## OPEN ACCESS

## EDITED AND REVIEWED BY

Malcolm Ronald Starkey,  
Department of Immunology and  
Pathology, Monash University,  
Australia

## \*CORRESPONDENCE

Dominique M. A. Bullens  
dominique.bullens@kuleuven.be

## SPECIALTY SECTION

This article was submitted to  
NK and Innate Lymphoid Cell Biology,  
a section of the journal  
Frontiers in Immunology

RECEIVED 30 August 2022

ACCEPTED 01 September 2022

PUBLISHED 21 September 2022

## CITATION

Jonckheere A-C, Seys SF, Steelant B,  
Decaestecker T, Dekoster K, Cremer J,  
Dilissen E, Schols D, Iwakura Y,  
Vande Velde G, Breynaert C,  
Schrijvers R, Vanoirbeek J,  
Ceuppens JL, Dupont LJ and  
Bullens DMA (2022) Corrigendum:  
Innate lymphoid cells are required to  
induce airway hyperreactivity in a  
murine neutrophilic asthma model.  
*Front. Immunol.* 13:1032423.  
doi: 10.3389/fimmu.2022.1032423

## COPYRIGHT

© 2022 Jonckheere, Seys, Steelant,  
Decaestecker, Dekoster, Cremer,  
Dilissen, Schols, Iwakura, Vande  
Velde, Breynaert, Schrijvers,  
Vanoirbeek, Ceuppens, Dupont  
and Bullens. This is an open-  
access article distributed under the  
terms of the [Creative Commons  
Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use,  
distribution or reproduction in other  
forums is permitted, provided the  
original author(s) and the copyright  
owner(s) are credited and that the  
original publication in this journal is  
cited, in accordance with accepted  
academic practice. No use,  
distribution or reproduction is  
permitted which does not comply with  
these terms.

# Corrigendum: Innate lymphoid cells are required to induce airway hyperreactivity in a murine neutrophilic asthma model

Anne-Charlotte Jonckheere<sup>1</sup>, Sven F. Seys<sup>1</sup>, Brecht Steelant<sup>1</sup>,  
Tatjana Decaestecker<sup>2</sup>, Kaat Dekoster<sup>3</sup>, Jonathan Cremer<sup>1</sup>,  
Ellen Dilissen<sup>1</sup>, Dominique Schols<sup>4</sup>, Yoichiro Iwakura<sup>5</sup>,  
Greetje Vande Velde<sup>3</sup>, Christine Breynaert<sup>1</sup>, Rik Schrijvers<sup>1</sup>,  
Jeroen Vanoirbeek<sup>6</sup>, Jan L. Ceuppens<sup>1</sup>, Lieven J. Dupont<sup>2,7</sup>  
and Dominique M. A. Bullens<sup>1,8\*</sup>

<sup>1</sup>Department of Microbiology, Immunology and Transplantation, Allergy and Clinical Immunology Research Group, KU Leuven, Leuven, Belgium, <sup>2</sup>Department of Chronic Diseases, Metabolism and Ageing, Laboratory of Respiratory Diseases and Thoracic Surgery, KU Leuven, Leuven, Belgium,

<sup>3</sup>Department of Imaging and Pathology, Biomedical MRI Unit/Molecular Small Animal Imaging Center (MoSAIC), KU Leuven, Leuven, Belgium, <sup>4</sup>Department of Microbiology, Immunology and Transplantation, Laboratory of Virology and Chemotherapy, KU Leuven, Leuven, Belgium, <sup>5</sup>Centre for Animal Disease Models, Research Institute for Biomedical Sciences, Tokyo University of Science, Chiba, Japan, <sup>6</sup>Department of Public Health and Primary Care, Centre for Environment and Health, KU Leuven, Leuven, Belgium, <sup>7</sup>Clinical Division of Respiratory Medicine, UZ Leuven, Leuven, Belgium, <sup>8</sup>Clinical Division of Paediatrics, UZ Leuven, Leuven, Belgium

## KEYWORDS

innate lymphoid cells (ILCs), non-allergic asthma, murine model, neutrophilic inflammation, airway hyperreactivity

## A corrigendum on

### Innate lymphoid cells are required to induce airway hyperreactivity in a murine neutrophilic asthma model

by Jonckheere A-C, Seys SF, Steelant B, Decaestecker T, Dekoster K, Cremer J, Dilissen E, Schols D, Iwakura Y, Vande Velde G, Breynaert C, Schrijver R, Vanoirbeek J, Ceuppens JL, Dupont LJ, Bullens DMA (2022) *Front. Immunol.* 13:849155. doi: 10.3389/fimmu.2022.849155.

In the published article, there was an error made in the provided concentration lipopolysaccharide (LPS) applied endonasally in the murine model for neutrophilic asthma.

A correction has been made to Material and Methods section, “*Neutrophilic asthma model*”. The sentence previously stated:

“Wild-type, SCID, Rag2<sup>-/-</sup> γC<sup>-/-</sup>, and IL-17A<sup>-/-</sup> mice were endonasally challenged with 2 μg LPS (in a solution of 50 μl; 40 mg/ml) or 50 μl of saline (0.9% NaCl, B. Braun) on four consecutive days (Supplementary Figure 1A).”

The corrected sentence appears below:

“Wild-type, SCID, Rag2<sup>-/-</sup> γC<sup>-/-</sup>, and IL-17A<sup>-/-</sup> mice were endonasally challenged with 2 μg LPS (in a solution of 50 μl; 40 μg/ml) or 50 μl of saline (0.9% NaCl, B. Braun) on four consecutive days (Supplementary Figure 1A).”

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.