



Alteration of Endocrine Hormones and Antibody Responses in Different Spectrum of Tuberculosis Disease

Yosef Tsegaye^{1,2†}, Wasihun Admassu^{2,3,4†}, Abebe Edao¹, Samuel Kinde¹, Meaza Gentu⁴, Markos Negash⁴, Tadelo Wondmagegn⁴, Addisu Gize⁵, Martha Zewdie², Kidist Bobosha² and Liya Wassie^{2*}

OPEN ACCESS

Edited by:

Simone A. Joosten,
Leiden University Medical Center,
Netherlands

Reviewed by:

Thomas Lindstrom,
Statens Serum Institut (SSI), Denmark
Taylor Foreman,
National Institutes of Health (NIH),
United States

*Correspondence:

Liya Wassie
liya.wassie@ahri.gov.et

[†]These authors have contributed
equally to this work and share
first authorship

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Immunology

Received: 05 January 2022

Accepted: 07 February 2022

Published: 25 February 2022

Citation:

Tsegaye Y, Admassu W, Edao A,
Kinde S, Gentu M, Negash M,
Wondmagegn T, Gize A, Zewdie M,
Bobosha K and Wassie L (2022)
Alteration of Endocrine Hormones and
Antibody Responses in Different
Spectrum of Tuberculosis Disease.
Front. Immunol. 13:849321.
doi: 10.3389/fimmu.2022.849321

¹ Department of Medical Laboratory Science, Addis Ababa University, Addis Ababa, Ethiopia, ² Mycobacterial Diseases Research Directorate, Armauer Hansen Research Institute, Addis Ababa, Ethiopia, ³ Immunology and Molecular Biology Unit, Jimma University Medical Center, Jimma, Ethiopia, ⁴ Department of Immunology, University of Gondar, Gondar, Ethiopia, ⁵ Department of Microbiology, St Paul's Hospital Millennium Medical College, Addis Ababa, Ethiopia

Effective control of *Mycobacterium tuberculosis* (*Mtb*) infection is mediated by multifaceted factors that involve both the endocrine and immune system. Profiling hormones and antibodies in different stages of TB provides insight in the pathogenesis of the disease. In this study, we profiled endocrine hormones (dehydroepiandrosterone (DHEA), cortisol, testosterone, estradiol, growth hormone and leptins) and *Mtb* strain H37RV lipoarabinomannan (LAM)-specific antibody levels in plasma samples, collected from pulmonary TB (PTB) patients, TB lymphadenitis (TBLN) patients and latently infected (QFT-positive) or uninfected (QFT-negative) apparently healthy individuals using ELISA. Plasma levels of leptin and DHEA were significantly low in PTB and TBLN patients compared to healthy controls ($P < 0.0001$ and $P = 0.02$, respectively), whereas these levels significantly increased following anti-TB treatment ($P = 0.002$ and $P = 0.0001$, respectively) among TB patients. The levels of estradiol and testosterone significantly improved following anti-TB treatment ($P = 0.03$ and $P = 0.0003$, respectively), whereas cortisol and growth hormones declined significantly ($P < 0.05$). Similarly, LAM-specific IgG, IgM and IgA were significantly higher in PTB patients compared to other groups, whereas levels of IgG1 subtype were significantly higher among LTBI groups compared to both TB patients and QFT-negative individuals ($P < 0.0001$). Overall, we observed significantly variable levels of endocrine hormones as well as immunoglobulins across the spectrum of TB illness and such profiling has a significant contribution in selection of effective biomarkers that have roles in TB treatment monitoring or diagnostics. Although this study did not show a functional association between hormones and antibodies, alterations in the levels of these biomarkers suggest the key roles these markers play in TB pathogenesis.

Keywords: hormones, antibodies, tuberculosis, immunity, pathogenesis

INTRODUCTION

The spectrum of tuberculosis (TB) pathogenesis often lies on interplay between the bacterium causing the illness, *Mycobacterium tuberculosis* (*Mtb*), and the host defense system, which is also maneuvered by cells of the immune system. In both experimental models and human patients, hormones are shown to act as immunomodulators and play a significant role in the pathogenesis of a disease (1). Hormones and the immune system share receptors and ligands that interact through cytokine- and hormone-producing cells, influencing the course of infection processes (2). Studies have demonstrated that specific hormones, such as cortisol, growth hormones and dehydroepiandrosterone (DHEA) correlate with tuberculosis (TB) treatment outcomes (3, 4). On the other hand, although antibody responses are often regarded as inferior to cell-mediated immunity, their role in protection against TB has been demonstrated in several studies (5–9). Profiling these two components of the body's defense system across a spectrum of TB illness is important to better understand TB pathogenesis and provide an additional insight to the ongoing biomarker pool investigations in TB control efforts. Tuberculosis is often characterized by a spectrum of infection stages and illnesses (10). In this study, we aimed to profile levels of lipoarabinomannan (LAM)-specific immunoglobulins (Igs) and selected hormones in different spectra of *Mtb* infections and/or TB diseases (latently un/infected individuals and TB patients) using enzyme-linked immunosorbent assay (ELISA). Lipoarabinomannan, a glycolipid, is a significant part of mycobacterial surface antigen, having roles in virulence, pathogenesis, and known to elicit immunological response (11). Its repetitive D-arabinofuranose residues allows it to direct B cell stimulatory functions (12), making it a suitable antigen choice in our study.

MATERIALS AND METHODS

Study Setting

A pool of 200 plasma samples, collected between 2005 and 2013, were systematically retrieved (based on quality, adequate sample availability, and linked clinical and socio-demographic data) from a biorepository at the Armauer Hansen Research Institute (AHRI) laboratory. Of these, 40 newly diagnosed microscopy or culture confirmed pulmonary TB (PTB) patients (before and after anti-TB treatment), 40 TB lymphadenitis patients (TBLN), 40 latently

infected (QuantiFERON (QFT) TB Gold test -positive) and 40 uninfected (QFT-negative) individuals were randomly retrieved. Based on the repository database, all samples were collected from a cohort of voluntary, adult, HIV negative individuals, who resided in Addis Ababa, the capital city of Ethiopia, Hossana and Butajira areas, (which are ~230 Km and 130 Km, respectively, southwest of Addis Ababa). Majority of the PTB patients had cough (38/40), night sweating (32) and hemoptysis in a few cases (8/40). Similarly, night sweating was the most common symptom exhibited (22/40) in TBLN cases, while cough (5/40) and hemoptysis (4/40) were rare occurrences. The most commonly observed sites of lymph node enlargements in TBLN cases were cervical area (12/40) and neck (11/40); however, other sites including inguinal, submandibular, clavicular and axial sites were also involved. Additional datasets are summarized in **Table 1**. Samples with a clinical history of severe malnourishment, anemia and other debilitating illnesses were excluded during the initial screening and enrollment of participants. Also, samples with icteric and turbid plasma were not included in these analyses.

Plasma Sample Preparation and Quantification of Hormone and Antibody Concentrations Using Enzyme-Linked Immunosorbent Assay (ELISA)

Once retrieved from -80°C freezers, plasma samples were thawed on ice and spun at 1500 rpm for 5 minutes to transfer clear plasma supernatants into new collection tubes that were kept at 4°C until assayed by enzyme-linked immunosorbent assay (ELISA). Concentrations of hormones (DHEA, cortisol, testosterone, estradiol, growth hormone and leptin) or LAM-specific antibody (IgG, IgM, IgA or IgG subclasses) levels were measured using ELISA according to the manufacturer's instructions (Eagle Bioscience, Italy).

Briefly, 96-well ELISA plates, pre-coated with hormones of interest (Eagle Bioscience, Italy), were incubated with plasma samples either at room temperature or at 37°C, depending on the hormones of interest and washed with diluted phosphate buffer saline (PBS) containing 0.05% Tween 20, followed by addition of enzyme-substrate conjugates for antibody binding. Similarly, Immulon I 96-well high-binding polystyrene microtiter plates (Thermo Fisher Scientific, U.S.A.) were coated with 1µg/ml of *M. tuberculosis* strain H37RV purified mannosylated lipoarabinomannan (LAM) (generously donated by BEI Resources, <https://www.beiresources.org/>) in 0.05M carbonate/bicarbonate coating buffer (pH 9.6) and incubated overnight at 4°C. The following day, plates were washed

TABLE 1 | Demographic and clinical features of study participants.

Parameter	Study groups			
	QFT-negative (n = 40)	QFT-positive (n = 40)	PTB (n = 40)	TBLN (n = 40)
Age (years mean ± SD)	20.95 ± 7.6	24.5 ± 9.7	26.3 ± 7.8	29.61 ± 14.5
Sex	Male (N, %)	22 (55)	22 (55)	12 (30)
	Female (N, %)	18 (45)	18 (45)	18 (45)
Presence of BCG scar (N, %)	17 (42.5)	13 (32.5)	6 (15)	5 (12.5)*
Body mass index (kg/m ²) (mean ± SD)	19.0 ± 3.2	20.2 ± 4.2	18.3 ± 3	Not Available

QFT, QuantiFERON; PTB, pulmonary tuberculosis; TBLN, tuberculosis lymphadenitis; *data not available.

with PBS containing 0.1% of Tween-20 and blocked using 0.1% bovine serum albumin (BSA). Then plasma samples and standards were added into wells after dilution (for Ig analyses sample dilutions ranged from 1:25,000 to 1:100), followed by addition of enzyme-substrate conjugates for antibody binding. In both cases, a color change was developed that was later captured as optical density (OD) using an ELISA reader (reading at 450 nm against a reference wavelength at 620 nm). All laboratory assays were done using commercially available kits following the manufacturers' instructions and standard operating procedures. Results were also interpreted according to the manufacturers' instructions. Plasma concentrations of hormones or antibodies were finally extrapolated from the standard curve and all statistical analyses were done using extrapolated values from the standard curves, assuming best curve fit (with highest R values). For diluted samples, final concentrations were calculated by multiplying the extrapolated concentration with dilution factor.

Ethics Approval

This study was performed in accordance with the Declaration of Helsinki and approved by the AHRI/ALERT Ethics Review Committee (Ref. No. PO07/18 and PO08/18), by the ethics committees of the Addis Ababa University, Department of Medical Laboratory Sciences (Ref. No. DRERC/326/18/MLS) and School of Biomedical and Laboratory Science, University of Gondar (UOG). Only anonymized samples were used to ensure privacy and confidentiality of study participants. In addition, institutional permission was obtained from AHRI to access biorepository samples and linked archived data.

Statistical Analyses

In this study, all plasma levels of hormones (cortisol, DHEA, growth hormone, estradiol, testosterone and leptin) and Igs (IgG, IgG1, IgG2, IgG3, IgA, IgM) are presented in concentrations and were considered as dependent variables, whereas variables including clinical presentations and socio-demographic data (such as age, sex, BMI, BCG vaccination status) were considered as independent variables. Statistical analyses were done using GraphPad Prism 7.01 and SPSS version 26: comparisons between groups were done using non-parametric statistics (One-way ANOVA (Kruskal-Wallis test) followed by Dunnett's multiple test comparison). Correlation between hormones and Ig levels with clinical status were done using Spearman correlational analysis. To compare hormone profiles or Igs in PTB patients before and after treatments, a nonparametric, Wilcoxon matched-pairs signed rank test was used. Multivariate analysis was also used to compare Ig levels between groups. Data are summarized using descriptive statistics, frequency tables and figures and a P-value less than 0.05 was considered statistically significant.

RESULTS

Plasma Levels of Hormones Among the Study Groups

The levels of DHEA were significantly lower in PTB patients compared to QFT-negative and TBLN patients ($P < 0.02$ and

$P = 0.01$, respectively) (**Figure 1A**). Leptin level was also significantly lower in PTB and TBLN patients compared to apparently healthy individuals who were either QFT-positive or negative (**Figure 1B**). On the other hand, plasma cortisol level was significantly higher in PTB and TBLN patients compared to apparently healthy groups, while its level was even higher in PTB patients compared to the TBLN cases (**Figure 1C**). In a similar way, age and sex adjusted plasma levels of testosterone, estradiol and growth hormone levels were measured; however, no apparent difference was observed between the different study groups (Data not shown).


M. tuberculosis LAM-Specific Antibody Profiles Among the Study Groups

In a similar trend, we compared levels of concentrations of immunoglobulins in plasma of TB patients (both PTB and TBLN) and apparently healthy individuals, who were either latently infected (QFT-positive) or non-infected (QFT-negative). We observed a significant difference in the levels of IgG, IgM and IgA (**Figure 2A**), where PTB patients exhibited higher levels of the immunoglobulins compared to the other study groups. Similarly, levels of IgG subtypes, particularly IgG2 and IgG3, were relatively higher in PTB patients (**Figure 2B**). On the other hand, all the apparently healthy groups (regardless of the QFT responses) had higher levels of IgG1 subclass compared to both TB patient groups ($P < 0.0001$) (**Figure 2B**).

Multiple regression was performed between hormone levels and clinical and other demographic variables (presence or absence of BCG scar, age, sex, BMI); however, no apparent association was observed, except for cortisol, whose levels poorly correlated with age ($R^2 = 0.24$, $p = 0.004$), sex ($R^2 = 0.24$, $p = 0.04$) and BMI ($R^2 = 0.24$, $p = 0.0001$) and testosterone, whose levels strongly correlated with BMI ($R^2 = 0.670$, $P = 0.0001$). A similar regression analysis between antibody levels and socio-demographic characteristics showed no apparent associations between the two variables; however the associations were more visible between antibody levels and types with underlying mycobacterial infections and disease states as shown in **Figure 2**. On the other hand, a Spearman correlation analysis was performed between levels of LAM-specific Ig responses and presence or absence of BCG scar and the analyses indicated variable degrees of correlation between Ig levels (particularly IgG, IgM, IgA and IgG2) and presence of BCG scar with correlation coefficients ($\rho = 0.65$, $P < 0.0001$; $\rho = 0.59$, $P < 0.0001$; $\rho = 0.39$, $P = 0.005$ and $\rho = 0.59$, $P < 0.0001$, respectively). However, a similar analysis between hormone levels and some of the independent variables showed no correlation between hormones levels and other variables (Data not shown).

Impact of Anti-TB Treatment on Plasma Levels of Hormones and LAM-Specific Immunoglobulins in PTB Patients

To assess the impact of anti-TB treatment on plasma hormone concentrations, we compared the median plasma levels of hormones in PTB patients at baseline (before initiation of anti-TB treatment) and following completion of treatment (after 6 months of therapy) and observed significant alterations in the plasma levels of hormones after anti-TB treatment (**Figure 3**). The median levels



of DHEA, leptin, testosterone and estradiol were significantly elevated following completion of treatment, whereas there was an apparent decline in the median levels of cortisol and growth hormones after treatment completion (Figure 3).


Similarly, plasma antibody levels were compared in PTB patients before and after completion of anti-TB treatment; however, no apparent difference was observed, except for IgM, where the levels significantly declined following anti-TB treatment ($P=0.001$) (Data not shown).

DISCUSSION

How hormones interact with the immune systems dictate the way optimal immune responses are also induced during host-pathogen interactions (1). Such optimal responses are often triggered through changes in the secretion patterns of hormones as well as the immune system. The main aim of this study was to profile hormone and antibody responses and measure how their levels alter during a spectrum of illnesses such as TB (latent TB infection, pulmonary and extrapulmonary TB disease or following treatment completion). Though cell-mediated immunity demonstrates a major role in protection against TB, antibodies are also recently reported to play a significant role in bacterial clearance and enhanced phagolysosomal maturation during TB infection (7). In this study, we profiled plasma levels of selected endocrine hormones (cortisol, dehydroepiandrosterone, leptin, human growth hormone, testosterone, and estradiol) and antibodies (IgG, IgG1, IgG2, IgG3, IgA, IgM) in tuberculosis patients with varied forms of the disease (pulmonary TB and TB lymphadenitis) and in comparison to apparently healthy individuals, who were further categorized as latently infected or uninfected.

The findings of our study suggest that active TB disease is characterized by elevated plasma levels of cortisol and decreased DHEA, resulting in a greater cortisol/DHEA ratio when compared to both the healthy controls and latently infected groups. This has been also evident from earlier studies, where cortisol level was significantly higher in TB patients compared to healthy individuals and that this increased level also correlated with disease severity, whereas it was the reverse for DHEA (13, 14). Previous studies also showed that cortisol promotes T-helper 2 (Th2) immune activity, partly by suppressing Th1 cells and that its natural antagonist, DHEA, is shown to promote Th1 cytokine production, while interfering with Th2 cytokine synthesis (15, 16). Because infection with intracellular pathogens such as TB necessitates the expression of Th1 cytokines such as IFN- γ , a central macrophage-activating cytokine that is also involved in immune protection against *MTB* (17), an increase in plasma levels of cortisol and a decrease in DHEA levels may impair the immune response against TB, favoring increased susceptibility and disease progression (18).

Furthermore, the plasma levels of cortisol and DHEA were considerably altered following therapy, with cortisol levels dramatically decreasing and DHEA increasing. Others have also reported a similar finding that cortisol levels decline throughout TB treatment in cured patients but remain unchanged in patients with a failed treatment outcome, whereas DHEA levels steadily increase in the cured groups






FIGURE 3 | Median plasma levels of hormones in PTB patients before and after anti-TB treatment.

during treatment (3). These changes demonstrate the importance of these hormones in the immunopathogenesis of tuberculosis. Similar alterations have been also reported from animal studies, where levels of DHEA improved, following treatment with DHEA sulfate, boosting the production of IFN- γ (19).

Similar like other reports (14, 20, 21), the TB patients had considerably lower plasma levels of leptin when compared to both latently infected and uninfected groups. However, this level significantly improved in PTB patients following anti-TB treatments (3). On the other hand, a few other studies have reported an increase or no change in leptin levels in TB patients (22, 23). The reasons for this disparity could be related to nutritional or dietary factors, genetic, environmental or treatment initiation time. Other studies have also demonstrated the inhibitory role persistent activation of pro-inflammatory cytokines play on leptin synthesis in TB patients (24). Because leptin concentration is also linked to adipose tissue mass (25), the lower leptin levels seen in TB patients and its change after therapy could be linked to disease-associated weight loss and the association between leptin levels and wasting seen in most TB patients.

Our analysis on the plasma levels of sex hormones (testosterone and estradiol) showed no apparent difference among the study groups; however, others have shown a considerably lower testosterone levels in TB patients (4, 13), which was also associated with other pro-inflammatory

cytokines such as IFN- γ , IL-6, and TGF- β (26). However, there was a significant alteration in the levels of both testosterone and estradiol concentrations in PTB patients following anti-TB treatment and this was also consistent with earlier reports (3). Although no significant difference was observed in the levels of growth hormones (GH) between the study groups, the level significantly declined after treatment in PTB patients. However, an earlier study showed higher levels of GH in TB patients compared to healthy individuals (14). On the other hand, others reported the roles GHs play in stimulating human monocytes to destroy *Mtb* by serving as a human macrophage-activating factor (27).

Similarly, plasma immunoglobulin levels were measured in individuals exhibiting different clinical presentations and exposure to TB and here *Mtb-derived* LAM was used as an immunogen to examine specific Ig levels. The findings of our study indicate that there is a significant difference in antibody profiles in different disease states, possibly suggesting for their role as potential biomarkers in future vaccines or diagnostics developments (28, 29). While there are no tools to specifically distinguish latent TB infection from active disease, some of these markers, particularly LAM-specific IgG and IgA responses, seem to play that role in TB, as also reported by others (5, 29).

This difference in the levels of IgG might not be the same, however, if HIV becomes a factor, as reported earlier (30), where HIV-positive TB patients exhibited a significantly lower LAM-

specific IgG subclasses compared to non-infected TB patients. Similarly, PTB patients exhibited a significantly higher levels of LAM-specific IgA compared to apparently healthy controls, who were either QFT-positive or negative. Earlier studies have also reported significantly higher levels of *Mtb* antigen-specific IgA responses in TB patients compared to healthy controls (31–34). In the current study, latently infected individuals showed a higher level of IgG1 titer, whereas earlier studies reported higher IgG1 and IgG2 levels in PTB patients (35), indicating their functional role during TB disease (9). The increased levels of IgG1 subtype in LTBI groups indicate its distinct glycosylation characteristics that influence binding of Fc receptors on effector cells, which may play roles during bacterial containment (40).

In this study, we also compared the breadth of the antibodies in PTB patients before and after treatment and the levels of LAM-specific IgM significantly dropped following treatment, possibly implying for the impact of treatment on *Mtb* clearance and containment of active infection. This also possibly implies to the role of IgM as a useful biomarker for treatment monitoring in TB patients.

In conclusion, lower levels of anabolic hormones such as DHEA and leptin and elevated levels of cortisol could mediate the gradual debilitation and other disabling symptoms associated with TB. Furthermore, the observation that anti-LAM Ig responses (as shown in antibody titer, type and subtype) differ significantly during TB stages, indicates a valuable roles these antibodies play during TB pathogenesis. The altered plasma levels of hormones as well as immunoglobulins during treatment suggest that evaluating these biomarkers could have prognostic importance, particularly in the evaluation of treatment outcomes, biomarker research and bacterial containment. Overall, profiling such biomarkers in a spectrum of an illness, particularly in a disease like TB, will have a significant contribution in future biomarker evaluations for treatment monitoring or diagnostics. However, the observed differences in the levels of these markers could not account for intrinsic factors related to the use of repository specimens, missing data on dietary intake or menstrual cycles/ovulation, particularly in female participants. This study does not show the functional association between hormones and antibodies; however, the observed changes in the levels of these biomarkers could possibly indicate their key roles in TB pathogenesis. A large-scale study might therefore be warranted to better understand the interaction between the endocrine and the immune system, particularly the innate immunity.

REFERENCES

- Webster J, Tonelli L, Sternberg EM. Neuroendocrine Regulation of Immunity. *Annu Rev Immunol* (2002) 20:125–63. doi: 10.1146/annurev.immunol.20.082401.104914
- Haddad JJ, Saadé NE, Safieh-Garabedian B. Cytokines and Neuro-Immune-Endocrine Interactions: A Role for the Hypothalamic-Pituitary-Adrenal Revolving Axis. *J Neuroimmunol* (2002) 133:1–19. doi: 10.1016/S0165-5728(02)00357-0
- Kleynhans L, Ruzive S, Ehlers L, Thiarit L, Chegou NN, Conradie M, et al. Changes in Host Immune-Endocrine Relationships During Tuberculosis

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by The AHRI/ALERT Ethics Review Committee, The ethics committee of the Addis Ababa University, Department of Medical Laboratory Sciences and School of Biomedical and Laboratory Science, University of Gondar. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

YT and WA did all the laboratory work, analyzed data, and wrote the 1st draft of the manuscript. AE, SK, MG, MN, TW, AG, MZ, and KB analyzed the data, edited the manuscript, and supervised the work. LW conceived and designed the study, analyzed the data, supervised the study, and edited the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by EDCTP grant through Collaboration and Integration of Tuberculosis Vaccine Trials in Europe and Africa (TBTEA) Consortium [grant number MS. 2010.18000.002] and core funds from Armauer Hansen Research Institute (AHRI), received from SIDA and NORAD. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

ACKNOWLEDGMENTS

The authors are grateful to all study participants and study nurses (Alemayehu Kifle, Bamlak Tesema and Haregewoin Yeshitela) from AHRI. The following reagent was obtained through BEI Resources, NIAID, NIH: *Mycobacterium tuberculosis*, Strain H37Rv, Purified Lipoarabinomannan (LAM), NR-14848.

Treatment in Patients With Cured and Failed Treatment Outcomes. *Front Immunol* (2017) 8:690. doi: 10.3389/fimmu.2017.00690

- Fernández R, Díaz A, D'Attilio L, Bongiovanni B, Santucci N, Bertola D, et al. An Adverse Immune-Endocrine Profile in Patients With Tuberculosis and Type 2 Diabetes. *Tuberculosis* (2016) 101:95–101. doi: 10.1016/j.tube.2016.09.001
- Li H, Wang X-x, Wang B, Fu L, Liu G, Lu Y, et al. Latently and Uninfected Healthcare Workers Exposed to TB Make Protective Antibodies Against *Mycobacterium Tuberculosis*. *Proc Natl Acad Sci* (2017) 114:5023–8. doi: 10.1073/pnas.1611776114
- Lu LL, Chung AW, Rosebrock TR, Ghebremichael M, Yu WH, Grace PS, et al. A Functional Role for Antibodies in Tuberculosis. *Cell* (2016) 167:433–43. doi: 10.1016/j.cell.2016.08.072

7. de L, Costello A, Kumar A, Narayan V, Akbar M, Ahmed S, et al. Does Antibody to Mycobacterial Antigens, Including Lipoarabinomannan, Limit Dissemination in Childhood Tuberculosis? *Trans R Soc Trop Med Hygiene* (1992) 86:686–92. doi: 10.1016/0035-9203(92)90192-F
8. Kumar SK, Singh P, Sinha S. Naturally Produced Opsonizing Antibodies Restrict the Survival of Mycobacterium Tuberculosis in Human Macrophages by Augmenting Phagosome Maturation. *Open Biol* (2015) 5:150171. doi: 10.1098/rsob.150171
9. Zimmermann N, Thormann V, Hu B, Köhler AB, Imai-Matsushima A, Lochter C, et al. Human Isotype-Dependent Inhibitory Antibody Responses Against Mycobacterium Tuberculosis. *EMBO Mol Med* (2016) 8:1325–39. doi: 10.15252/emmm.201606330
10. Choreño-Parra JA, Weinstein LI, Yunis EJ, Zúñiga J, Hernández-Pando R. Thinking Outside the Box: Innate-and B Cell-Memory Responses as Novel Protective Mechanisms Against Tuberculosis. *Front Immunol* (2020) 11:226. doi: 10.3389/fimmu.2020.00226
11. Fukuda T, Matsumura T, Ato M, Hamasaki M, Nishiuchi Y, Murakami Y, et al. Critical Roles for Lipomannan and Lipoarabinomannan in Cell Wall Integrity of Mycobacteria and Pathogenesis of Tuberculosis. *MBio* (2013) 4:1–11. doi: 10.1128/mBio.00472-12
12. Mishra AK, Driessen NN, Appelmelk BJ, Besra GS. Lipoarabinomannan and Related Glycoconjugates: Structure, Biogenesis and Role in Mycobacterium Tuberculosis Physiology and Host–Pathogen Interaction. *FEMS Microbiol Rev* (2011) 35:1126–57. doi: 10.1111/j.1574-6976.2011.00276.x
13. Del Rey A, Mahuad CV, Bozza VV, Bogue C, Farroni MA, Bay ML, et al. Endocrine and Cytokine Responses in Humans With Pulmonary Tuberculosis. *Brain Behav Immun* (2007) 21:171–9. doi: 10.1016/j.bbi.2006.06.005
14. Santucci N, D’Attilio L, Kovalevski L, Bozza V, Besedovsky H, Del Rey A, et al. A Multifaceted Analysis of Immune-Endocrine-Metabolic Alterations in Patients With Pulmonary Tuberculosis. *PLoS One* (2011) 6:e26363. doi: 10.1371/journal.pone.0026363
15. Kroboth PD, Salek FS, Pittenger AL, Fabian TJ, Frye RF. DHEA and DHEA-s: A Review. *J Clin Pharmacol* (1999) 39:327–48. doi: 10.1177/00912709922007903
16. Miyaura H, Iwata M. Direct and Indirect Inhibition of Th1 Development by Progesterone and Glucocorticoids. *J Immunol* (2002) 168:1087–94. doi: 10.4049/jimmunol.168.3.1087
17. Collins HL, Kaufmann SH. The Many Faces of Host Responses to Tuberculosis. *Immunology* (2001) 103:1. doi: 10.1046/j.1365-2567.2001.01236.x
18. Pérez AR, Bottasso O, Savino W. The Impact of Infectious Diseases Upon Neuroendocrine Circuits. *Neuroimmunomodulation* (2009) 16:96–105. doi: 10.1159/000180264
19. Ribeiro F, Lopes RP, Nunes CP, Maito F, Bonorino C, Bauer ME. Dehydroepiandrosterone Sulphate Enhances Igg and Interferon-Gamma Production During Immunization to Tuberculosis in Young But Not Aged Mice. *Biogerontology* (2007) 8:209–20. doi: 10.1007/s10522-006-9069-z
20. Santucci N, D’Attilio L, Besedovsky H, Del Rey A, Bay ML, Bottasso O. A Clinical Correlate of the Dysregulated Immunoendocrine Response in Human Tuberculosis. *Neuroimmunomodulation* (2010) 17:184–7. doi: 10.1159/000258719
21. Yurt S, Erman H, Korkmaz G, Kosar A, Uysal P, Gelisgen R, et al. The Role of Feed Regulating Peptides on Weight Loss in Patients With Pulmonary Tuberculosis. *Clin Biochem* (2013) 46:40–4. doi: 10.1016/j.clinbiochem.2012.09.008
22. Kim JH, Lee C-T, Yoon HI, Song J, Shin WG, Lee JH. Relation of Ghrelin, Leptin and Inflammatory Markers to Nutritional Status in Active Pulmonary Tuberculosis. *Clin Nutr* (2010) 29:512–8. doi: 10.1016/j.clnu.2010.01.008
23. Yüksel İ, Şencan M, Sebila Dökmetaş H, Dökmetaş İ, Ataseven H, Yönm Ö. The Relation Between Serum Leptin Levels and Body Fat Mass in Patients With Active Lung Tuberculosis. *Endocrine Res* (2003) 29:257–64. doi: 10.1081/ERC-120025033
24. Matarese G, Moschos S, Mantzoros CS. Leptin in Immunology. *J Immunol* (2005) 174:3137–42. doi: 10.4049/jimmunol.174.6.3137
25. Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR, et al. Serum Immunoreactive-Leptin Concentrations in Normal-Weight and Obese Humans. *N Engl J Med* (1996) 334:292–5. doi: 10.1056/NEJM199602013340503
26. Bini EI, D’Attilio L, Marquina-Castillo B, Mata-Espinosa D, Díaz A, Marquez-Velasco R, et al. The Implication of Pro-Inflammatory Cytokines in the Impaired Production of Gonadal Androgens by Patients With Pulmonary Tuberculosis. *Tuberculosis* (2015) 95:701–6. doi: 10.1016/j.tube.2015.06.002
27. Dorshkind K, Horseman ND. The Roles of Prolactin, Growth Hormone, Insulin-Like Growth Factor-I, and Thyroid Hormones in Lymphocyte Development and Function: Insights From Genetic Models of Hormone and Hormone Receptor Deficiency. *Endocrine Rev* (2000) 21:292–312. doi: 10.1210/edrv.21.3.0397
28. Kimuda SG, Nalwoga A, Levin J, Franken KL, Ottenhoff TH, Elliott AM, et al. Humoral Responses to Rv1733c, Rv0081, Rv1735c, and Rv1737c Dsr Regulon-Encoded Proteins of Mycobacterium Tuberculosis in Individuals With Latent Tuberculosis Infection. *J Immunol Res* (2017) 2017:1–8. doi: 10.1155/2017/1593143
29. Wang S, Wu J, Chen J, Gao Y, Zhang S, Zhou Z, et al. Evaluation of Mycobacterium Tuberculosis-Specific Antibody Responses for the Discrimination of Active and Latent Tuberculosis Infection. *Int J Infect Dis* (2018) 70:1–9. doi: 10.1016/j.ijid.2018.01.007
30. Da Costa C, Khanolkar-Young S, Elliott A, Wasunna K, McAdam K. Immunoglobulin G Subclass Responses to Mycobacterial Lipoarabinomannan in HIV-Infected and Non-Infected Patients With Tuberculosis. *Clin Exp Immunol* (1993) 91:25–9. doi: 10.1111/j.1365-2249.1993.tb03348.x
31. Baumann R, Kaempfer S, Chegou NN, Oehlmann W, Spallek R, Loxton AG, et al. A Subgroup of Latently Mycobacterium Tuberculosis Infected Individuals is Characterized by Consistently Elevated Iga Responses to Several Mycobacterial Antigens. *Mediators Inflamm* (2015) 2015:1–10. doi: 10.1155/2015/364758
32. Legesse M, Ameni G, Medhin G, Mamo G, Franken K, Ottenhoff T, et al. Iga Response to ESAT-6/CFP-10 and R V2031 Antigens Varies in Patients With Culture-Confirmed Pulmonary Tuberculosis, Healthy M Ycobacterium Tuberculosis-Infected and non-Infected Individuals in a Tuberculosis Endemic Setting, E Thiopia. *Scand J Immunol* (2013) 78:266–74. doi: 10.1111/sji.12080
33. Abebe F, Belay M, Legesse M, Franken KLMC, Ottenhoff THM. Iga and IgG Against Mycobacterium Tuberculosis Rv2031 Discriminate Between Pulmonary Tuberculosis Patients, Mycobacterium Tuberculosis-Infected and Non-Infected Individuals. *PLoS One* (2018) 13(1):e0190989. doi: 10.1371/journal.pone.0190989
34. Nahas AA, Lima Mid.S, Goulart IMB, Goulart LR. Anti-Lipoarabinomannan-Specific Salivary Iga as Prognostic Marker for Leprosy Reactions in Patients and Cellular Immunity in Contacts. *Front Immunol* (2018) 9:1205. doi: 10.3389/fimmu.2018.01205
35. Hussain R, Kifayet A, Chiang TJ. Immunoglobulin G1 (Igg1) and Igg3 Antibodies are Markers of Progressive Disease in Leprosy. *Infect Immun* (1995) 63:410–5. doi: 10.1128/iai.63.2.410-415.1995

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Tsegaye, Admassu, Edao, Kinde, Gentu, Negash, Wondmagegn, Gize, Zewdie, Bobosha and Wassie. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.