



Soluble P2X7 Receptor Is Elevated in the Plasma of COVID-19 Patients and Correlates With Disease Severity

Julio García-Villalba^{1†}, Laura Hurtado-Navarro^{1†}, Alejandro Peñín-Franch¹, Cristina Molina-López¹, Laura Martínez-Alarcón¹, Diego Angosto-Bazarra¹, Alberto Baroja-Mazo¹ and Pablo Pelegrin^{1,2*}

¹ Biomedical Research Institute of Murcia (IMB-Arixaca), University Clinical Hospital Virgen Arrixaca, Murcia, Spain,

² Department of Biochemistry and Molecular Biology B and Immunology, Faculty of Medicine, University of Murcia, Murcia, Spain

OPEN ACCESS

Edited by:

Jianguo Wu,
Wuhan University, China

Reviewed by:

Talia H. Swartz,
Icahn School of Medicine at Mount
Sinai, United States
Anna Lisa Giuliani,
University of Ferrara, Italy

*Correspondence:

Pablo Pelegrin
pablo.pelegrin@imib.es

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Inflammation,
a section of the journal
Frontiers in Immunology

Received: 11 March 2022

Accepted: 19 April 2022

Published: 18 May 2022

Citation:

García-Villalba J, Hurtado-Navarro L,
Peñín-Franch A, Molina-López C,
Martínez-Alarcón L,
Angosto-Bazarra D, Baroja-Mazo A
and Pelegrin P (2022) Soluble P2X7
Receptor Is Elevated in the
Plasma of COVID-19 Patients and
Correlates With Disease Severity.
Front. Immunol. 13:894470.
doi: 10.3389/fimmu.2022.894470

Inflammation is a tightly coordinated response against bacterial and viral infections, triggered by the production of pro-inflammatory cytokines. SARS-CoV-2 infection induces COVID-19 disease, characterized by an inflammatory response mediated through the activation of the NLRP3 inflammasome, which results in the production of IL-1 β and IL-18 along with pyroptotic cell death. The NLRP3 inflammasome could be also activated by sterile danger signals such as extracellular ATP triggering the purinergic P2X7 receptor. Severe inflammation in the lungs of SARS-CoV-2-infected individuals is associated with pneumonia, hypoxia and acute respiratory distress syndrome, these being the causes of death associated with COVID-19. Both the P2X7 receptor and NLRP3 have been considered as potential pharmacological targets for treating inflammation in COVID-19. However, there is no experimental evidence of the involvement of the P2X7 receptor during COVID-19 disease. In the present study, we determined the concentration of different cytokines and the P2X7 receptor in the plasma of COVID-19 patients and found that along with the increase in IL-6, IL-18 and the IL-1 receptor antagonist in the plasma of COVID-19 patients, there was also an increase in the purinergic P2X7 receptor. The increase in COVID-19 severity and C-reactive protein concentration positively correlated with increased concentration of the P2X7 receptor in the plasma, but not with the IL-18 cytokine. The P2X7 receptor was found in the supernatant of human peripheral blood mononuclear cells after inflammasome activation. Therefore, our data suggest that determining the levels of the P2X7 receptor in the plasma could be a novel biomarker of COVID-19 severity.

Keywords: COVID-19, SARS-CoV-2, inflammation, P2X7, purinergic receptors, pyroptosis

INTRODUCTION

Coronavirus disease 2019 (COVID-19) develops after infection with the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and according to the Johns Hopkins University COVID-19 dashboard, as of 10 February 2022 there have been over 400 million infections and 5.7 million deaths worldwide (coronavirus.jhu.edu/map.html). SARS-CoV-2 primarily infects cells in the respiratory

tract and causes different degrees of symptomatology, with the severe cases developing into pneumonia with hypoxia and acute respiratory distress syndrome (ARDS) (1). The lungs of COVID-19 patients with ARDS show considerable infiltration of neutrophils and monocytes and an increase in the concentration of several pro-inflammatory cytokines in the blood (1–6). The nucleotide-binding oligomerization domain, leucine-rich repeat and pyrin domain-containing-protein 3 (NLRP3) inflammasome controls the release of bioactive interleukin (IL)-1 β and IL-18 cytokines by activating caspase-1 and inducing pyroptosis cell death (7, 8). Caspase-1 also cleaves gasdermin D (GSDMD) that induces pores in the plasma membrane allowing the release of IL-1 β and IL-18 (7, 9, 10). The NLRP3 inflammasome is activated by different signals, including triggering of the P2X7 receptor by extracellular ATP, the phagocytosis of particulate matter and infection with specific viruses or bacteria (11–13). The SARS-CoV-2 virus activates NLRP3 (14–18), and this inflammasome has been implicated in the production of pro-inflammatory cytokines in severe cases of COVID-19 and is thus directly associated with the severity of ARDS (15, 19, 20). A clinical trial in COVID-19 patients with anakinra (the recombinant IL-1 receptor antagonist) has found that it decreases the severity of the respiratory failure by reducing the inflammatory response (21). Mechanistically, the SARS-CoV-2 viroporin encoded by *ORF3a* triggers cellular K⁺ efflux and activates NLRP3 (22), with decreased intracellular K⁺ being a well-known activator of the NLRP3 inflammasome (23). Also, the SARS-CoV-2 N protein and the *ORF8*-encoded protein are able to interact with NLRP3 and induce its activation (24, 25). On the one hand, during SARS-CoV-2 infection there is a reprogramming of the host macrophages, which means that they can be primed for the NLRP3 inflammasome by the virus spike (S) protein (26). On the other hand, the SARS-CoV-2 virus blocks the inflammasome pathway at different levels; for example, the envelope (E) protein impairs NLRP3 priming (27), the non-structural proteins 1 and 13 suppress the activation of the NLRP3 inflammasome (28) and the N protein cleaves GSDMD within the N-terminal lytic domain, thus inactivating pyroptosis (29). Therefore, there is a fine balance between positive and negative routes during NLRP3 inflammasome activation in cases of SARS-CoV-2 infection. Different review studies also suggest that the P2X7 receptor could be an important molecular pathway activated during the COVID-19 disease, partly due to the activation of the NLRP3 inflammasome (30, 31). Activation of P2X7 also leads to the cellular shedding of this receptor (32) and, during different inflammatory diseases, an increase in the concentration of a soluble form of the receptor in the plasma (32–34). However, in SARS-CoV-2 infection there is no experimental evidence regarding the potential involvement of the P2X7 receptor.

In the present study, we have found that during active COVID-19 disease there are elevated concentrations in the plasma of IL-18, the IL-1 receptor antagonist (IL-1Ra) and a soluble form of the purinergic P2X7 receptor. IL-1Ra and the P2X7 receptor both increased with the severity of COVID-19 symptoms, but only the P2X7 receptor correlated with CRP concentration in the plasma of COVID-19 patients.

RESULTS

Plasma samples were collected from a cohort of 208 individuals with confirmed SARS-CoV-2 infection and different degrees of COVID-19 disease gravity between September 2020 and March 2021 in the Region of Murcia in the southeast region of Spain (**Table 1**). The largest group among these individuals were those with moderate COVID-19 symptoms (N = 128), followed by those with mild symptoms (N = 41) and those with severe symptoms (N = 39). The control cohort used samples that had been collected during the period 2016–2019 (pre-COVID-19 pandemic) also in the Region of Murcia in the southeast region of Spain. These samples were gathered from 69 individuals without SARS-CoV-2 infection and without inflammatory or infection symptoms (**Table 1**). The level of the acute phase C-reactive protein (CRP) was above the standard threshold in the plasma of 90.9% of the patients infected with SARS-CoV-2 (**Figure 1A**). However, procalcitonin (PCT) was above the standard healthy threshold in only 10.6% of the COVID-19 patients (**Figure 1A**), whereas ferritin was above the standard healthy threshold in 71.2% of the COVID-19 patients (taking into account male and female thresholds, **Figure 1A**). As expected, the IL-6 cytokine was also elevated in the plasma of patients infected with SARS-CoV-2 when compared to healthy control individuals (**Figure 1B**), and IL-15, IL-1RA and IL-18 were also elevated (**Figure 1B**), a finding that is in line with previous publications (3, 16). However, other cytokines or chemokines, such as CCL2/MCP-1 and IL-2, were not elevated in the plasma of the cohort of SARS-CoV-2 infected patients when compared to healthy individuals (**Figure 1C**). Increases in these have been associated with mortality only in COVID-19 patients (3), and our cohort confirms that CCL2/MCP-1 was also significantly increased in the COVID-19 patients who die (237.4 ± 12.4 vs. 421.9 ± 59.0 pg/ml CCL2 mean \pm sem in COVID-19 survival vs. exitus; $p = 0.018$ Mann–Whitney test). However, values for IL-2 failed to associate with COVID-19 mortality (not shown), and this could probably be because the IL-2 concentration was very low, close to the detection threshold.

The presence of the soluble P2X7 receptor was significantly increased in the plasma of SARS-CoV-2 patients when compared to healthy individuals (**Figure 1D**). However, other markers of pyroptosis (HMGB1, ASC and GSDMD) were not elevated in the plasma of SARS-CoV-2-infected patients when compared to healthy individuals (**Figure 1D**). Of particular note is that only 27.5% of healthy individuals and 25% of COVID-19 patients presented a positive detection of GSDMD in the plasma, suggesting that GSDMD was under the detection threshold in the majority of individuals and irrespective of the infection. Similarly, HMGB1 was detected in only 58% of healthy donors and in 74% of COVID-19 patients. The detection rates of all the other identified proteins were very similar among healthy donors and COVID-19 patients, with a detection range above 80% of the samples analysed.

The concentration of CRP in the plasma of COVID-19 patients significantly increased in line with the categories of illness severity (**Figure 2A**). Illness severity categories were defined as mild,

TABLE 1 | Demographic and clinical characteristics of the individuals included in this study.

	Healthy donors	Total COVID-19	Mild COVID-19	Moderate COVID-19	Severe COVID-19
N	69	208	41	128	39
Age , mean (range) ± SD	48.26 (20–94) ± 18.44	64.31 (20–92) ± 15.56	72.86 (42–89) ± 14.16	61.20 (25–91) ± 14.16	71.49 (20–92) ± 17.07
<i>p</i> value		<i>p</i> < 0.0001 vs. healthy donors			
Gender , N (%)					
Male	33 (47.8%)	88 (42.3%)	24 (58.5%)	71 (55.5%)	25 (64.1%)
Female	36 (52.2%)	120 (57.7%)	17 (41.5%)	57 (44.5%)	14 (35.9%)
<i>p</i> value		<i>p</i> > 0.05 vs. healthy donors			
Clinical data					
Patients with available clinical data (%)		89.9	80.5	92.9	89.7
Days of hospitalization, mean (range) ± SD		8.123 (1–133) ± 11.07	3.78 (1–13) ± 2.631	8.151 (1–34) ± 6.782	12.11 (1–133) ± 21.68
N (%)		5 (2.4%)	1 (0.5%)	1 (2.44%)	3 (7.7%)
Heart rate ≥125 bpm		20 (9.62%)	1 (0.5%)	16 (39%)	3 (7.7%)
Respiratory rate >24 bpm		42 (20.19%)	5 (2.4%)	29 (70.73%)	8 (20.51%)
SPO ₂ <94%					
Supplemental O ₂ , N (%)		157 (75.5%)	15 (36.5%)	105 (82.1%)	37 (94.8%)
Nasal cannula		98 (47.1%)	36.5%	59.3%	17.9%
Face mask ventilation		44 (21.2%)	–	21.1%	43.5%
Non-invasive mechanical Ventilation/intubation		15 (7.2%)	–	1.6%	33.3%
Critical unit/ICU, N (%)		39 (18.75%)	–	1.6%	94.8%
Clinical lung disease diagnosis to COVID-19, N (%)					
None		15 (7.2%)	21.9%	4.6%	–
Pneumonia		44 (21.2%)	31.7%	14.1%	33.3%
Bilateral pneumonia		52 (25.0%)	14.6%	32.1%	12.8%
Bronchopneumonia		95 (45.7%)	29.2%	49.3%	51.2%
Pulmonary embolism		1 (0.5%)	–	–	2.5%
Atelectasis lamellose		1 (0.5%)	2.4%	–	–
Deaths, N (%)		29 (13.9%)	–	–	74.3%
Comorbidities, N (%)					
Hypertension		98 (47.1%)	41.4%	45.3%	58.9%
Dyslipidaemia		72 (34.6%)	24.3%	33.6%	48.7%
Diabetes		49 (23.6%)	9.7%	26.6%	28.2%
Hypothyroidism		31 (14.9%)	26.8%	7.1%	28.2%
Obesity		32 (15.4%)	12.2%	17.2%	12.8%
Heart disease		40 (19.2%)	26.8%	10.9%	38.4%
Bronchial asthma		18 (8.7%)	2.4%	8.6%	15.4%
Hyperuricemia		9 (4.3%)	4.8%	5.6%	–
Immunocompromised		6 (2.9%)	4.8%	1.6%	5.1%
Hepatitis C		2 (0.9%)	2.4%	0.8%	–
Gilbert's disease		1 (0.5%)	–	0.8%	–
Crohn's disease		1 (0.5%)	–	0.8%	–
Glaucoma		1 (0.5%)	–	0.8%	–
Chagas disease		1 (0.5%)	2.4%	–	–




FIGURE 1 | Inflammatory cytokines and biomarkers are markedly induced in COVID-19 patients. **(A)** Concentration of C-reactive protein (CRP), procalcitonin (PCT) and ferritin in plasma of COVID-19 patients. **(B, C)** Plasma concentrations of IL-6, IL-15, IL-1RA, IL18 **(B)**, CCL2 and IL-2 **(C)** in healthy donors (HD) and COVID-19 patients. **(D)** Plasma concentration of P2X7 receptor and markers of pyroptosis (ASC, GSDMD, and HMGB1) in healthy donors (HD) and COVID-19 patients. Each dot represents an individual patient; data are represented as mean (grey bars) \pm sem. Dotted lines represent the normal range for each parameter analysed (the upper line in Ferritin's graph corresponds to the normal range for healthy men and bottom line for healthy women). For **(B–D)**, **** $p < 0.0001$; *ns*, no significant difference ($p > 0.05$) with the Mann–Whitney test.

moderate and severe/critical in line with the clinical spectrum of SARS-CoV-2 infection defined in the latest COVID-19 treatment guidelines published by the National Institute of Health. A total of 41 mild, 128 moderate and 39 severe COVID-19 cases were identified (**Table 1**). However, PCT was significantly elevated in only the severe COVID-19 group (**Figure 2A**) and ferritin was increased in the mild and severe groups (**Figure 2A**).

The P2X7 receptor and IL-1RA were significantly increased in the plasma of COVID-19 patients across all severity groups when compared to healthy controls (**Figure 2B**). IL-6 and IL-15 were not increased in the mild COVID-19 group but were significantly elevated in the moderate and severe groups (**Figure 2C**). IL-18 was significantly increased only in moderate COVID-19 patients (**Figure 2C**). The increase in the P2X7 receptor in the plasma of




FIGURE 2 | P2X7 receptor and IL-1RA increase with COVID-19 disease severity. **(A)** Plasma concentration of C-reactive protein (CRP), procalcitonin (PCT) and ferritin in association with the severity of COVID-19 manifestation (mild, moderate or severe). **(B, C)** Concentration of P2X7 receptor, IL-1RA **(B)**, IL-6, IL-15 and IL-18 **(C)** in plasma of healthy donors (HD) and COVID-19 patients in association with the severity of disease manifestation (mild, moderate or severe). **(D)** Concentration of P2X7 receptor in the plasma of HD and COVID-19 patients in different age groups. **(E)** Concentration of P2X7 receptor in the plasma of COVID-19 patients with hypothyroidism, asthma, hypertension, diabetes or dyslipidaemia. **(F)** Concentration of P2X7 receptor in the plasma of severe COVID-19 patients with good or poor outcome (death or prolonged intensive care unit stay). Each dot represents an individual patient; data are represented as mean (grey bars) \pm sem; dotted lines represent the normal range for each parameter analysed (the upper line in the Ferritin graph corresponds to normal range for healthy men, and the bottom line corresponds to normal range for healthy women); * $p < 0.05$; ** $p < 0.005$; *** $p < 0.0005$; **** $p < 0.0001$; ns, no significant difference ($p > 0.05$) with the Kruskal–Wallis test for **(A–D)** and Mann–Whitney test for **(E, F)**.

COVID-19 patients was higher in the 60- to 80-year-old group (**Figure 2D**) and in COVID-19 patients with diabetes or dyslipidaemia (**Figure 2E**). However, the P2X7 receptor concentration in the plasma was not affected in COVID-19 patients with hypothyroidism, asthma or hypertension (**Figure 2E**). Furthermore, the P2X7 receptor was significantly increased in the plasma of COVID-19 severe patients who had a poor outcome (death or prolonged intensive care unit stay) (**Figure 2F**) and a multivariate logistic regression analysis showed that an increase in the concentration of soluble P2X7 is associated with severe COVID-19 symptoms (OR = 1.04; [95% CI: 1.013–1.060]; $p = 0.002$). This multivariate analysis also revealed that patients affected by asthma increases 4.4 times of having severe COVID-19 (OR = 4.4; [95% CI: 1.575–12.186]; $p = 0.005$) or age increases 4.2 times of having severe COVID-19 (OR = 4.2; [95% CI: 1.597–11.139]; $p = 0.004$).

The amount of soluble P2X7 receptor and the concentration of the cytokines IL-6 and IL-15 positively correlated with the amount

of CRP in the plasma (**Figures 3A, B**). The positive correlation between the P2X7 receptor and CRP increased in the moderate and severe COVID-19 groups, but in contrast, the positive IL-6 and IL-15 correlation with CRP decreased in these groups (**Figure 3C**). Neither IL-1RA nor IL-18 cytokines correlated with CRP when all COVID-19 patients were analysed (**Figures 3A, B**) or when the moderate and severe groups were separated (not shown). Although CRP also positively correlated with ferritin concentration (**Figure 3D**), ferritin had a weaker correlation with the P2X7 receptor and no correlation with IL-18 (**Figure 3D**). However, the soluble P2X7 receptor positively correlated with the different cytokines analysed, the best correlation being when the presence of IL-18 increased in the plasma (**Figure 3E**). This suggests that the soluble P2X7 receptor increases with the severity of COVID-19 illness.

P2X7 receptor release has been observed in human macrophages after treatment with the semi-selective P2X7 receptor agonist




FIGURE 3 | P2X7 receptor detected in the plasma of COVID-19 patients correlates with the C-reactive protein. **(A, B)** Correlation between the concentration of C-reactive protein (CRP) and the quantification of the P2X7 receptor, IL-1RA **(A)**, IL-6, IL-15 and IL-18 **(B)** in plasma from COVID-19 patients; disease severity is represented with yellow circles (mild), orange circles (moderate) and red circles (severe). **(C)** Correlation between the concentration of CRP and the quantification of the P2X7 receptor, IL-6 and IL-15 in plasma from COVID-19 patients with moderate (orange circles) and severe (red circles) COVID-19 manifestation. **(D)** Correlation between the concentration of ferritin and CRP, P2X7 receptor and IL-18 in plasma from COVID-19 patients; disease severity is represented with yellow circles (mild), orange circles (moderate) and red circles (severe). **(E)** Correlation between the concentration of the P2X7 receptor and IL-1RA, IL-6, IL-15 and IL-18 from plasma of COVID-19 patients; disease severity is represented with yellow circles (mild), orange circles (moderate) and red circles (severe). Each dot represents an individual patient; Spearman correlation for all panels.

benzoyl ATP (32); we therefore asked if the ATP treatment of human peripheral blood mononuclear cells (PBMCs) would also result in the shedding of the P2X7 receptor. Our results showed that ATP was able to increase the concentration of the P2X7 receptor found in the supernatant of LPS-treated PBMCs (**Figure 4**). This LPS and ATP treatment also induced the activation of the NLRP3 inflammasome (11), as was evidenced by the release of IL-1 β (**Figure 4**). Furthermore, the P2X7 receptor and IL-1 β were also released after the activation of a P2X7 receptor-independent inflammasome, such as the pyrin inflammasome (**Figure 4**). The pyrin inflammasome is activated by toxins that block RhoA GTPase, such as *Clostridium difficile* enterotoxin B (TcdB). RhoA blockage leads to the dephosphorylation of pyrin and liberates it from the inhibitory 14–3–3 protein, which in turn leads to the activation of the pyrin inflammasome (35). These results support the notion that inflammasome-dependent caspase-1 activation is linked to the shedding of the P2X7 receptor from human blood cells, and given the inflammasome is activated during COVID-19 infection (14–16, 18, 20, 26), this could be the cause of the elevated levels of the P2X7 receptor in the plasma of these patients.

DISCUSSION

Severe COVID-19 symptoms are related to an exacerbated inflammatory response mediated by the production of several pro-inflammatory cytokines (1, 3). In the present study, we corroborate in a cohort of COVID-19 patients from the southeast of Spain an elevated concentration of the cytokines IL-6, IL-15, IL-18 and IL-1RA. Together with these cytokines, we found that levels of the purinergic receptor P2X7 also increase in the plasma of symptomatic COVID-19 patients, that these levels increase as the severity of the disease increases and that they correlate with the levels of the acute phase protein CRP, different cytokines and poor outcome. However, our cohort present a relatively low mortality and

association of the concentration of P2X7 receptor in the plasma with mortality would require further validation in larger cohorts. The activation of the P2X7 receptor leads to the activation of the NLRP3 inflammasome, the release of the IL-1 β and IL-18 cytokines (11, 13), and the shedding of the P2X7 receptor in microparticles (32). Soluble P2X7 receptor found in cell supernatants is therefore associated with extracellular vesicles (32), and is therefore expected to be similar to the cellular plasma-membrane bound receptor. However, there is no data on the potential function of this shed form of the P2X7 receptor in extracellular vesicles, although circulatory exosomes from COVID-19 patients are able to activate the NLRP3 inflammasome (36). In the present study, we show that human peripheral blood mononuclear cells release the P2X7 receptor when treated with ATP. This suggests that potential NLRP3 inflammasome activation and subsequent pyroptosis could be partially responsible for the release of microparticles loaded with P2X7 receptor, because the endosomal sorting complexes required for transport (ESCRT) machinery induces the shedding of GSDMD-damaged membranes (7, 37). In support of this hypothesis, we also describe in this study how the activation of the Pyrin inflammasome by TcdB and the subsequent pyroptosis (which is independent of the P2X7 receptor) also leads to the release of the P2X7 receptor. Increased concentrations of soluble P2X7 receptor in the blood have been described as positively correlating with the concentration of CRP (32, 34), with the soluble P2X7 receptor being elevated during infection and sepsis (32, 33), and during temporal lobe epilepsy [34]. Our results agree with those previous studies, and describe how infection with SARS-CoV-2 also increases the soluble P2X7 receptor in the plasma, which positively correlates with CRP concentration. In fact, recent studies have reported the activation of the inflammasome during SARS-CoV-2 infection and relate it to the severity of the COVID-19 infection (18). All this evidence suggests that detecting the P2X7 receptor could be a promising novel blood-biomarker for inflammatory processes and help diagnose and determine the severity of




FIGURE 4 | Treatment with ATP or *Clostridium difficile* toxin B induces the release of the P2X7 receptor. P2X7 receptor (left) or IL-1 β (right) from human peripheral blood mononuclear cells from healthy individuals treated for 3 h with LPS (1.6 μ g/ml) and then treated with ATP (5 mM) for 45 min or with *Clostridium difficile* toxin B (TcdB, 1 μ g/ml) for 1 h. Each dot represents data from an independent individual.

COVID-19. The increase in the P2X7 receptor in the plasma of COVID-19 patients was higher in those with diabetes and dyslipidaemia, which in turn associates the P2X7 receptor with metabolic diseases (38, 39). In addition, targeting the P2X7 receptor with drug-like pharmacological antagonists has been suggested as a promising strategy for treating severe-inflammatory and neurological complications in COVID-19 patients (30, 40).

The concentration of IL-6 has been clinically used to evaluate the severity of COVID-19 (3). Also, IL-6 has been used as a therapeutic target, and IL-6-blocking antibodies have been widely used to treat COVID-19 patients (41). IL-6 is a cytokine produced downstream from IL-1 β , whereas IL-1RA is a cytokine that blocks IL-1 signalling as it binds to the type I IL-1 receptor and prevents IL-1 from triggering its receptor (42). In diseases satisfactorily treated with anti-IL-1 therapy, such as sepsis or autoinflammatory syndromes, the blood concentrations of IL-1 β are very low and usually not detected (33, 43). According to previous publications and despite the evidence of inflammasome activation during COVID-19, IL-1 β has not been detected in the blood samples of COVID-19 patients (5, 16). In this regard, detecting IL-1RA has been proposed as a systemic indicator of IL-1 β production, and our study, in agreement with previous studies (4, 16), shows an important increase in IL-1RA in the plasma of COVID-19 patients. These results suggest that there is potential production of IL-1 β during COVID-19 infection (6).

While the axes IL-1 β /IL-6 link to CRP production in bacterial infections, IL-18 production links to transferrin in viral infection (44). However, in the plasma of SARS-CoV-2-infected patients we found increased levels of both axes, IL-6/CRP and IL-18/ferritin; however, although IL-6 positively correlated with CRP, IL-18 failed to correlate with ferritin in COVID-19 patients. P2X7 correlated with CRP, a finding supported by previous studies (32, 34), but not with ferritin, denoting that the shedding of the P2X7 receptor is somehow related to the IL-6/CRP inflammatory programme in both bacterial (45) and SARS-CoV-2 viral (this study) infections.

In conclusion, our study shows that the soluble P2X7 receptor concentration increases in the plasma of COVID-19 and positively correlates with disease severity and CRP protein concentration. This correlation was not found for IL-18 or other cytokines, although these were found to increase in the plasma of COVID-19 patients. Furthermore, the P2X7 receptor was released from human PBMCs upon inflammasome activation, suggesting that the P2X7 receptor could be a novel blood biomarker for COVID-19 severity with poor outcome.

MATERIAL AND METHODS

Human Samples

Samples and data from patients included in this study, who gave written informed consent, were collected, processed and provided by the *Biobanco en Red de la Región de Murcia*, BIOBANC-MUR, were registered on the *Registro Nacional de Biobancos* with registration number B.0000859 and were processed following standard operating procedures with the appropriate approval of the Ethical Committees (2021-5-10-HCUVA). The study included 208 recovered individuals who had previously given positive RT-

PCR tests, positive serology and/or positive rapid antigen tests for SARS-CoV-2. The plasma samples were collected in the days following hospital admission or diagnosis of SARS-CoV-2 infection. For research purposes across studies, patients with COVID-19 were divided according to the clinical spectrum of SARS-CoV-2 infection into three categories of disease severity, namely mild, moderate and severe/critical, as described in the latest National Institutes of Health COVID-19 treatment guidelines (www.covid19treatmentguidelines.nih.gov, accessed 11/25/2021). Mild illness is attributed to hospitalized patients who have any of the various symptoms associated with COVID-19 (such as headache, nausea, sore throat, fever, cough, loss of taste and smell, malaise or muscle pain) but who have no shortness of breath, dyspnoea, oxygen supplementation or abnormal chest imaging. Moderate illness is attributed to hospitalized patients who show evidence of lower respiratory disease during clinical assessment or chest radiograph and who have an oxygen saturation (SpO₂) \geq 94% on room air at sea level requiring, in some cases, oxygen therapy. Severe/critical illness is attributed to hospitalized patients in the Critical Care Unit who have SpO₂ <94% on room air at sea level, a respiratory rate >24 breaths/min, highest level of supplemental oxygen required, bilateral infiltrates and individuals who die from respiratory failure caused by COVID-19 disease, septic shock and/or organ dysfunction. The degree of oxygen supplementation increases depending on the severity of the disease, although the oxygen needs vary depending on the degree of hypoxia in each patient. The negative control group consisted of 69 healthy blood donors from whom plasma samples were collected before the COVID-19 pandemic, meaning that they were free of COVID-19. Sera were stored at -80°C until use. Mononuclear cells from healthy donors were purified after the donors had signed their informed consent agreement, and whole peripheral blood samples were collected ($n = 3$).

Reagents

The different reagents used in this assay and their sources were ultrapure *Escherichia coli* lipopolysaccharide (LPS) serotype 0111:B4 (InvivoGen, San Diego, CA, USA), adenosine 5'-triphosphate (ATP) from Sigma-Aldrich (St. Louis, MO, USA) and *Clostridium difficile* toxin B (TcdB) from Enzo Life Sciences (Farmingdale, NY, USA).

PBMC Stimulation

Human peripheral blood mononuclear cells were collected using Ficoll Histopaque-1077 (Sigma-Aldrich) and cultured in Opti-MEM Reduced Serum Media (Invitrogen, Carlsbad, CA, USA). PBMCs from patients were left unstimulated or stimulated with 1.6 μ g/ml of LPS at 37°C and then subsequently stimulated with ATP 5 mM for 45 min or TcdB 1 μ g/ml for 1 h.

ELISA

Individual culture cell-free supernatants were collected and clarified by centrifugation. The concentration of human soluble P2X7 was tested by ELISA following the manufacturer's instructions (Cusabio, Houston, TX, USA). Plasma levels of human ASC, HMGB1, GSDMD and P2X7 were also tested by ELISA (Cusabio for ASC and P2X7, Aviva Systems Biology (San Diego, CA, USA) for

GSDMD and Arigo Biolaboratories (Hsinchu, Taiwan) for HMGB1). Results were read in a Synergy Mx (BioTek, Winooski, VT, USA) plate reader at 450 nm and corrected at 540 nm.

Multiplex

Multiplexing in human serum for IL-15, IL-18, MCP-1, IL-2, IL-6 and IL-1RA was performed using the Luminex color-coded antibody-immobilized beads from Merck Millipore (Burlington, MA, USA) following the manufacturer's indications, and the results were analysed in a Luminex MAGPIX instrument (Luminex Corporation, Austin, TX, USA).

Statistics and Data Analysis

Statistical analyses were performed using GraphPad Prism 9 (GraphPad Software Inc., La Jolla, CA, USA). Normality of the samples was determined using the D'Agostino and Pearson omnibus K2 normality test. Outliers from data sets were identified by the ROUT method with $Q = 1\%$ and were eliminated from the analysis and representation. The non-parametric Mann–Whitney test was used to compare differences among two non-paired groups, and the Kruskal–Wallis test was used to compare differences between three or more groups. The χ^2 test was used to determine whether there was a significant difference between clinical variables among groups of patients. Non-parametric correlations were made according to the Spearman correlation coefficient. All data are shown as mean values, and error bars represent standard error. To identify risk factors associated with SARS-CoV-2, severity multivariable analysis was done with SPSS software version 23 (SPSS Inc., Chicago, IL, USA). Age (groups of 20–60 and 60–95 years old), hypothyroidism, asthma, hypertension, diabetes and dyslipidaemia were reported as categorical variables, and the concentration of the soluble P2X7 receptor in the plasma was reported as a continuous variable. We performed a multivariable logistic regression with those variables with a significant difference compared using the χ^2 test, with SARS-CoV-2 severity (mild and moderate, vs. severe) as the dependent variable. The R^2 Cox coefficient was used to assess the proportion of variation explained by the variables studied. For this context, an R^2 Cox coefficient of 0 explains that there is no relationship between the variables and the variability in the data and, whereas an R^2 Cox coefficient of 1 suggests that the variables in the regression explain 100% of the variability in the data.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

REFERENCES

- Zhang Q, Bastard P. COVID Human Genetic Effort, Cobat A, Casanova JL, et al. Human Genetic and Immunological Determinants of Critical COVID-19 Pneumonia. *Nature* (2022) 603:587–98. doi: 10.1038/s41586-022-04447-0
- Guo Y-R, Cao QD, Hong ZS, Tan YY, Chen SD, Jin HJ, et al. The Origin, Transmission and Clinical Therapies on Coronavirus Disease 2019 (COVID-19) Outbreak – an Update on the Status. *Mil Med Res* (2020) 7:11. doi: 10.1186/s40779-020-00240-0

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Comité de Ética en Investigación con Medicamentos, Hospital Clínico Universitario Virgen de la Arrixaca. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

JG-V, LH-N, AP-F and CM-L executed the experiment. JG-V, LH-N and LM-A collected the samples. JG-V, LH-N, AP-F, CM-L, DA-B, AB-M and PP analysed the data. AB-M, DA-B and PP conceived the experiments. JG-V, LH-N and PP prepared the figures and wrote the paper. PP provided the funding and overall supervision of the study. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by grants to PP from MCIN/AEI/10.13039/501100011033 (grant PID2020-116709RB-I00), Fundación Séneca (grants 20859/PI/18, 21081/PDC/19 and 0003/COVI/20) the Instituto Salud Carlos III (grant DTS21/00080), the EU Horizon 2020 project PlasticHeal (grant 965196) and European Research Council (grant ERC-2019-PoC 899636). LH-N was supported by the fellowship 21214/FPI/19 (Fundación Séneca, Región de Murcia, Spain), and CM-L was funded by the fellowship PRE2018-086824 (Ministerio Economía y Competitividad).

ACKNOWLEDGMENTS

We are particularly grateful for the generous contribution of the donors and the collaboration of the Biobank Network of the Region of Murcia, BIOBANC-MUR, which is registered on the *Registro Nacional de Biobancos* with registration number B.0000859. BIOBANC-MUR is supported by the *Instituto de Salud Carlos III* (grant PT20/00109), by the *Instituto Murciano de Investigación Biosanitaria* and by the *Consejería de Salud de la Comunidad Autónoma de la Región de Murcia*. We would like to thank all the members of Dr. Pelegrin's laboratory for comments and suggestions.

- Abers MS, Delmonte OM, Ricotta EE, Fintzi J, Fink DL, de Jesus AAA, et al. An Immune-Based Biomarker Signature is Associated With Mortality in COVID-19 Patients. *JCI Insight* (2021) 6:e144455. doi: 10.1172/jci.insight.144455
- Wilson JG, Simpson LJ, Ferreira AM, Rustagi A, Roque J, Asuni A, et al. Cytokine Profile in Plasma of Severe COVID-19 Does Not Differ From ARDS and Sepsis. *JCI Insight* (2020) 5:1–6. doi: 10.1172/jci.insight.140289
- Del Valle DM, Kim-Schulze S, Huang HH, Beckmann ND, Nirenberg S, Wang B, et al. An Inflammatory Cytokine Signature Predicts COVID-19

- Severity and Survival. *Nat Med* (2020) 26:1636–43. doi: 10.1038/s41591-020-1051-9
6. McElvaney OJ, McEvoy NL, McElvaney OF, Carroll TP, Murphy MP, Dunlea DM, et al. Characterization of the Inflammatory Response to Severe COVID-19 Illness. *Am J Respir Crit Care Med* (2020) 202:812–21. doi: 10.1164/rccm.202005-1583OC
 7. Broz P, Pelegrin P, Shao F. The Gasdermins, a Protein Family Executing Cell Death and Inflammation. *Nat Rev Immunol* (2020) 20:143–57. doi: 10.1038/s41577-019-0228-2
 8. Swanson KV, Deng M, Ting JP. The NLRP3 Inflammasome: Molecular Activation and Regulation to Therapeutics. *Nat Rev Immunol* (2019) 19:477–89. doi: 10.1038/s41577-019-0165-0
 9. Evavold CL, Ruan J, Tan Y, Xia S, Wu H, Kagan JC. The Pore-Forming Protein Gasdermin D Regulates Interleukin-1 Secretion From Living Macrophages. *Immunity* (2018) 48:35–44.e6. doi: 10.1016/j.immuni.2017.11.013
 10. Xia S, Zhang Z, Magupalli VG, Pablo JL, Dong Y, Vora SM, et al. Gasdermin D Pore Structure Reveals Preferential Release of Mature Interleukin-1. *Nature* (2021) 593:607–11. doi: 10.1038/s41586-021-03478-3
 11. Di Virgilio F, Dal Ben D, Sarti AC, Giuliani AL, Falzoni S. The P2X7 Receptor in Infection and Inflammation. *Immunity* (2017) 47:15–31. doi: 10.1016/j.immuni.2017.06.020
 12. Christgen S, Place DE, Kanneganti TD. Toward Targeting Inflammasomes: Insights Into Their Regulation and Activation. *Cell Res* (2020) 30:315–27. doi: 10.1038/s41422-020-0295-8
 13. Pelegrin P. P2X7 Receptor and the NLRP3 Inflammasome: Partners in Crime. *Biochem Pharmacol* (2021) 187:114385. doi: 10.1016/j.bcp.2020.114385
 14. Kroemer A, Khan K, Plassmeyer M, Alpan O, Haseeb MA, Gupta R, et al. Inflammasome Activation and Pyroptosis in Lymphopenic Liver Patients With COVID-19. *J Hepatol* (2020) 73:1258–62. doi: 10.1016/j.jhep.2020.06.034
 15. Rodrigues TS, de Sá KSG, Ishimoto AY, Becerra A, Oliveira S, Almeida L, et al. Inflammasomes are Activated in Response to SARS-CoV-2 Infection and are Associated With COVID-19 Severity in Patients. *J Exp Med* (2021) 218:e20201707. doi: 10.1084/jem.20201707
 16. Junqueira C, Crespo Á, Ranjbar S, Lewandrowski M, Ingber J, de Lacerda LB, et al. FcγR-Mediated SARS-CoV-2 Infection of Monocytes Activates Inflammation. *Nature* (2022). doi: 10.1038/s41586-022-04702-4
 17. Kucia M, Ratajczak J, Bujko K, Adamiak M, Ciechanowicz A, Chumak V, et al. An Evidence That SARS-Cov-2/COVID-19 Spike Protein (SP) Damages Hematopoietic Stem/Progenitor Cells in the Mechanism of Pyroptosis in Nlrp3 Inflammasome-Dependent Manner. *Leukemia* (2021) 35:3026–29. doi: 10.1038/s41375-021-01332-z
 18. Vora SM, Lieberman J, Wu H. Inflammasome Activation at the Crux of Severe COVID-19. *Nat Rev Immunol* (2021) 21:694–703. doi: 10.1038/s41577-021-00588-x
 19. Amin S, Aktar S, Rahman MM, Chowdhury MMH. NLRP3 Inflammasome Activation in COVID-19: An Interlink Between Risk Factors and Disease Severity. *Microbes Infect* (2021) 110:104913. doi: 10.1016/j.micinf.2021.104913
 20. Courjon J, Dufies O, Robert A, Bailly L, Torre C, Chirio D, et al. Heterogeneous NLRP3 Inflammasome Signature in Circulating Myeloid Cells as a Biomarker of COVID-19 Severity. *Blood Adv* (2021) 5:1523–34. doi: 10.1182/bloodadvances.2020003918
 21. Kyriazopoulou E, Panagopoulos P, Metallidis S, Dalekos GN, Poulakou G, Gatselis N, et al. An Open Label Trial of Anakinra to Prevent Respiratory Failure in Covid-19. *Elife* (2021) 10:1–21. doi: 10.7554/eLife.66125
 22. Xu H, Akinyemi IA, Chitre SA, Loeb JC, Lednický JA, McIntosh MT, et al. SARS-CoV-2 Viroprotein Encoded by ORF3a Triggers the NLRP3 Inflammatory Pathway. *Virology* (2022) 568:13–22. doi: 10.1016/j.virol.2022.01.003
 23. Tapia-Abellán A, Angosto-Bazarrá D, Alarcón-Vila C, Baños MC, Hafner-Bratkovič I, Oliva B, et al. Sensing Low Intracellular Potassium by NLRP3 Results in a Stable Open Structure That Promotes Inflammasome Activation. *Sci Adv* (2021) 7:eabf4468. doi: 10.1126/sciadv.abf4468
 24. Wu X, Manske MK, Ruan G, Nowakowski KE, Abeykoon JP, Tang X, et al. Secreted ORF8 Is a Pathogenic Cause of Severe Covid-19 and Potentially Targetable With Select NLRP3 Inhibitors. *bioRxiv* (2021). doi: 10.1101/2021.12.02.470978
 25. Pan P, Shen M, Yu Z, Ge W, Chen K, Tian M, et al. SARS-CoV-2 N Protein Promotes NLRP3 Inflammasome Activation to Induce Hyperinflammation. *Nat Commun* (2021) 12:4664. doi: 10.1038/s41467-021-25015-6
 26. Theobald SJ, Simonis A, Georgomanolis T, Kreer C, Zehner M, Eisfeld HS, et al. Long-Lived Macrophage Reprogramming Drives Spike Protein-Mediated Inflammasome Activation in COVID-19. *EMBO Mol Med* (2021) 13:e14150. doi: 10.15252/emmm.202114150
 27. Yalcinkaya M, Liu W, Islam MN, Kotini AG, Gusarova GA, Fidler TP, et al. Modulation of the NLRP3 Inflammasome by Sars-CoV-2 Envelope Protein. *Sci Rep* (2021) 11:24432. doi: 10.1038/s41598-021-04133-7
 28. Kim N-E, Kim D-K, Song Y-J. SARS-CoV-2 Nonstructural Proteins 1 and 13 Suppress Caspase-1 and the NLRP3 Inflammasome Activation. *Microorganisms* (2021) 9:494. doi: 10.3390/microorganisms9030494
 29. Ma J, Zhu F, Zhao M, Shao F, Yu D, Ma J, et al. SARS-CoV-2 Nucleocapsid Suppresses Host Pyroptosis by Blocking Gasdermin D Cleavage. *EMBO J* (2021) 40:1–17. doi: 10.15252/emboj.2021108249
 30. Di Virgilio F, Tang Y, Sarti AC, Rossato M. A Rationale for Targeting the P2X7 Receptor in Coronavirus Disease 19. *Br J Pharmacol* (2020) 177:4990–4. doi: 10.1111/bph.15138
 31. Hasan D, Shono A, Van Kalken CK, Van derSpek PJ, Krenning EP, Kotani T, et al. A Novel Definition and Treatment of Hyperinflammation in COVID-19 Based on Purinergic Signalling. *Purinergic Signal* (2021) 18:13–59. doi: 10.1007/s11302-021-09814-6
 32. Giuliani AL, Berchan M, Sanz JM, Passaro A, Pizzicotti S, Vultaggio-Poma V, et al. The P2X7 Receptor Is Shed Into Circulation: Correlation With C-Reactive Protein Levels. *Front Immunol* (2019) 10:1–7. doi: 10.3389/fimmu.2019.00793
 33. Martínez-García JJ, Martínez-Banaclocha H, Angosto-Bazarrá D, de Torre-Minguela C, Baroja-Mazo A, Alarcón-Vila C, et al. P2X7 Receptor Induces Mitochondrial Failure in Monocytes and Compromises NLRP3 Inflammasome Activation During Sepsis. *Nat Commun* (2019) 10:2711. doi: 10.1038/s41467-019-10626-x
 34. Conte G, Menéndez-Méndez A, Bauer S, El-Naggag H, Alves M, Nicke A, et al. Circulating P2X7 Receptor Signaling Components as Diagnostic Biomarkers for Temporal Lobe Epilepsy. *Cells* (2021) 10:2444. doi: 10.3390/cells10092444
 35. Masters SL, Lagou V, Jéru I, Baker PJ, Van Eyck L, Parry DA, et al. Familial Autoinflammation With Neutrophilic Dermatitis Reveals a Regulatory Mechanism of Pyrin Activation. *Sci Transl Med* (2016) 8:332ra45. doi: 10.1126/scitranslmed.aaf1471
 36. Sur S, Steele R, Isbell TS, Ray R, Ray RB. Circulatory Exosomes From COVID-19 Patients Trigger NLRP3 Inflammasome in Endothelial Cells. *bioRxiv* (2022). doi: 10.1101/2022.02.03.479081
 37. Rühl S, Shkarina K, Demarco B, Heilig R, Santos JC, Broz P, et al. ESCRT-Dependent Membrane Repair Negatively Regulates Pyroptosis Downstream of GSDMD Activation. *Science* (2018) 362:956–60. doi: 10.1126/science.aar7607
 38. Jain S, Jacobson KA. Purinergic Signaling in Diabetes and Metabolism. *Biochem Pharmacol* (2021) 187:114393. doi: 10.1016/j.bcp.2020.114393
 39. Solini A, Novak I. Role of the P2X7 Receptor in the Pathogenesis of Type 2 Diabetes and its Microvascular Complications. *Curr Opin Pharmacol* (2019) 47:75–81. doi: 10.1016/j.coph.2019.02.009
 40. Ribeiro DE, Oliveira-Giacomelli Á, Glaser T, Arnaud-Sampaio VF, Andrejew R, Dieckmann L, et al. Hyperactivation of P2X7 Receptors as a Culprit of COVID-19 Neuropathology. *Mol Psychiatry* (2021) 26:1044–59. doi: 10.1038/s41380-020-00965-3
 41. Cortegiani A, Ippolito M, Greco M, Granone V, Protti A, Gregoretti C, et al. Rationale and Evidence on the Use of Tocilizumab in COVID-19: A Systematic Review. *Pulmonology* (2021) 27:52–66. doi: 10.1016/j.pulmoe.2020.07.003
 42. Dinarello CA. Interleukin-1. *Cytokine Growth Factor Rev* (1997) 8:253–65. doi: 10.1016/S1359-6101(97)00023-3
 43. Baroja-Mazo A, Martín-Sánchez F, Gómez AI, Martínez CM, Amores-Iniesta J, Compan V, et al. The NLRP3 Inflammasome is Released as a Particulate Danger Signal That Amplifies the Inflammatory Response. *Nat Immunol* (2014) 15:738–48. doi: 10.1038/ni.2919
 44. Slaats J, ten Oever J, van deVeerdonk FL, Netea MG. IL-1β/IL-6/CRP and IL-18/Ferritin: Distinct Inflammatory Programs in Infections. *PLoS Pathog* (2016) 12:1–13. doi: 10.1371/journal.ppat.1005973
 45. Holub M, Lawrence DA, Andersen N, Davidová A, Beran O, Marečková V, et al. Cytokines and Chemokines as Biomarkers of Community-Acquired

Bacterial Infection. *Mediators Inflamm* (2013) 2013:190145. doi: 10.1155/2013/190145

Conflict of Interest: PP is consultant of Glenmark Pharmaceutical, Monte Rosa Trx and Viva *In Vitro* Diagnostics SL. LH-N, LM-A, DA-B, AB-M and PP are cofounders of Viva *In Vitro* Diagnostics SL.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of

the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 García-Villalba, Hurtado-Navarro, Peñín-Franch, Molina-López, Martínez-Alarcón, Angosto-Bazarra, Baroja-Mazo and Pelegrin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.