



OPEN ACCESS

APPROVED BY
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE

Mei Li
✉ mei@igbmc.fr

RECEIVED 06 July 2023

ACCEPTED 07 July 2023

PUBLISHED 18 July 2023

CITATION

Hachem CE, Marschall P, Hener P,
Karnam A, Bonam SR, Meyer P, Flatter E,
Birling M-C, Bayry J and Li M (2023)
Corrigendum: IL-3 produced by T cells is
crucial for basophil extravasation
in hapten- induced allergic
contact dermatitis.
Front. Immunol. 14:1254157.
doi: 10.3389/fimmu.2023.1254157

COPYRIGHT

© 2023 Hachem, Marschall, Hener, Karnam,
Bonam, Meyer, Flatter, Birling, Bayry and Li.
This is an open-access article distributed
under the terms of the [Creative Commons
Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use,
distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Corrigendum: IL-3 produced by T cells is crucial for basophil extravasation in hapten- induced allergic contact dermatitis

Carole El Hachem¹, Pierre Marschall¹, Pierre Hener¹,
Anupama Karnam², Srinivasa Reddy Bonam², Pierre Meyer¹,
Eric Flatter¹, Marie-Christine Birling³, Jagadeesh Bayry^{2,4}
and Mei Li^{1*}

¹Institut de Génétique et de Biologie Moléculaire et Cellulaire, Centre National de la Recherche Scientifique UMR7104, Institut National de la Santé et de la Recherche Médicale U1258, Université de Strasbourg, Illkirch, France, ²Institut National de la Santé et de la Recherche Médicale, Centre de Recherche des Cordeliers, Sorbonne Université, Université de Paris, Paris, France, ³Institut Clinique de la Souris, Illkirch, France, ⁴Department of Biological Sciences & Engineering, Indian Institute of Technology Palakkad, Palakkad, India

KEYWORDS

basophil, IL-3, allergy, skin, extravasation, integrin, retinoic acid

A Corrigendum on

IL-3 produced by T cells is crucial for basophil extravasation in hapten-induced allergic contact dermatitis

By Hachem CE, Marschall P, Hener P, Karnam A, Bonam SR, Meyer P, Flatter E, Birling M-C, Bayry J and Li M (2023) *Front. Immunol.* 14:1151468. doi: 10.3389/fimmu.2023.1151468

In the published article, there was an error in the plasmid number and website link. A correction has been made to **Materials and methods**, *Mice*. This sentence previously stated:

“In order to obtain an Il3 “2 in 1” allele (tm1a, Figure S1), we acquired and modified an IMPC plasmid PG00140_W_3_C02 (<https://www.mousephenotype.org/data/genes/MGI:1923649>)”

The corrected sentence appears below:

“In order to obtain an Il3 “2 in 1” allele (tm1a, Figure S1), we acquired and modified an IMPC plasmid ETPG00275_W_2_F02 (<https://www.mousephenotype.org/data/genes/MGI:96552>)”

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.