



OPEN ACCESS

EDITED BY

Francesco Sabbatino,
University of Salerno, Italy

REVIEWED BY

Mizue Terai,
Thomas Jefferson University, United States
Sarah Coupland,
University of Liverpool, United Kingdom

*CORRESPONDENCE

Klaus Georg Griewank
✉ klaus.griewank@uk-essen.de

[†]These authors have contributed
equally to this work and share
last authorship

RECEIVED 06 February 2024

ACCEPTED 07 May 2024

PUBLISHED 05 June 2024

CITATION

Matull J, Placke J-M, Lodde G, Zaremba A,
Utikal J, Terheyden P, Pföhler C, Herbst R,
Kreuter A, Welzel J, Kretz J, Möller I,
Sucker A, Paschen A, Livingstone E, Zimmer L,
Hadaschik E, Ugurel S, Schadendorf D,
Thielmann CM and Griewank KG (2024)
Clinical and genetic characteristics of *BAP1*-
mutated non-veal and uveal melanoma.
Front. Immunol. 15:1383125.
doi: 10.3389/fimmu.2024.1383125

COPYRIGHT

© 2024 Matull, Placke, Lodde, Zaremba, Utikal,
Terheyden, Pföhler, Herbst, Kreuter, Welzel,
Kretz, Möller, Sucker, Paschen, Livingstone,
Zimmer, Hadaschik, Ugurel, Schadendorf,
Thielmann and Griewank. This is an open-
access article distributed under the terms of
the [Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction
in other forums is permitted, provided the
original author(s) and the copyright owner(s)
are credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Clinical and genetic characteristics of *BAP1*-mutated non-veal and uveal melanoma

Johanna Matull^{1,2}, Jan-Malte Placke¹, Georg Lodde¹,
Anne Zaremba¹, Jochen Utikal^{3,4,5}, Patrick Terheyden⁶,
Claudia Pföhler⁷, Rudolf Herbst⁸, Alexander Kreuter²,
Julia Welzel⁹, Julia Kretz¹, Inga Möller¹, Antje Sucker¹,
Annette Paschen¹, Elisabeth Livingstone¹, Lisa Zimmer¹,
Eva Hadaschik¹, Selma Ugurel¹, Dirk Schadendorf^{1,10,11},
Carl Maximilian Thielmann^{1†} and Klaus Georg Griewank^{1*†}

¹Department of Dermatology, University Hospital Essen, University of Duisburg-Essen, Germany & German Cancer Consortium (Deutsches Konsortium für Translationale Krebsforschung, DKTK), Essen, Germany, ²Department of Dermatology, Venereology and Allergology, Helios St. Elisabeth Hospital Oberhausen, University Witten/Herdecke, Oberhausen, Germany, ³Skin Cancer Unit, German Cancer Research Center (Deutsches Krebsforschungszentrum, DKFZ), Heidelberg, Germany,

⁴Department of Dermatology, Venereology and Allergology, University Medical Center Mannheim, Ruprecht-Karl University of Heidelberg, Heidelberg, Germany, ⁵German Cancer Research Center (Deutsches Krebsforschungszentrum, DKFZ) Hector Cancer Institute at the University Medical Center Mannheim, Mannheim, Germany, ⁶Department of Dermatology, University Hospital Schleswig-Holstein, Lübeck, Germany, ⁷Department of Dermatology, Saarland University Medical School, Homburg, Germany, ⁸Skin Cancer Unit, Helios Klinikum Erfurt, Erfurt, Germany, ⁹Department of Dermatology and Allergology, University Hospital Augsburg, Augsburg, Germany, ¹⁰Comprehensive Cancer Center (Westdeutsches Tumorzentrum), University Hospital Essen, Essen & National Center for Tumor Diseases (NCT) West, Essen, Germany, ¹¹Research Center One Health, University Duisburg-Essen, Essen, Germany

Background: Screening for gene mutations has become routine clinical practice across numerous tumor entities, including melanoma. *BAP1* gene mutations have been identified in various tumor types and acknowledged as a critical event in metastatic uveal melanoma, but their role in non-veal melanoma remains inadequately characterized.

Methods: A retrospective analysis of all melanomas sequenced in our department from 2014–2022 (n=2650) was conducted to identify *BAP1* mutated samples. Assessment of clinical and genetic characteristics was performed as well as correlations with treatment outcome.

Results: *BAP1* mutations were identified in 129 cases and distributed across the entire gene without any apparent hot spots. Inactivating *BAP1* mutations were more prevalent in uveal (55%) compared to non-veal (17%) melanomas. Non-veal *BAP1* mutated melanomas frequently exhibited UV-signature mutations and had a significantly higher mutation load than uveal melanomas. *GNAQ* and *GNA11* mutations were common in uveal melanomas, while MAP-Kinase mutations were frequent in non-veal melanomas with *NF1*, *BRAF* V600 and *NRAS* Q61 mutations occurring in decreasing frequency, consistent with a strong UV association. Survival outcomes did not differ among non-veal melanoma patients based on whether they received targeted or immune checkpoint therapy, or if their tumors harbored inactivating *BAP1* mutations.

Conclusion: In contrast to uveal melanomas, where *BAP1* mutations serve as a significant prognostic indicator of an unfavorable outcome, *BAP1* mutations in non-uveal melanomas are primarily considered passenger mutations and do not appear to be relevant from a prognostic or therapeutic perspective.

KEYWORDS

BAP1, non-uveal melanoma, uveal melanoma, mutation profiling, immunotherapy

1 Introduction

Melanoma, a highly aggressive skin cancer with poor prognosis once metastasized, leads to approximately 55,500 deaths annually worldwide (1). Treatment options for advanced disease were limited for decades, but therapeutic breakthroughs, such as the introduction of immune checkpoint inhibitors (ICI) and targeted therapies (TT), have significantly improved progression-free and overall survival rates. Essential to their development was a better understanding of tumor immunology, genetics, and the widespread use of high-throughput sequencing in clinical routine (2).

Melanoma exhibits one of the highest mutation frequencies among all cancers, with a particularly diverse range of genetic alterations (2, 3). The Cancer Genome Atlas has proposed a genetic classification of melanoma into four subtypes based on mutations in *BRAF*, *NRAS*, *NFI* and triple-wild-type melanomas (4). While some mutations have clear therapeutic implications, such as *BRAF V600E*, the clinical relevance of the majority of identified mutations remains poorly defined.

Mutations in the BRCA-1 associated protein 1 (*BAP1*) gene were recognized as relevant in various cancer types, including uveal melanoma, mesothelioma and renal cell carcinoma. *BAP1* is a ubiquitin carboxy-terminal hydrolase encoded by the *BAP1* gene, located on the short arm of chromosome 3. It was discovered by Jensen and colleagues in 1998 for its ability to bind to BRCA-1 and enhance its tumor suppressive activity (5, 6).

Over the years, *BAP1* has been found to act independently as a tumor suppressor through its de-ubiquinating activity, which regulates target genes involved in transcription, cell cycle control, DNA damage repair, apoptosis, and cell metabolism (7). Germline *BAP1* mutations cause the *BAP1* predisposition syndrome (*BAP1*-TPDS), associated with a high susceptibility to various malignancies, such as uveal melanoma, malignant mesothelioma, cutaneous melanoma, renal cell carcinoma, and other tumors (8).

BAP1 inactivation is strongly linked to a higher metastatic risk and poor prognosis in uveal melanoma, mutated in 84% of metastatic cases (9, 10). However, in non-uveal melanoma, the role of *BAP1* in tumorigenesis and its prognostic significance, particularly in cutaneous melanoma, has been controversial. Low *BAP1* mRNA expression levels were reported to be associated with worse survival in some cutaneous melanoma patient cohorts, while

in others, low *BAP1* mRNA expression was associated with better overall survival (11, 12).

Current research suggests that loss of *BAP1* may have a growth-sustaining effect, making it a potential therapeutic target (13). This study aims to further understand the role of *BAP1* and its implications on clinical course in non-uveal and uveal melanoma by examining a multicenter cohort and correlating clinical and survival information in the respective patients.

2 Materials and methods

2.1 Patient identification

The next-generation sequencing reports from a total of 2650 melanoma patients analyzed at the Department of Dermatology, University Hospital Essen, were reviewed to identify patients harboring *BAP1* mutations (n=129). Of those, 60 tissue samples and related clinical data were obtained from the Westdeutsche Biobank Essen (11–4715-BO), and 69 from the prospective multicenter translational study Tissue Registry in Melanoma (ADOREG/TRIM; NCT05750511; CA209–578; 15–6566-BO) conducted by the German Dermatological Cooperative Oncology Group. Existing data of *BAP1* wildtype melanoma samples (n=1215) were analyzed for comparison of mutational load and mutation types. Tumors were classified as per the American Joint Committee on Cancer (AJCC 8th) staging system (14). Histological evaluation was carried out by local board-certified dermatopathologists. The study was conducted in accordance with the Declaration of Helsinki and was approved by the local ethics committee of the University of Duisburg-Essen (ethics approval no. 21–9873-BO).

2.2 DNA isolation

Formalin-fixed, paraffin-embedded (FFPE) specimens were prepared in 10 µm sections and deparaffinized according to standard procedures. After air-drying, the tumor tissue was manually macrodissected from sections (15). Genomic DNA was isolated applying the QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions.

2.3 Targeted sequencing

Sequencing was performed using a 30-gene custom amplicon-based panel as previously described, covering known melanoma-related gene mutations including *BAP1* (Supplementary Table 1) (16).

To eliminate questionable low frequency background mutation calls, mutations were reported only if ≥ 10 reads reported the mutated variant, coverage of the mutation site was ≥ 30 reads and the frequency of mutated reads was $\geq 10\%$. The average read coverage of the targeted area achieved in the study was 1773x. All samples were sequenced using an Illumina MiSeq and analyzed with the same software (CLC) by the same team over the past eight years. In 2018, there was a transition from PCR-based amplification to an oligo-capture-based technique by Integrated DNA Technologies (IDT).

2.4 Statistical analysis

Associations between covariates were investigated using chi-squared and Fisher's exact tests as indicated. Continuous variables are presented as mean with standard deviation or as median with range, as appropriate. Categorical variables are presented as counts and percentages. Survival data were analyzed using the Kaplan-Meier method with log-rank testing. Progression-free survival (PFS) was calculated from date of systemic treatment initiation to date of progression, or death. Censoring occurred upon change of therapeutic regimen or date of last follow-up.

Overall survival (OS) was calculated from the first date of stage IV diagnosis or start of ICI/TT therapy until death or last patient contact (censored observation), respectively. Tests with *P*-values less than .05 were considered statistically significant. Statistical analyses were performed using Microsoft Excel, GraphPad Prism (version 9), SPSS 27.0 (IBM Corp., Armonk NY, USA), R (R version 4.0.3 (2020–10–10)) and RStudio (17).

3 Results

3.1 Sample cohort

Among a cohort of 2650 melanoma patients, 129 patients harboring a *BAP1* mutation (*BAP1_{mut}*) were identified and included in this study. Of those, 116 (89.9%) cases were categorized as non-veal melanoma (NUM) based on the origin of the primary tumor (cutaneous (n=98), mucosal (n=6), meningeal (n=1), or occult (n=11)). Two additional cases with missing primary location information were considered NUM based on mutational pattern. Eleven (8.5%) cases were of uveal origin.

3.1.1 *BAP1_{mut}* non-veal melanoma

In the non-veal melanoma subgroup (n=118) median age at first diagnosis was 60 years (range 22–82) and 65.3% (n=77) patients were male (Table 1). In patients with cutaneous

melanoma and documented primary (n=45), the most common reported localization was the lower extremity (n=16; 35.6%). Trunk, head and neck and upper extremity were less frequent (n=13; n=12; n=4, respectively).

Of all patients receiving systemic therapy (n=56), anti-PD-1 monotherapy was most frequently administered as first-line treatment (20 cases, 35.7%). CTLA4/PD-1 blockade and BRAF/MEK targeted therapy was less common (13 and 8 cases, respectively). In 15 cases (26.8%) other therapeutic regimens were used including chemotherapy-based regimens, anti-CTLA-4 monotherapy, BRAF inhibitor monotherapy and combination therapy of anti-PD1 and BRAF/MEK inhibitors.

Activating mutations in *BRAF* V600, *NRAS* Q61 or mutations in *NFI* were detected in 32 (27.1%), 31 (26.3%) and 71 (60.2%) samples, respectively. Activating mutations in *GNAQ*/*GNA11* genes were less common with mutations in 3 and 4 samples (2.5% and 3.4%, respectively) (Figure 1A, Supplementary Table 2). *BAP1* mutations were inactivating frameshift or nonsense (hereafter abbreviated and termed "INAC") in 16.9% (n=20) of cases.

3.1.2 *BAP1_{mut}* uveal melanoma

In this subgroup, 5 patients were female and 6 were male. Median age at diagnosis was 65 years (range 43–84) (Table 2). Neither age at first diagnosis nor sex differed significantly between NUM and uveal melanoma subgroup (Table 3).

Combined CTLA4/PD-1 blockade was administered in 3 of 4 cases as first-line non-adjuvant systemic therapy.

No *BRAF*, *NRAS* or *NFI* mutations were detected in uveal melanoma samples. *GNAQ* and *GNA11* mutations were regularly present with mutations in 7 (63.6%) and 3 samples (27.3%), respectively. *BAP1* mutations were detected in all 11 samples with 54.5% inactivating mutations (n=6).

3.2 Targeted next generation sequencing

271 *BAP1* mutations were identified in the 129 examined samples (Supplementary Table 4). Non-veal melanomas frequently harbored more than one *BAP1* mutation (n=54, 45.8%), while only 3 samples of uveal origin (27.3%) harbored two or more (Supplementary Table 5). Mutations in *BAP1* were distributed equally without clustering or hotspots. The primary catalytic domain of *BAP1* protein harbored both inactivating and missense mutations (Figure 2). Uveal melanomas harbored significantly more inactivating (frameshift or nonsense) mutations than non-veal (54.5% and 16.9%, p=0.003).

3.2.1 *BAP1_{mut}* non-veal melanoma

Mutations in other genes were identified in 117 NUM tumor samples (97.5%). *BRAF* mutations were found in more than half of the cases (n=62, 53.4%) with activating V600E and V600K mutations in 32 (27.1%) and 3 samples (2.5%), respectively. *NRAS* mutations were found in 54 samples (45.8%), of which 33 (28.0%) were activating Q61/G12 mutations. *KRAS* mutations were less frequent with 5 activating mutations (4.2, 1 G12V, 3 G12D, 1

TABLE 1 Clinical characteristics of patients with *BAP1*_{mut} non-veal melanoma (n=118).

Variable, n (%)	
Age at first diagnosis, n (%)	
Mean (+/- SD)	60.4 (+/- 15.0)
Range	22 – 82
≤60 years	53 (44.9)
>60 years	65 (55.1)
Sex, n (%)	
Female	41 (34.7)
Male	77 (65.3)
Mutated oncogene, n (%)	
<i>BRAF</i> V600E	32 (27.1)
<i>NRAS</i> Q61	31 (26.3)
<i>NFI</i>	71 (60.2)
<i>GNAQ</i>	3 (2.5)
<i>GNA11</i>	4 (3.4)
<i>BAP1</i>	118 (100)
Primary tumor site, n (%)	
Cutaneous	98 (83.1)
Mucosal	6 (5.1)
Occult	11 (9.3)
Meningeal	1 (0.8)
Unknown	2 (1.7)
Location cutaneous tumor, n (%)*	
Trunk	13 (28.9)
Lower extremity	16 (35.6)
Upper extremity	4 (8.9)
Head and neck	12 (26.7)
Subtype cutaneous tumor, n (%)	
SSM	18 (18.4)
NMM	30 (30.6)
ALM	11 (11.2)
LMM	1 (1.0)
Desmoplastic	5 (5.1)
Unclassified melanoma	33 (33.7)
Ulceration of primary, n (%)	
Present	43 (39.0)
Absent	46 (36.4)
Unknown	29 (24.6)
Sentinel Lymph Node Biopsy, n (%)	

(Continued)

TABLE 1 Continued

Variable, n (%)	
Sentinel Lymph Node Biopsy, n (%)	
Positive	42 (35.6)
Negative	28 (23.7)
Not performed	48 (40.7)
PD-L1, n (%)	
Positive	40 (33.9)
Negative	54 (45.8)
Not performed	24 (20.3)
Tumor Thickness, n (%)	
Mean ± SD	3.38 ± 1.26
< 1 mm	9 (7.6)
1 - 2 mm	23 (19.5)
2 - 4 mm	29 (24.6)
> 4 mm	28 (23.7)
Unknown	29 (24.6)
First-line non-adjuvant systemic therapy, n (%)	
Anti-PD1 monotherapy	20 (35.7)
Anti-PD1 + anti-CTLA-4	13 (23.2)
unknown	8 (14.3)
other	15 (26.8)
Stage at therapy start, n (%)	
III	7 (12.5)
M1a	2 (3.6)
M1b	8 (14.3)
M1c	18 (32.1)
M1d	4 (7.1)
unknown	17 (30.4)
Tissue used for analysis, n (%)*	
Primary	65 (55.1)
Metastasis	33 (28.0)
Unknown	20 (17.0)

* Sums may not add to 100 because of rounding.

G13S). *NFI* mutations were present in 71 samples (60.2%) and activating *TERT*-promoter mutations in 54 samples (45.8%) (Supplementary Table 4). Other frequently mutated genes included *ARID1A* (65.3%), *ARID2* (59.3%), and *SMARCA4* (58.5%). Less frequent mutations were reported in various genes including *SF3B1*, *KIT*, *TERT*, *TP53*, *WT1*, *PIK3CA*, *FBXW7*, *GNA11*, *CTNNB1*, *PIK3R1*, *MAP2K1*, *MITF*, *IDH1*, *MAP2K2*, *GNAQ*, *PTEN*, *EZH1*, *RAC1* and *CDK4* (Figure 3).




FIGURE 1 Characteristics of *BAP1_{mut}* melanoma. Distribution of activating gene mutations in *BAP1_{mut}* non-veal (left) and uveal (right) melanoma tumor samples (A). Left: *BAP1_{mut}* melanoma harbored more mutations than *BAP1_{wt}* melanoma. Middle: Within the group of *BAP1_{mut}* melanoma, non-veal tumors exhibited higher mutation numbers than tumors of uveal origin. Right: Non-veal *BAP1_{mut}* tumors from cutaneous sites showed the highest number of mutations compared with tumors of uveal origin and mucosal, meningeal or occult origin (subsumed as “other”) (B). Uveal *BAP1_{mut}* tumor samples exhibited the lowest amount of C>T substitutions compared to both non-veal *BAP1_{mut}* and *BAP1_{wt}* melanomas (C). Statistical tests were performed using Welch’s t test and Dunnett’s test. Data is shown as mean ± SEM. *p < 0.05, ****p < 0.0001.

3.2.2 *BAP1_{mut}* uveal melanoma

No *BRAF*, *NRAS*, *NF1* or *TERT* promoter mutations were detected, though all tumor samples harbored additional mutations (Figure 4). *GNAQ* and *GNA11* mutations were frequent with 7 (63.6%) and 4 mutations (36.4%) and predominantly activating (100% of *GNAQ* mutations and 75% of *GNA11*). Mutations affecting codon 209 in *GNAQ* were Q209L (n=3), Q209P (n=2) and Q209R (n=1). One sample harbored an activating R183Q mutation. In *GNA11* all codon 209 mutations were Q209L (n=2). One sample harbored an activating R183C mutation in Exon 4. More than half of detected *BAP1* mutations were found to be inactivating. Rarer mutations identified were *SF3B1*, *ARID1A*, and *SMARCA4*.

3.3 Mutational load and ultraviolet signature mutations

BAP1_{mut} melanomas (n=129) exhibited a significantly higher number of mutations compared to *BAP1_{wt}* melanomas (n=1215) with 33.1 versus 4.1 mutations per sample. Within the group of *BAP1_{mut}* melanomas, uveal melanomas demonstrated lower mutation frequencies compared to *BAP1_{mut}* NUM (3.3 mutations versus 35.9 mutations per sample). Upon subgroup analysis of the non-veal *BAP1_{mut}* cohort, cutaneous melanomas exhibited a higher mutational load compared to those of mucosal, meningeal, or occult origin (mean 39.5 and 18.2 mutations per sample, respectively) (Figure 1B).

TABLE 2 Clinical characteristics of patients with *BAP1*_{mut} uveal melanoma (n=11).

Variable, n (%)	
Age at first diagnosis, n (%)	
Median (+/- SD)	65.3 (+/- 12.1)
Range	43–84
≤60 years	4 (36.4)
>60 years	7 (63.6)
Sex, n (%)	
Female	5 (45.5)
Male	6 (54.5)
Mutated oncogene, n (%)	
<i>GNAQ</i>	7 (63.6)
<i>GNA11</i>	4 (36.4)
<i>BAP1</i>	11 (100)
PD-L1, n (%)*	
Positive	2 (18.2)
Negative	5 (45.5)
Not performed	4 (36.4)
First-line non-adjuvant systemic therapy, n (%)	
Anti-PD1 monotherapy	1 (25.0)
Anti-PD1 + anti-CTLA-4	3 (75.0)
Other	0 (0)
Unknown	0 (0)
Stage at therapy start, n (%)	
III	1 (25.0)
M1a	0 (0)
M1b	1 (25.0)
M1c	2 (50.0)
M1d	0 (0)
Unknown	0 (0)
Tissue used for analysis, n (%)	
Primary	3 (27.3)
Metastasis	8 (72.7)

*Sums may not add to 100 because of rounding.

*BAP1*_{mut} NUM showed significantly more C>T alterations than *BAP1*_{wt} melanomas. Uveal *BAP1*_{mut} tumor samples were found to exhibit the lowest amount of C>T substitutions compared to both non-uveal *BAP1*_{mut} and *BAP1*_{wt} melanomas (Figure 1C).

3.4 Survival analysis and treatment response

Survival analysis showed a median overall survival time of 38.0 months for all included patients with stage IV *BAP1*_{mut} tumors with matching survival data (n=81). Comparison of OS between patients with *BAP1*_{mut} uveal and non-uveal melanoma revealed a longer survival for those with NUM, though nonsignificant (41.2 and 44.7, respectively, p=0.26) (Figure 5A).

3.4.1 *BAP1*_{mut} non-uveal melanoma

Survival rates of NUM patients receiving immunotherapy as first non-adjuvant therapy (n=29) were 7.4 (mPFS) and 28.1 months (mOS), respectively. Patients receiving targeted therapies (n=7) as first-line therapy had a mPFS of 11.3 and mOS of 37.0 months. Comparison of survival rates between ICI-cohort and TT-cohort did not show any significant difference in either PFS or OS: p=0.73 and p=0.76, respectively (Figures 5B, C).

Further analysis of OS in patients with stage IV *BAP1*_{mut} NUM depending on mutation-type showed a median OS of 57.0 months for patients with inactivating *BAP1* mutations (n=15) and 44.7 months for those with other mutation-types (n=50). The observed difference was not statistically significant (p=0.61) (Figure 5D).

3.4.2 *BAP1*_{mut} uveal melanoma

A case-by-case analysis for uveal melanoma patients was performed to evaluate treatment response (Supplementary Table 3). All patients with first-line non-adjuvant systemic therapy received ICI-based regimens (n=4). Treatment response to ICI was progressive disease in three patients (75%). One patient (25%) exhibited a partial response (this tumor harbored a *GNA11* R183C and a *BAP1* R385* mutation, Supplementary Table 3).

3.5 *BAP1*_{mut} non-uveal melanoma with a uveal mutation signature

In seven cases *BAP1*_{mut} non-uveal tumors were identified harboring activating *GNAQ* or *GNA11* mutations. Four tumors were of cutaneous origin, two occult and one melanocytoma of the central nervous system. Therapies were diverse and follow-up data incomplete (Supplementary Table 2).

4 Discussion

Our study aimed to investigate the genetic characteristics of *BAP1*_{mut} melanoma based on a cohort of 129 uveal and non-uveal melanoma patient cases, and to correlate these with clinicopathological data and outcomes.

To the best of our knowledge, this study is the largest to date investigating *BAP1*_{mut} non-uveal melanoma and contains the most detailed genetic analysis of this melanoma subtype.

TABLE 3 Comparison of clinical characteristics between *BAP1_{mut}* non-uveal and uveal melanoma patients.

Variable, n (%)	non-uveal (n=118)	uveal (n=11)	p-value
Age at first diagnosis, n (%)			.303
Mean +/- SD	60.4 (+/- 15.0)	65.3 (+/- 12.1)	
Range	22 - 82	43-84	
≤60 years	53 (44.9)	4 (36.4)	
>60 years	65 (55.1)	7 (63.6)	
Sex, n (%)			.482
Female	41 (34.7)	5 (45.5)	
Male	77 (65.3)	6 (54.5)	
Mutation distribution, n (%)			.003
Inactivating (INAC)	20 (16.9)	6 (54.5)	
other	98 (83.1)	5 (45.5)	
PD-L1, n (%)*			.221
Positive	42 (35.6)	2 (18.2)	
Negative	28 (23.7)	5 (45.5)	
Not performed	48 (40.7)	4 (36.4)	

* Sums may not add to 100 because of rounding.

Among *BAP1_{mut}* non-uveal melanoma cases, we observed a predominance of nodular melanoma as the most prevalent histopathological subtype, and a skewed distribution of tumor thickness towards thicker tumors. This finding is noteworthy as superficial spreading melanomas typically represent the prevailing subtype in Western countries (18). Mucosal and uveal melanomas were overrepresented compared to *BAP1_{wt}* cohorts, fitting existing data (9, 19–22).

Mutation patterns varied substantially between non-uveal and uveal samples. Uveal *BAP1_{mut}* melanomas exhibited significantly lower numbers of accompanying mutations and no evidence of UV-induced mutagenesis (23). In contrast, non-uveal *BAP1_{mut}* melanomas had a higher mutational burden and number of UV-signature mutations (C>T/CC>>TT transitions) than *BAP1_{wt}*

melanomas, indicating preferential tumor occurrence in sun-exposed skin (20).

Analysis of uveal *BAP1_{mut}* samples revealed a significantly lower total number of mutations, lacking common cutaneous driver mutations, while harboring known uveal melanoma driver mutations (10, 23). Genomic patterns of *BAP1_{mut}* non-uveal melanomas differed substantially from those of uveal origin in terms of mutational load and driver oncogenes: *NF1*, *BRAF*, and *NRAS* mutations were frequent, often with numerous co-mutations. *NF1* was the most common concomitant mutation. High mutation numbers and frequent *NF1* mutations may suggest that *BAP1_{mut}* non-uveal melanomas tend to be hypermutated tumors (24, 25). Previous reports on *BAP1_{mut}* cutaneous melanocytic tumors have indicated higher frequencies of concurrent *BRAF* V600E mutations compared to our cohort (26). It will be interesting to see if other, larger studies can validate this finding.

BAP1 mutations are associated with poor prognosis in uveal melanoma, but their prognostic value in non-uveal melanoma remains controversial (10, 27, 28). Recent studies have shown that *BAP1* mutations are associated with an inflammatory tumor microenvironment and increased immune cell infiltration, suggesting a potential role as a predictive biomarker for immunotherapy response (6, 29–32). Furthermore, it is well-documented that *BAP1* mutations in uveal melanoma strongly correlate with *BAP1* expression in immunohistochemical staining (33). However, we did not observe a significant difference in overall survival of stage IV non-uveal melanoma patients harboring *BAP1* mutations compared to published *BAP1* wildtype cohorts (16, 24). Within the cohort of uveal melanoma, a case-by-case analysis of four patients revealed a poor response to immunotherapy, consistent with previous studies, showing low efficacy of anti-PD-1 and anti-CTLA-4 therapies in uveal melanoma (9, 34). Overall survival independent of treatment in uveal melanoma patients, calculated from the initial diagnosis of stage IV, was relatively long compared to other cohorts of metastatic uveal melanoma patients reported previously (35). We believe this is partly due to selection bias, likely caused by the small number of patients with metastatic uveal melanoma treated in our department.

Although very rare, non-uveal melanoma with a uveal melanoma gene mutation signature can occur. These entities, termed “blue-nevus like melanoma” if cutaneous, or “primary




FIGURE 2 Lollipop mutation graph demonstrates the distribution of mutations throughout the *BAP1* gene with missense mutations shown in green, inactivating (Nonsense or frame-shift mutations) in black, and frameshift mutations in brown.




FIGURE 3 Mutation distribution in *BAP1_{mut}* non-veal melanoma. Green: mutations known or assumed to be activating. Red: loss of function mutations. Blue: known activating mutations in the TERT promoter region.

central nervous system melanoma” if derived from the central nervous system, behave similarly to uveal melanoma (36). Our cohort encompassed seven cases; however, limited case number and follow-up did not allow a representative comparison. In these tumors, *BAP1* mutations should not be seen as passenger mutations but relevant markers of metastasis and prognosis (37, 38).

Our study has certain limitations. We conducted sequencing on both primary tumors and metastases, and our assay may not have detected deletions involving entire exons, potentially resulting in missed identification of *BAP1* alterations in some patients. Due to the retrospective study design and long data collection period as well as advances in sequencing technology over the years, there might be variations in the mutation detection rate or

characterization accuracy over time. In addition, changes in treatment standards have occurred, making the interpretation of survival analysis challenging for this study. The cohort we analyzed was heterogeneous and consisted of various types of melanoma, including cutaneous, mucosal, occult, and meningeal melanoma. Furthermore, due to the retrospective nature of this study, we did not have access to comprehensive immunohistochemical staining for this cohort, which could have provided additional information, such as whether loss of protein expression is a good surrogate for *BAP1* mutation status in non-veal melanoma, as has been well demonstrated for uveal melanoma.

Although our findings are based on the largest cohort of *BAP1_{mut}* non-veal melanomas to date, larger, preferably prospective studies are needed to validate our results.




FIGURE 4 Mutation distribution in *BAP1_{mut}* uveal melanoma. Green: mutations known or assumed to be activating. Red: loss of function mutations.




FIGURE 5

Patients with stage IV $BAP1_{mut}$ non-uveal melanoma ($n=73$) did not show a difference in overall survival compared to patients with stage IV $BAP1_{mut}$ uveal melanoma ($n=8$) (A). Survival rates of $BAP1_{mut}$ non-uveal melanoma patients receiving immunotherapy as first non-adjuvant therapy compared to those receiving targeted therapies showed no significant difference in either progression-free or overall survival (B, C). Patients with inactivating $BAP1$ mutations did not differ in overall survival compared to those with other mutation types (D).

Our analysis demonstrates that, except for rare cases such as non-uveal melanomas exhibiting a uveal melanoma mutation signature and cases involving germline mutations, where $BAP1$ mutations are associated with poor prognosis or familial predisposition syndromes, respectively, $BAP1$ mutations in non-uveal melanomas are typically passenger mutations. These mutations are predominantly found in heavily mutated tumors and do not appear to have any significant prognostic or therapeutic value.

Data availability statement

The data analyzed in this study is subject to the following licenses/restrictions: The data underlying this article will be shared on reasonable request to the corresponding author. Requests to access these datasets should be directed to klaus.griewank@uk-essen.de.

Ethics statement

The studies involving humans were approved by ethics committee of the University of Duisburg-Essen (ethics approval no. 21-9873-BO). The studies were conducted in accordance with

the local legislation and institutional requirements. The human samples used in this study were acquired from a by-product of routine care or industry. Written informed consent for participation was not required from the participants or the participants' legal guardians/next of kin in accordance with the national legislation and institutional requirements.

Author contributions

JM: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing. J-MP: Data curation, Writing – review & editing. GL: Writing – review & editing. AZ: Writing – review & editing. JU: Data curation, Writing – review & editing. PT: Data curation, Writing – review & editing. CP: Data curation, Writing – review & editing. RH: Data curation, Writing – review & editing. AK: Data curation, Writing – review & editing. JW: Data curation, Writing – review & editing. JK: Data curation, Writing – review & editing. IM: Data curation, Writing – review & editing. AS: Data curation, Writing – review & editing. AP: Data curation, Writing – review & editing. EL: Data curation, Writing – review & editing. LZ: Data curation, Writing – review & editing. EH: Data curation, Writing – review & editing. SU: Data curation, Writing – review & editing. DS: Data curation, Writing – review & editing.

CT: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Supervision, Writing – review & editing. KG: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Supervision, Writing – review & editing.

Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article. This work was in part supported by Bristol Myers Squibb for the multicenter translational study “Tissue Registry in Melanoma” (TRIM) within the framework of the skin cancer registry ADOREG of the German Dermatologic Cooperative Oncology Group (DeCOG). J-MP was supported by the DFG (German Research Foundation)funded Clinician Scientist Program of the University Medicine Essen Clinician Scientist Academy (UMEA) (FU 356/12-1).

Acknowledgments

The authors are indebted to all patients and their relatives. Human biological samples and related data were obtained from the Westdeutsche Biobank Essen (11–4715-BO, n=60), and from the prospective multicenter translational study Tissue Registry in Melanoma (ADOREG/TRIM; NCT05750511; CA209–578; 15–6566-BO, n=69) conducted by the German Dermatological Cooperative Oncology Group (DeCOG).

Conflict of interest

JM: Declares travel support from Bristol Myers Squibb, Novartis and Sun Pharmaceutical Industries, outside the submitted work. J-MP: served as consultant and/or has received honoraria from Bristol-Myers Squibb, Novartis, Sanofi and received travel support from Bristol-Myers Squibb, Novartis, Pierre Fabre and Therakos, outside the submitted work. GL: Declares travel support from Sun Pharma, outside the submitted work. AZ: Declares travel support from Novartis, Sanofi Genzyme, and Bristol-Myers Squibb, outside the submitted work. JU: Is on the advisory board or has received honoraria and travel support from Amgen, Bristol Myers Squibb, GSK, Immunocore, LeoPharma, Merck Sharp and Dohme, Novartis, Pierre Fabre, Roche, Sanofi outside the submitted work. PT: served as consultant and/or received honoraria form Almirall, Bristol Myers Squibb, Biofrontera, Curevac, Kyowa Kirin, Merck, Merck Sharp & Dohme, Novartis, Pierre-Fabre, Roche, Sanofi, 4SC, and travel support from Bristol Myers Squibb outside the submitted work. CP: Received honoraria speaker honoraria and advisory-board honoraria and travel support from BMS, MSD, Novartis, Merck Serono, Pierre Fabre, Sunpharma, AbbVie, LEO, and

Kyona Kirin, outside the submitted work. RH: Is an employee of Helios Kliniken Erfurt GmbH. JW: Received honoraria and travel support from Almirall, Bristol Myers Squibb, Novartis, Pierre Fabre and Merck Sharp & Dohme, outside the submitted work. EL: Served as consultant and/or has received honoraria from Bristol-Myers Squibb, Merck Sharp & Dohme, Novartis, Pierre-Fabre, Sanofi, Sunpharma, Takeda and travel support from Bristol-Myers Squibb, Pierre Fabre, Sunpharma and Novartis, outside the submitted work. LZ: Served as consultant and/or has received honoraria from Bristol-Myers Squibb, Merck Sharp & Dohme, Novartis, Pierre-Fabre, Sunpharma and Sanofi; Research funding to institution: Novartis; travel support from Merck Sharp & Dohme, Bristol- Myers Squibb, Amgen, Pierre-Fabre, Sunpharma and Novartis, outside the submitted work. SU: Research support from Bristol Myers Squibb and Merck Serono; speakers and advisory board honoraria from Bristol Myers Squibb, Merck Sharp & Dohme, Merck Serono, and Novartis; meeting and travel support from Almirall, Bristol-Myers Squibb, IGEA Clinical Biophysics, Merck Sharp & Dohme, Novartis, Pierre Fabre, and Sun Pharma, outside the submitted work. DS: Reports personal fees and non-financial support from Roche/Genentech, grants, personal fees, non-financial support and other from BMS, personal fees from Merck Sharp & Dohme, personal fees and non-financial support from Merck Serono, grant, personal fees and non-financial support from Amgen, personal fees from Immunocore, personal fees from Incyte, personal fees from 4SC, personal fees from Pierre Fabre, personal fees and non-financial support from Sanofi/Regeneron, personal fees from Array BioPharma, personal fees from Pfizer, personal fees from Philogen, personal fees from Regeneron, personal fees from Nektar, personal fees from Sandoz, grants, personal fees and non-financial support from Novartis, personal fees and non-financial support from SunPharma, Replimune, Helsinn, OncoSec and InFlaRx outside the submitted work.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2024.1383125/full#supplementary-material>

References

- Schadendorf D, van Akkooi ACJ, Berking C, Griewank KG, Gutzmer R, Hauschild A, et al. Melanoma. *Lancet*. (2018) 392:971–84. doi: 10.1016/S0140-6736(18)31559-9
- Davis EJ, Johnson DB, Sosman JA, Chandra S. Melanoma: What do all the mutations mean? *Cancer*. (2018) 124:3490–9. doi: 10.1002/cncr.31345
- Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, et al. Signatures of mutational processes in human cancer. *Nature*. (2013) 500:415–21. doi: 10.1038/nature12477
- Cancer Genome Atlas N. Genomic classification of cutaneous melanoma. *Cell*. (2015) 161:1681–96. doi: 10.1016/j.cell.2015.05.044
- Jensen DE, Proctor M, Marquis ST, Gardner HP, Ha SI, Chodosh LA, et al. BAP1: a novel ubiquitin hydrolase which binds to the BRCA1 RING finger and enhances BRCA1-mediated cell growth suppression. *Oncogene*. (1998) 16:1097–112. doi: 10.1038/sj.onc.1201861
- Louie BH, Kurzrock R. BAP1: Not just a BRCA1-associated protein. *Cancer Treat Rev*. (2020) 90:102091. doi: 10.1016/j.ctrv.2020.102091
- Wang A, Papneja A, Hyrcza M, Al-Habeeb A, Ghazarian D. Gene of the month: BAP1. *J Clin Pathol*. (2016) 69:750–3. doi: 10.1136/jclinpath-2016-203866
- Rai K, Pilarski R, Cebulla CM, Abdel-Rahman MH. Comprehensive review of BAP1 tumor predisposition syndrome with report of two new cases. *Clin Genet*. (2016) 89:285–94. doi: 10.1111/cge.12630
- Yang J, Manson DK, Marr BP, Carvajal RD. Treatment of uveal melanoma: where are we now? *Ther Adv Med Oncol*. (2018) 10:1758834018757175. doi: 10.1177/1758834018757175
- Decatur CL, Ong E, Garg N, Anbunathan H, Bowcock AM, Field MG, et al. Driver mutations in uveal melanoma: Associations with gene expression profile and patient outcomes. *JAMA Ophthalmol*. (2016) 134:728–33. doi: 10.1001/jamaophthalmol.2016.0903
- Kumar R, Taylor M, Miao B, Ji Z, Njauw JC, Jonsson G, et al. BAP1 has a survival role in cutaneous melanoma. *J Invest Dermatol*. (2015) 135:1089–97. doi: 10.1038/jid.2014.528
- Liu-Smith F, Lu Y. Opposite roles of BAP1 in overall survival of uveal melanoma and cutaneous melanoma. *J Clin Med*. (2020) 9:411. doi: 10.3390/jcm9020411
- Luo X, Xu Y, Li Y, Zhang G, Huang S, Liu X, et al. BAP1 deletion abrogates growth and metastasis of murine cutaneous melanoma. *Melanoma Res*. (2021) 31:119–29. doi: 10.1097/CMR.0000000000000714
- Keung EZ, Gershenwald JE. The eighth edition American Joint Committee on Cancer (AJCC) melanoma staging system: implications for melanoma treatment and care. *Expert Rev Anticancer Ther*. (2018) 18:775–84. doi: 10.1080/14737140.2018.1489246
- Griewank KG, Westekemper H, Murali R, Mach M, Schilling B, Wiesner T, et al. Conjunctival melanomas harbor BRAF and NRAS mutations and copy number changes similar to cutaneous and mucosal melanomas. *Clin Cancer Res*. (2013) 19:3143–52. doi: 10.1158/1078-0432.CCR-13-0163
- Thielmann CM, Matull J, Roth S, Placke JM, Chorti E, Zaremba A, et al. Genetic and clinical characteristics of ARID1A mutated melanoma reveal high tumor mutational load without implications on patient survival. *Cancers (Basel)*. (2022) 14:2090. doi: 10.3390/cancers14092090
- Team RC. *R: A language and environment for statistical computing*. R foundation for statistical computing. (Vienna). (2010).
- Elder DE, Bastian BC, Cree IA, Massi D, Scolyer RA. The 2018 world health organization classification of cutaneous, mucosal, and uveal melanoma: Detailed analysis of 9 distinct subtypes defined by their evolutionary pathway. *Arch Pathol Lab Med*. (2020) 144:500–22. doi: 10.5858/arpa.2019-0561-RA
- Murali R, Wilmott JS, Jakrot V, Al-Ahmadie HA, Wiesner T, McCarthy SW, et al. BAP1 expression in cutaneous melanoma: a pilot study. *Pathology*. (2013) 45:606–9. doi: 10.1097/PAT.0b013e3283653818
- Anderson WF, Pfeiffer RM, Tucker MA, Rosenberg PS. Divergent cancer pathways for early-onset and late-onset cutaneous Malignant melanoma. *Cancer*. (2009) 115:4176–85. doi: 10.1002/cncr.24481
- McLaughlin CC, Wu XC, Jemal A, Martin HJ, Roche LM, Chen VW. Incidence of noncutaneous melanomas in the U.S. *Cancer*. (2005) 103:1000–7. doi: 10.1002/cncr.20866
- Song H, Wang L, Lyu J, Wu Y, Guo W, Ren G. Loss of nuclear BAP1 expression is associated with poor prognosis in oral mucosal melanoma. *Oncotarget*. (2017) 8:29080–90. doi: 10.18632/oncotarget.16175
- Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, et al. Mutations of the BRAF gene in human cancer. *Nature*. (2002) 417:949–54. doi: 10.1038/nature00766
- Thielmann CM, Chorti E, Matull J, Murali R, Zaremba A, Lodde G, et al. NF1-mutated melanomas reveal distinct clinical characteristics depending on tumour origin and respond favourably to immune checkpoint inhibitors. *Eur J Cancer*. (2021) 159:113–24. doi: 10.1016/j.ejca.2021.09.035
- Cirenajwis H, Lauss M, Ekedahl H, Torngren T, Kvist A, Saal LH, et al. NF1-mutated melanomas harbor distinct clinical and biological characteristics. *Mol Oncol*. (2017) 11:438–51. doi: 10.1002/1878-0261.12050
- Piris A, Mihm MC Jr, Hoang MP. BAP1 and BRAFV600E expression in benign and Malignant melanocytic proliferations. *Hum Pathol*. (2015) 46:239–45. doi: 10.1016/j.humpath.2014.10.015
- Harbour JW, Onken MD, Roberson ED, Duan S, Cao L, Worley LA, et al. Frequent mutation of BAP1 in metastasizing uveal melanomas. *Science*. (2010) 330:1410–3. doi: 10.1126/science.1194472
- Ewens KG, Kanetsky PA, Richards-Yutz J, Purrzellera J, Shields CL, Ganguly T, et al. Chromosome 3 status combined with BAP1 and EIF1AX mutation profiles are associated with metastasis in uveal melanoma. *Invest Ophthalmol Vis Sci*. (2014) 55:1600–7. doi: 10.1167/iovs.14-14550
- Hirsch TZ, Negulescu A, Gupta B, Caruso S, Noblet B, Couchy G, et al. BAP1 mutations define a homogeneous subgroup of hepatocellular carcinoma with fibrolamellar-like features and activated PKA. *J Hepatol*. (2020) 72:924–36. doi: 10.1016/j.jhep.2019.12.006
- Shrestha R, Nabavi N, Lin YY, Mo F, Anderson S, Volik S, et al. BAP1 haploinsufficiency predicts a distinct immunogenic class of Malignant peritoneal mesothelioma. *Genome Med*. (2019) 11:8. doi: 10.1186/s13073-019-0620-3
- Ladanyi M, Sanchez Vega F, Zauderer M. Loss of BAP1 as a candidate predictive biomarker for immunotherapy of mesothelioma. *Genome Med*. (2019) 11:18. doi: 10.1186/s13073-019-0631-0
- Spencer KR, Wang J, Silk AW, Ganesan S, Kaufman HL, Mehnert JM. Biomarkers for immunotherapy: current developments and challenges. *Am Soc Clin Oncol Educ Book*. (2016) 35:e493–503. doi: 10.1200/EDBK_160766
- Koopmans AE, Verdijk RM, Brouwer RW, van den Bosch TP, van den Berg MM, Vaarwater J, et al. Clinical significance of immunohistochemistry for detection of BAP1 mutations in uveal melanoma. *Mod Pathol*. (2014) 27:1321–30. doi: 10.1038/modpathol.2014.43
- Heppt MV, Amaral T, Kahler KC, Heinzerling L, Hassel JC, Meissner M, et al. Combined immune checkpoint blockade for metastatic uveal melanoma: a retrospective, multi-center study. *J Immunother Cancer*. (2019) 7:299. doi: 10.1186/s40425-019-0800-0
- Rodriguez-Vidal C, Fernandez-Diaz D, Fernandez-Marta B, Lago-Baameiro N, Pardo M, Silva P, et al. Treatment of metastatic uveal melanoma: Systematic review. *Cancers (Basel)*. (2020) 12:2557. doi: 10.3390/cancers12092557
- Griewank KG, Muller H, Jackett LA, Emberger M, Moller I, van de Nes JA, et al. SF3B1 and BAP1 mutations in blue nevus-like melanoma. *Modern pathology: an Off J United States Can Acad Pathology Inc*. (2017) 30:928–939. doi: 10.1038/modpathol.2017.23
- Smit KN, Jager MJ, de Klein A, Kiliç E. Uveal melanoma: Towards a molecular understanding. *Prog Retin Eye Res*. (2020) 75:100800. doi: 10.1016/j.preteyeres.2019.100800
- Griewank KG, Koelsche C, van de Nes JAP, Schrimpf D, Gessi M, Moller I, et al. Integrated genomic classification of melanocytic tumors of the central nervous system using mutation analysis, copy number alterations, and DNA methylation profiling. *Clin Cancer Res*. (2018) 24:4494–504. doi: 10.1158/1078-0432.CCR-18-0763