



OPEN ACCESS

EDITED AND REVIEWED BY

Luis Pedro Sousa Inês,
Coimbra Hospital and University Center,
Portugal

*CORRESPONDENCE

Nasrin Asgari
✉ nasgari@health.sdu.dk

†PRESENT ADDRESS

Mads Nikolaj Olesen,
Lundbeck A/S, Copenhagen, Denmark

RECEIVED 14 May 2024

ACCEPTED 27 May 2024

PUBLISHED 04 June 2024

CITATION

Wegener MAB, Möller S, Olesen MN,
Madsen JS, Sorensen GL, Voss A and Asgari N
(2024) Corrigendum: Microfibrillar-associated
protein 4 interaction with inflammation and
clinical characteristics in neuropsychiatric
systemic lupus erythematosus.
Front. Lupus 2:1432884.
doi: 10.3389/flupu.2024.1432884

COPYRIGHT

© 2024 Wegener, Möller, Olesen, Madsen,
Sorensen, Voss and Asgari. This is an open-
access article distributed under the terms of
the [Creative Commons Attribution License
\(CC BY\)](#). The use, distribution or reproduction
in other forums is permitted, provided the
original author(s) and the copyright owner(s)
are credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Corrigendum: Microfibrillar-associated protein 4 interaction with inflammation and clinical characteristics in neuropsychiatric systemic lupus erythematosus

Maria Alexandra Bandehkhoda Wegener^{1,2}, Sören Möller^{3,4},
Mads Nikolaj Olesen^{1,2,5,6†}, Jonna Skov Madsen^{2,7},
Grith Lykke Sorensen^{5,6,8}, Anne Voss^{3,9,10} and Nasrin Asgari^{1,2,3,5*}

¹Department of Neurology, Slagelse Hospital, Slagelse, Denmark, ²Institute of Regional Health Research, University of Southern Denmark, Odense, Denmark, ³Open Patient Data Explorative Network, Odense University Hospital & Research Unit OPEN, Odense, Denmark, ⁴The Department of Clinical Research, University of Southern Denmark, Odense, Denmark, ⁵Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark, ⁶The Department of Clinical Immunology, Odense University Hospital, Odense, Denmark, ⁷Department of Clinical Biochemistry and Immunology, Lillebaelt Hospital, Vejle, Denmark, ⁸Cancer and Inflammation Research, Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark, ⁹Department of Rheumatology, Odense University Hospital, Odense, Denmark, ¹⁰Research Unit of Rheumatology, University of Southern Denmark, Odense, Denmark

KEYWORDS

systemic lupus erythematosus, neuropsychiatric SLE, inflammation, microfibrillar-associated protein 4, accumulated disease damage

A Corrigendum on

Microfibrillar-associated protein 4 interaction with inflammation and clinical characteristics in neuropsychiatric systemic lupus erythematosus

Wegener MAB, Möller Sören, Olesen MN, Madsen JS, Sorensen GL, Voss A and Asgari N (2024). Front. Lupus 2:1386256. doi: 10.3389/flupu.2024.1386256

In the published article, there was an error. “Data not shown” should have been deleted. A correction has been made to **3 Results**, *3.5 Markers of BBB disruption*. This sentence previously stated:

“Additionally, MMP-2 was significantly elevated in NPSLE patients with CNS manifestations compared to PNS manifestations ($p = 0.010$, data not shown).”

The corrected sentence appears below:

“Additionally, MMP-2 was significantly elevated in NPSLE patients with CNS manifestations compared to PNS manifestations ($p = 0.010$).”

In the published article, there was an error. The below sentences in the Discussion should have been deleted. A correction has been made to the **4 Discussion**. The below sentences have been removed:

“Interestingly, MFAP4 was associated with SLICC (accumulated disease damage) in the total SLE cohort suggesting that MFAP4 may reflect inflammation. Interestingly, MFAP4 was associated with SLICC in the total SLE cohort suggesting that MFAP4 may reflect inflammation.”

In the published article, there was an error in the punctuation of the of caption Figure 4. The corrected legend appears below.

Figure 4. Correlation of MFAP4 with MMP-7 and VEGF in NPSLE subgroup and non-NPSLE subgroup. A significant association was found between MMP-7 and MFAP4 within NPSLE patients (A). Likewise, a significant association was found between VEGF and MFAP4 within NPSLE patients (C). No association was found between MMP-7 and MFAP4 (B) nor between VEGF and MFAP4 (D) in the non-NPSLE subgroup. Grey zone, confidence band. MFAP4, microfibrillar-associated protein 4; MMP-7, matrix metalloproteinase 7; VEGF, vascular endothelial growth factor; SLE, systemic lupus erythematosus; NPSLE, neuropsychiatric systemic lupus erythematosus.

The authors apologize for these errors and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.