



Benthic Cyanobacterial Diversity and Antagonistic Interactions in Abrolhos Bank: Allelopathy, Susceptibility to Herbivory, and Toxicity

OPEN ACCESS

Edited by:

Mirko Mutalipassi,
Anton Dohrn Zoological Station, Italy

Reviewed by:

Ethan C. Cissell,
Florida State University, United States
Olivier P. Thomas,
National University of Ireland Galway,
Ireland
Lucia Barra,
Zoological Station Anton Dohrn, Italy

*Correspondence:

Paulo Sergio Salomon
pssalomon@gmail.com

† Present address:

Renato Crespo Pereira,
Instituto de Pesquisas Jardim
Botânico do Rio de Janeiro, Rio de
Janeiro, Brazil

Specialty section:

This article was submitted to
Marine Biotechnology,
a section of the journal
Frontiers in Marine Science

Received: 06 October 2021

Accepted: 23 November 2021

Published: 06 January 2022

Citation:

Ribeiro FV, Caires TA,
Simões MAA, Hargreaves PI,
Vilela LB, Fistarol GO,
Cazelgrandi Junior AB,
Pereira-Filho GH, Moura RL,
Pereira RC and Salomon PS (2022)
Benthic Cyanobacterial Diversity
and Antagonistic Interactions
in Abrolhos Bank: Allelopathy,
Susceptibility to Herbivory,
and Toxicity.
Front. Mar. Sci. 8:790277.
doi: 10.3389/fmars.2021.790277

Felipe de Vargas Ribeiro^{1,2}, Taiara Aguiar Caires³,
Marcela Alvarenga de Almeida Simões⁴, Paulo liboshi Hargreaves⁴,
Livia Bonetti Villela^{4,5}, Giovana de Oliveira Fistarol⁴,
Alexandre Brauns Cazelgrandi Junior⁶, Guilherme Henrique Pereira-Filho⁷,
Rodrigo Leão de Moura^{4,8}, Renato Crespo Pereira^{2†} and Paulo Sergio Salomon^{4,8*}

¹ Programa de Pós-Graduação em Dinâmica dos Oceanos e da Terra, Universidade Federal Fluminense, Niterói, Brazil, ² Departamento de Biologia Marinha, Universidade Federal Fluminense, Niterói, Brazil, ³ Departamento de Ciências Humanas e Tecnologias, Universidade do Estado da Bahia, Salvador, Brazil, ⁴ Departamento de Biologia Marinha, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil, ⁵ Programa de Pós Graduação em Genética, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil, ⁶ Programa de Pós-Graduação em Ecologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil, ⁷ Departamento de Ciências do Mar, Universidade Federal de São Paulo, São Paulo, Brazil, ⁸ Laboratório de Avaliação e Monitoramento da Biodiversidade, Núcleo Rogério Valle de Produção Sustentável (SAGE), Instituto Alberto Luiz Coimbra de Pós-Graduação e Pesquisa de Engenharia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

Benthic cyanobacterial mats (BCMs) are conspicuous components of coral reef communities, where they play key ecological roles as primary producers among others. BCMs often bloom and might outcompete neighboring benthic organisms, including reef-building corals. We investigated the cyanobacterial species composition of three BCMs morphotypes from the marginal reef complex of Abrolhos Bank (Southeastern Brazil). Also, we assessed their allelopathic effects on coral zooxanthellae, their susceptibility to herbivory by fish, and their toxicity to brine shrimp nauplii. Morphology and 16S rDNA sequencing unveiled the cyanobacteria *Moorena bouillonii*, *Okeania erythroflocculosa*, *Adonisia turfae*, *Leptolyngbya* sp., and *Halomicronema* sp. as components of BCMs from Abrolhos. BCMs cell-free filtrates and extracts exerted an allelopathic effect by reducing the growth of the *ex hospite* *Symbiodinium* sp. in culture. BCMs-only treatments remained untouched in field susceptibility assays in contrast to macroalgae only and mixed BCMs-macroalgae treatments that had the macroalgae fully removed by reef fish. Crude aqueous extracts from BCMs were toxic to brine shrimps in acute assays. Besides unveiling the diversity of BCMs consortia in Abrolhos, our results cast some light on their allelopathy, antiherbivory, and toxicity properties. These antagonistic interactions might promote adverse cascading effects during benthic cyanobacteria blooms and in gradual shifts to BCMs-dominated states.

Keywords: BCMs, Oscillatoriales, Synechococcales, zooxanthellae, interspecific interactions, Southwestern Atlantic, turbid reefs, chemical mediation

INTRODUCTION

Benthic cyanobacterial mats (BCMs) are ubiquitous in coral reef ecosystems, where they contribute to critical functions such as nitrogen fixation, sulfate reduction, and primary production (Charpy et al., 2007). The term BCMs is used to define a range of cyanobacteria-rich mats that might display diverse morphologies, from periphytic biofilms and horizontally-spreading mats over sediment and rubble, to tufts and fronds of various heights, up to tens of centimeters (Graham and Wilcox, 2000; Charpy et al., 2007; Cissell et al., 2019; Cissell and McCoy, 2021). Filamentous cyanobacteria can also be found entangled in other structurally more complex benthic assemblages, namely turf (Connell et al., 2014). These mats can be formed by a single genus (Paul et al., 2005) or by a consortium of multiple taxa (Echenique-Subiabre et al., 2015), including protists and non-cyanobacterial, heterotrophic prokaryotes, and viruses (Cissell and McCoy, 2021). Despite its relevance, the identity of BCMs components is often overlooked by morphology-based taxonomy (Engene et al., 2013a). The globally distributed genus *Lyngbya* C. Agardh ex Gomont is a foremost example of the cryptic diversity within benthic marine cyanobacteria. The recent combination of molecular, chemical, and morphological analyses revealed three additional genera [*Moorena* Engene and Tronholm, *Okeania* N. Engene et al., and *Dapis* Engene et al. (2018)] and several new species within the *Lyngbya* complex (Engene et al., 2012, 2013b, 2018; Tronholm and Engene, 2019).

Increasing seawater warming and nutrient uploads stimulate BCMs growth which might lead to blooms in coastal areas (Sneed et al., 2017) and might be associated with the coral cover decline (Ford et al., 2018). Reef degradation generates newly available substrates easily occupied by fast-growing BCMs (e.g., Albert et al., 2005). BCMs can accentuate their presence even under limited space by growing directly over living organisms, such as scleractinian corals, fleshy algae, and seagrass (Yamamuro, 1999; Ritson-Williams et al., 2005; Puyana and Prato, 2013).

Cyanobacteria produce various cyclic and linear peptides, alkaloids, macro lactones, and heterocyclic compounds (Carroll et al., 2021). These chemicals can act as grazing deterrents against fishes (e.g., Thacker et al., 1997; Capper et al., 2016), sea urchins, mollusks, and crustaceans (see Leão et al., 2012), although some specialist sea hare (Capper and Paul, 2008; Ribeiro et al., 2017) and reef fish (Cissell et al., 2019) can actively graze on BCMs. Although cyanobacteria are an important source of nitrogen for herbivorous fish in marine (Yamamuro, 1999; Nicholson and Clements, 2020), and freshwater systems (Yang and Dudgeon, 2010), their toxicity and nutritional value is varied among taxa and within subpopulations (Nagarkar et al., 2004; Groendahl and Fink, 2017). Such variability also constrains the feeding behavior of consumers to meet their nutritional requirements (Mendes et al., 2018).

Cyanobacterial compounds also have allelopathic properties against competitors and may significantly impact the physiology of reef organisms (Thacker et al., 1998; Chadwick and Morrow, 2011), through microbiome disruption (Morrow et al., 2011). Such antagonistic interactions can generate associated cascading effects since competition strongly determines the structure and

composition of coral reef communities (Birkeland, 1997). Benthic cyanobacterial mat allelopathic effects in coral reef environments are still poorly investigated (Brumley et al., 2018). Few studies have demonstrated that BCMs and the allelochemicals they produce can damage the coral's tissue and microbiome (e.g., Titlyanov and Titlyanova, 2009; Morrow et al., 2011). In the nutrified and turbid reef systems of Abrolhos (Leão et al., 2016), BCMs have been associated with long-term impacts on coral growth (Ribeiro et al., 2018) and altered states of benthic dynamics (Teixeira et al., 2021).


Blooms of BCMs covering several benthic organisms have been observed in Abrolhos reefs during the summer (Ribeiro et al., 2017). Whereas the concentration of allelopathic chemicals depends on the BCMs assemblage structure (Brumley et al., 2018), the diffused compounds in benthic boundary layers could harm corals and alter trophic interactions (Brocke et al., 2015). Considering the coral tissue suppression by cyanobacteria previously reported (Ribeiro et al., 2018), we aimed to investigate the interaction between cyanobacteria and coral photosymbionts. Although symbiotic dinoflagellates are embedded in the gastrodermis, the constant mass transfer through boundary layers and tissues could expose them to toxic metabolites from the environment (Patterson, 1992). This pathway would be enhanced in the current global warming scenario (Teixeira et al., 2019) since tissue thickness is reduced in thermally stressed corals (Ainsworth et al., 2008), which increases mass transfer rates (Patterson, 1992).

We hypothesized that potentially toxic cyanobacteria within BCMs undermine coral physiology through the impairment of endosymbionts and disturb consumer–prey interactions. Here, we characterize the BCMs taxonomic composition of Abrolhos and evaluate BCM's allelopathy, susceptibility to herbivory, and toxicity.

MATERIALS AND METHODS

Sampling and Establishment of Cyanobacterial Cultures

Benthic cyanobacterial mats of three distinct morphotypes were manually collected by SCUBA divers between 3 and 8 m depth in the Abrolhos Bank (16°40', 19°40'S—39°10', 37°20'W) (Figure 1). Sampling was performed at two different sites on pinnacle reef tops: the nearshore Pedra de Leste (PLES) of Parcel das Paredes reef and the offshore site of Parcel dos Abrolhos reef (PAB). Throughout the text samples, PLES and PAB will be referred to as PLES_{BCM} and PAB_{BCM}, respectively. The BCMs cover and long-term survey data are described in Teixeira et al. (2021). An aliquot of each sample was rinsed with filtered (GF/F) seawater and snap-frozen or fixed in 1% glutaraldehyde. Additional aliquots were transferred to vials containing f/2 medium (Guillard, 1975), prepared with 0.22 μm filtered Abrolhos seawater or filtered seawater, and transported to the laboratory. Single cyanobacterial filaments were isolated in a flow cytometer (MoFlo, Beckman Coulter, United States) (Fistartol et al., 2018) or by manual micromanipulation under an inverted microscope in case of wider filaments. Collected



filaments were chosen to encompass distinct diameter diversities within conspicuous BCMs morphotypes in the Abrolhos Bank. Isolated filaments were transferred to *f/2* medium (Guillard, 1975), prepared with 0.22 μm filtered Abrolhos seawater, and kept at 26°C with a 14:10-h light/photoperiod and ca. 20 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$. Established cultures were deposited in the Culture Collection of Microorganisms at UFRJ (CCMR).

Taxonomic Characterization of the Benthic Cyanobacterial Mats

For each BCMs sample, single wide and narrow filaments were selected from the tufts under an inverted microscope (Leica DMi8[®]) and thoroughly rinsed with a sterile *f/2* medium. The selected filaments were photographed with a digital camera for further morphometric analysis using Leica Application Suite X 3.4.1.17822 (Leica[®], Mannheim, Germany) software. We measured a minimum of 20 filaments of each morphotype from the same sample, whenever possible, using the AxioVs40 v4.8.2.0 imaging system (Carl Zeiss, Jena, Germany). Single filaments were transferred to ca. 1 μL of medium to 0.5 mL in PCR tubes and digested with 0.5 μL of proteinase K (23 mg ml^{-1}) at 37°C for 1 h, followed by 80°C for 5 min to denature the enzyme. These digested mixtures were further used directly in PCR to amplify the 16S rDNA gene. For

cyanobacteria maintained in cultures, DNA was extracted from a cell pellet with 2% CTAB buffer (hexadecyl-trimethylammonium bromide, pH 8.0), followed by chloroform/isoamyl alcohol (24:1, v/v) to take off organic compounds. DNA was precipitated with 95% ethanol and 0.3 M NaOAc, rinsed two times with cold 70% ethanol, and resuspended in 25 μL of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0). After DNA extraction, the integrity of the genetic material was checked on a 1% agarose gel. DNA quantification was done in a Nanodrop 1000 spectrophotometer (Thermo Scientific, United States).

Amplification of the 16S rDNA was performed according to Lane (1991) by nested PCR, using external universal 16S rDNA primers (forward 27F 5'-AGAGTTTGTATCMTGGCTCAG-3'; reverse 1492R 5'-GGTTACCTTGTTACGACTT-3') and subsequently internal cyanobacteria-specific primers (forward CYA359F 5'-GGGGAATYTTCCGCAATGGG-3' and CYA106F 5'-CGGACGGGTGAGTAACGCGTGA-3'; reverse CYA781R 5'-GACTACTGGGTATCTAATCCCWTT-3'), following previously described procedures (Nübel et al., 1997). Fragments using CYA106F were chosen to be amplified and confirmed by electrophoresis in agarose gel. The PCR products were purified with ExoSAP-IT (Thermo Scientific, United States). Sample sequencing was obtained by the SANGER method (Applied Biosystem 3500). Contigs were assembled with SeqMan

Pro v.11.1, and electropherograms were visually checked for peak quality or reading errors and compared among them after alignments, and also to published sequences using the BLASTn tool in GenBank. In this study, 15 cyanobacterial sequences from Abrolhos were produced. A set of sequences from cultures with high similarity to these, sequences from type specimens, and sequences from closely related groups were retrieved to be used in phylogenetic reconstruction (49 sequences in total). Also, an external sequence from *E. coli* (NC000913) was chosen to root the topology. Sequences were aligned in ClustalW implemented in MEGA X (Kumar et al., 2018) with further manual refinement. Bayesian information criterion (BIC) determined the evolutionary substitution model with the model test, also implemented in MEGA X. Phylogenetic reconstructions were done by Bayesian and maximum likelihood (ML) methods. Mrbayes 3.2 (Ronquist et al., 2012) was used with 1 M of MCMC searching, discarding 10% as burn-in, and web-IQTREE (Trifinopoulos et al., 2016) with the UFBoot, ultrafast bootstrap approximation tool (Hoang et al., 2018), and the SH-aLRT test (Guindon et al., 2010). The Bayesian topology was edited in FigTree (Rambaut, 2018), and ML branch supports were added to the final figure.

Effect of Cell-Free Filtrates on Coral Zooxanthellae

To investigate the allelopathic effects of cyanobacterial chemicals on *Symbiodinium* sp. CCMR0100 (ITS2 type A4) (*viz.* Silva-Lima et al., 2015) isolated from the coral *Mussismilia braziliensis* (Verrill, 1868) in the same reef system, we exposed the photosymbionts to cell-free filtrates of cultured *Moorea bouillonii* (L. Hoffmann and Demoulin) Engene Tronholm ($N = 4$) and *Adonis turfae* Walter et al. ($N = 5$), isolated from PAB_{BCM}. Strains were grown in an *f*/2 medium at 26°C, ca. 20 $\mu\text{E m}^{-2}\text{s}^{-1}$, and a 14:10-h light/dark photoperiod. Cyanobacterial cultures were grown for ca. 1 month, depending on the strain, until enough biomass was accumulated. The cells were then removed by filtration through glass microfiber filters (GF/F, 25 mm). Aliquots of 9 ml of filtrate from each cyanobacterial culture were distributed into quadruplicate 20 ml glass vials containing 9 ml of *Symbiodinium* sp. culture (harvested in the late exponential phase) in an *f*/2 medium. Controls were made by adding *f*/2 medium to the symbiont cultures instead of cyanobacteria culture filtrate. Two milliliters of fresh medium were added to each treatment and control vial to avoid nutrient limitation. Each experimental unit had an initial dinoflagellate concentration of ca. 6×10^4 cells ml^{-1} . The experiment was run for 72 h at temperature and irradiance as described above. Cell counts were performed daily either by flow cytometry (BD Accuri C6) or by microscopy using Palmer–Maloney counting chambers. The allelopathic effect was assessed daily by measuring the difference in symbiont cell number between each treatment and its respective control. Negative effects resulted in significantly fewer cells in the treatment exposed to the filtrate than in the control. Positive effects resulted in more cells in the treatment flasks (*i.e.*, greater cell number than the control).

Effect of Benthic Cyanobacterial Mat Extracts on Coral Zooxanthellae

To evaluate the allelopathic properties of secondary metabolites stored intracellularly in comparison with tests using naturally exuded allelochemicals (see previous section), we extracted a pooled biomass of PAB_{BCM} using organic solvents. Approximately 300 g (wet weight) of BCMs fronds was frozen in the field, transported to the laboratory, and freeze-dried. At that site, the BCMs is predominantly composed of *M. bouillonii*, *A. turfae*, *Halomiconema* sp., and *Leptolyngbya* sp. (Table 1). Exhaustive and successive organic extractions were done using ethyl acetate/methanol (1:1), yielding a homogeneous extract. Aliquots of this extract were solubilized in DMSO (dimethyl sulfoxide) and added to vials containing *Symbiodinium* sp. CCMR0100 to a final concentration proportional to the cyanobacteria biovolume used to prepare the extract, plus three dilutions by a factor of 10, that is, 10, 100, and 1,000 times of initial concentration. *Symbiodinium* cells were exposed to these extracts following the same experimental design as explained above for the cell-free filtrates. Cells were counted using an automated imaging system (FlowCam, Fluid Imaging Technologies, Inc.) before the experiment and after 48 h of incubation with the extracts to characterize the allelopathic effect of the cyanobacterial chemicals.

Benthic Cyanobacterial Mat Susceptibility to Herbivory

To evaluate the susceptibility of PAB_{BCM} to consumption and the potential influence of BCMs presence on consumer–prey interactions, we performed susceptibility assays of macroalgae and cyanobacteria in the field. We hypothesized that reef fish would refrain from consuming PAB_{BCM} filaments due to their high toxicity and/or low nutritional value. A small piece (5–10 cm) of fresh macroalgae and cyanobacteria was fixed to a 20-cm length 3-strand polypropylene rope with both ends fastened to a fishing weight, according to conventional methods (Hay et al., 1988), with modifications. *Halymenia* sp. was selected as a palatable macroalgae reference to evaluate BCMs consumption since it was readily consumed in previous tests (the authors, unpublished). Assays recorded grazing in three different treatments: a mix of macroalgae and BCMs, BCMs only, and macroalgae. A GoPRO camera was fastened to the bottom, and all treatments were placed within the camera's field of view. The experimental design was conceived after running pilot studies in which we repeatedly observed no consumption of stranded BCMs, and full consumption of *Halymenia* when offered separately to consumers. Due to such strong contrast, we added the mixed treatment to evaluate the grazing behavior toward *Halymenia* baits in face of unpalatable BCMs of similar color covering the thalli. We performed three sequential trials, using a single replicate for each treatment in every trial ($n = 3$) and recorded each one for 3 h, independent of the total consumption of any given treatment. The consumer species, number of bites, and time before consumption of the total biomass offered in a single bait were identified and recorded in each video.

TABLE 1 | Morphological and ecological features of the filaments found in the BCMs of Abrolhos Bank, Brazil.

Taxon	Site	Depth (m)	Depth BCMs features	Proportion	Filament color	Filament width (μm)	Sheath	Trichome width (μm)	Cell length (μm)	Constriction	Apical cell	BCMs isolates
<i>M. bouillonii</i>	PAB	5–8	Fasciculate mats, long pink-brown or flame-like appearance	Dominant	Red, brown, pink, salmon or light red	42–61	Firm, hyaline, thin, or thick, sometimes lamellated	37.1–45.4	8.6–11.3	No or slightly constricted	Rounded	CCMR174; PAB1-5
<i>Halomicronema</i> sp.	PAB	5–8	NA	Few	Pale pink	0.5–0.8	Colorless, very thin, joined	0.5–0.8	1.6–4.4	Constricted	Conical-rounded	CCMR085
<i>A. turfae</i>	PAB/PLES	5–8	NA	Few	Dark pink or violet	0.8–1.1	Colorless, very thin, joined	0.8–1.1	1.2–2.8	Constricted or slightly constricted	Conical-rounded	CCMR080;81; 82;83;86
<i>O. erythroflocculosa</i>	PLES	3–6	Dark pink Fasciculate mats	Dominant	Dark pink or red	59–66.6	Firm, thick, hyaline	49–58.9	5.3–6.7	Slightly constricted or not constricted	Rounded	PLES1-2
<i>Leptolyngbya</i> sp.	PAB	3–6	NA	Few	Pale olive-green	0.6–0.9	Firm or diffluent, thin, colorless, joined	0.6–0.9	1.2–2.6	Slightly constricted	Conical-rounded	CCMR087

Benthic Cyanobacterial Mat Toxicity to *Artemia salina*

Samples of PAB_{BCM} (i.e., *Moorena*-dominated tufts, see Table 1) were snap-frozen and taken to the laboratory. The toxicity was assessed by exposing *Artemia salina* nauplii within 24 h after eclosion to crude aqueous extracts of BCMs biomass. The BCMs biomass was harvested on 25-mm fiberglass filters (GF/C), moistened with seawater, and preweighted. Filters containing the biomass were immediately weighted to calculate the wet weight of the cyanobacteria samples. Filters were then transferred to 15-ml polystyrene vials filled with 8 ml of filtered seawater. Cyanobacteria cells were disrupted by three cycles of sonication (1 min each with pulses of 1 s, full power, in an ice bath) interspersed with freeze–thaw at -18°C . Then, the homogenates were filtered on 25-mm fiberglass filters (GF/C) to remove cell debris, 9 ml of AMF was added to 3 ml of the cell-free filtrates, and 6 ml of AMF was added into six 1:1 dilutions of the homogenate. Dilution series were prepared to achieve six concentrations from a max mean ($n = 11$) concentration of 26, 29 mg f.w. eq. ml^{-1} . Aliquots of 2 ml from each filtrate dilution were added to a 5-mL well plate in triplicates. Each well contained 2 ml of filtered seawater and 20 *A. salina* nauplii, which were observed after 48 h and scored as alive (motile) or dead (non-motile).

The specific growth rates of *Symbiodinium* cultures in incubations were calculated based on the cell concentrations, according to Equation 1:

$$\mu (d^{-1}) = \frac{(\ln N_1 - \ln N_0)}{t} \quad (1)$$

Where N_1 and N_0 are the cell concentrations at t_1 and t_0 , and t is the time, in days, between sampling occasions.

Statistical Analyses

Allelopathy data were analyzed with ANOVA with growth rate as response variable and treatment as fixed factor, followed by Tukey's *post hoc* test. Validity of equal variance, normality, and independence assumptions on the error terms was confirmed using the Shapiro–Wilk test and visual inspection of the plotted residuals. Susceptibility data were analyzed with the Kruskal–Wallis non-parametric test to check for differences between treatments with *Halymenia* sp. Toxicity was measured as $\text{LC}_{50-48\text{ h}}$ (median lethal concentration with 95% confidence intervals after 48 h of exposure) calculated by the Trimmed Spearman–Karber method (Hamilton et al., 1977). All statistical analyses were performed in the R environment (R Core Team, 2016).

RESULTS

Taxonomic Characterization of the Benthic Cyanobacterial Mats

The BCMs were predominantly composed of *M. bouillonii* or *O. erythroflocculosa*, non-ramified cyanobacteria species enclosed in exopolysaccharide sheaths and exhibit discoid cells


arranged in wide trichomes, corresponding to the Oscillatoriales order (Komárek and Anagnostidis, 2005). In addition, *Adonisiaturfæ* Walter et al., *Leptolyngbya* sp., and *Halomicronema* sp., were also found in the BCMs. However, they belong to the Synechococcales order, which has thin, non-ramified filaments, with cells longer than wide.


The BCMs from PAB_{BCM} presented different macroscopic morphologies and cyanobacteria taxonomic compositions than those collected from PLES_{BCM}. In PAB, the BCMs form fasciculate tufts of flame-like appearance, dark-red contours, and a pale inner region (Figure 2A), or as long, pink-brown fasciculate tufts (Figure 2C). These BCMs were predominantly composed of *M. bouillonii* (Figures 2D,E) and a few entangled filaments of *A. turfæ* (Figure 2H) and *Halomicronema* sp. (Figure 2I). In BCMs from PLES, *O. erythroflocculosa* was the predominant taxon (Figure 2F) with *Leptolyngbya* sp. and *A. turfæ* (Figures 2G,H), as relatively less abundant taxa, forming dark pink fasciculate, horizontally-spreading mats (Figure 2B and Table 1).

Phylogenetic inference (Figure 3) grouped the 16S rDNA sequences from broad filaments into two well-supported

clades (maximum likelihood, UF-bootstrap percentage/Bayesian analyses, posterior probability), *Moorena* (97/0.99) and *Okeania* (99/1.00). The *Moorena* clade included the isolates CCMR0174, PAB1, 2, 3, 4, and 5, identified as *M. bouillonii*. The sequences of PLES1 and PLES2, identified as *O. erythroflocculosa*, clustered together in the *Okeania* clade (98/0.96). The seven sequences from thin filament isolates were distributed within two clades, one represented by *Halomicronema* sp. (92/0.95), including the strain CCMR0085, and the other (–/0.81) composed of the remaining six isolates, identified as *A. turfæ* (CCMR0080, CCMR0081, CCMR0082, CCMR0083, and CCMR0086) and *Leptolyngbya* sp. (CCMR0087).

The 16S rDNA sequences presented a considerable similarity to others already published, such as PLES1 and PLES2, whose sequences showed 100% similarity with *O. erythroflocculosa* KC986930 (Engene et al., 2013b). The sequences of PAB1, 2, 3, 4, 5, and CCMR0174 were 99–100% similar to *M. bouillonii* GU111927 (Engene et al., 2012). Regarding the thin filaments, sequences of strains CCMR0080, CCMR0081, CCMR0082, CCMR0083, and CCMR0086 previously identified as *A. turfæ* had 100% similarity to *Leptolyngbya* sp. JX481735






(Konstantinou et al., 2019). The sequence of the strain CCMR0085 presented 99% similarity to *Halomicronema* sp. AB257773 (Yamazaki et al., 2008), which was found in the skeleton of the reef-building coral *Goniastrea aspera* Verrill, 1866. The sequence of the strain CCMR0087 was 99% similar to *Leptolyngbya* sp. EU743968, collected on the coral *Orbicella annularis* Ellis and Solander, 1786, which is one of the species responsible for causing Black band disease (BBD) (Gantar et al., 2009).

Allelopathic Effect of Cyanobacterial Exudates and Benthic Cyanobacterial Mat Extracts

Growth rates of the *ex hospite* *Symbiodinium* sp. exposed to cell-free filtrates (i.e., exudates) of the cultured *M. bouillonii* and *A. turfae* were significantly lower than those of corresponding controls (Supplementary Table 1 and Figure 4). Cell-free filtrates from cultures of the remaining taxa did not cause significant effects on *Symbiodinium*. Organic extracts of PAB_{BCM} significantly reduced the growth rates of *Symbiodinium* sp. at natural concentrations after a 48-h incubation (Supplementary Table 1 and Figure 5).



Benthic Cyanobacterial Mat Susceptibility to Herbivory

There was no BCMs consumption by fish in susceptibility field assays performed in PAB (Table 2, Figure 6, and




FIGURE 5 | Growth rates, $m (d^{-1})$, for the incubation with crude extract in the natural concentration (1), diluted 10x (0.1), diluted 100x (0.01) diluted 1000x (0.001). Error bars are standard errors. *Statistical significance compared to controls, $p < 0.05$. Asterisks represent significant p -values (alpha = 5%): $p < 0.001$ ***.

TABLE 2 | Number of bites in the susceptibility to herbivory tests of BCMs and macroalgae in single and mixed treatments.

Treatment	Total bites	Mean (SD)	CI (Low–Up)
BCM + Macroalgae	61	20 (\pm 24)	16–26
BCMs	0	NA	NA
Macroalgae	27	9 (\pm 2)	6–13

Supplementary Video). Treatments with only macroalgae were consumed first, before the *Halymenia* pieces in mixed treatments. All BCMs in the experiment remained untouched until the end of the trial. BCMs present in mixed treatments was avoided during delicate targeting of the macroalgae thalli. *Halymenia* sp. in macroalgae-only and mixed treatments were consumed in all trials compared with no-consumption of BCMs-only treatments (Table 2). There was no difference in the total number of bites between macroalgae treatments (chi-square = 0.2, $df = 1$, $p > 0.05$). The angelfish *Pomacanthus paru* and the surgeon fish *Acanthurus bahianus* were the primary consumers. They had distinct strategies to avoid BCMs while grazing the macroalgae in mixed treatments. *Pomacanthus paru* have a protruding feeding apparatus that allowed them to break the bait and selectively suck the pieces of interest floating in the water. The surgeon fishes have a smaller mouth and repeatedly bite the *Halymenia* pieces, taking out small fragments. Other eventual consumers were the juvenile of the parrotfish *Sparisoma axillare*, the yellowtail snapper *Ocyurus chrysurus*, and the damselfish *Stegastes fuscus*, which consumed *Halymenia* only in the mixed treatment.

Benthic Cyanobacterial Mat Toxicity

Three out of 11 PAB_{BCM} samples caused more than 50% mortality to *A. salina* nauplii (Supplementary Figure 1). The LC₅₀-48 h for these three samples was 9.8 mg w.w. eq. ml⁻¹ (95% CI: 8.3–11.6 mg w.w. eq. ml⁻¹), 9.3 mg w.w. eq. ml⁻¹ (95% CI: 6.8–12.6 mg w.w. eq. ml⁻¹), and 3.2 mg w.w. eq. ml⁻¹ (95% CI: 2.5–4.1 mg w.w. eq. ml⁻¹). The other eight samples had LC₅₀-48 h values higher than the maximum concentration used in tests (i.e., > 13.7–32.6 mg w.w. eq. ml⁻¹).

DISCUSSION


Morphological and Molecular Characterization of the Cyanobacteria

Morphological and genetic analyses of cyanobacteria in the BCMs unveiled two species, *M. bouillonii* and *O. erythroflocculosa*, hitherto unknown to the Abrolhos reefs and the tropical Southwestern Atlantic (SWA). Species belonging to both genera have been reported in Florida, the Caribbean, and the Pacific regions (Engene et al., 2012). *Okeania erythroflocculosa* has been detected as an abundant component of the BCMs worldwide (Engene et al., 2012, 2013a). Despite the high gene abundance of Oscillatoriales cyanobacteria previously reported for Abrolhos cyanobacterial mats (Walter et al., 2016), the species have not been identified to date. The species-level identification of cyanobacteria with thin filaments requires diacritic features that are difficult to observe. Their genotypic diversity can be high within the different phenotypes, characterizing them as cryptotaxa (Engene et al., 2010; Cairns et al., 2018). Here, the BCMs cryptotaxa were identified as *A. turfae*, previously reported (Walter et al., 2020), and two strains closely related to *Leptolyngbya* sp. and *Halomicronema* sp. The 16S rDNA sequences of the two latter cyanobacteria showed great similarities with the sequences found in the skeleton of scleractinian corals (Yamazaki et al., 2008; Gantar et al., 2009).

Allelopathic Effect of Cyanobacterial Exudates and Benthic Cyanobacterial Mat Extracts

Laboratory experiments evidenced allelopathic effects of chemical substances from cultured strains of cyanobacteria (*M. bouillonii* and *A. turfae*), expressed by inhibiting the growth of the *ex-hospite* *Symbiodinium* sp. isolated from *M. braziliensis*. This negative effect on *Symbiodinium* was also evidenced in incubations with organic extracts from BCMs collected in Abrolhos. Similar incubation bioassays reported a decrease in the number of *Chlorella vulgaris* cells after exposure to freshwater cyanobacteria exudates (Leão et al., 2009). Growth inhibition potential was also reported by the metabolites hapalindole from *Fischerella* sp. and calothrixin A from *Calothrix* sp., which affected growth and inhibited the genetic replication of prokaryotic and eukaryotic cells in bioassays (Doan et al., 2000). Also, exudates from the Baltic Sea cyanobacteria *Nodularia spumigena* and *Anabaena* sp. exhibited antagonistic effects on cryptophytes (Suikkanen et al., 2004). Cyanobacterin, a metabolite from *Scytonema hofmanni*, and fischerelin A, from *Fischerella muscicula*, were reported to affect the growth of photoautotrophs by inhibition of photosystem II (Gleason, 1990). Cyanobacteria extracts also exerted allelopathic effects in the field, manifested through a decreased pigment concentration of endosymbiotic dinoflagellates and the consequent bleaching of the *Porites* coral (Puyana et al., 2019).

The decrease in growth rate of *Symbiodinium* sp. cultures supports previous evidence of an antagonistic relationship between BCMs and reef builders in Abrolhos (Ribeiro et al., 2018). The effects shown here are likely involved in



interactions at the BCMs-coral boundary, which accumulate and transfer metabolites (Wangpraseurt et al., 2012). The increase in cyanobacterial biomass observed during blooms could result in thicker diffusion boundary layers that enhance transfer mechanisms and have antagonistic effects on coral photosynthesis and respiration (Jorissen et al., 2016). Such enhanced mass transfer through boundary layers and tissues (Patterson, 1992) could expose *Symbiodinium* to combined stress due to molecules produced by heterotrophs under anoxic conditions within the BCMs, and exuded allelochemicals diffused during blooms (Brocke et al., 2015). An increase in BCMs cover would also be a threat to coral recruitment success (Ritson-Williams et al., 2020). Additionally,

allelopathic effects during episodic cyanobacteria blooms can be further aggravated after acute disturbances such as heatwaves (Teixeira et al., 2019) when both the host and photosymbionts are physiologically impaired due to oxidative stress (Oakley and Davy, 2018).

Benthic Cyanobacterial Mat Toxicity and Susceptibility to Consumers

The mortality of brine shrimp after exposure to aqueous extracts of Abrolhos BCMs at natural cyanobacterial densities unveils the potential toxic properties of these benthic structures. Variable toxicity within the same BCMs that we observed (i.e., 3 out of

11 samples caused >50% mortality) should not imply a lack of chemical defense by cyanobacteria, as spatial and seasonal variability of chemical defenses against herbivores is expected due to metabolic responses to changes in the environment (Puglisi et al., 2019). The high spatial variability of Abrolhos reefs likely contributes to a context dependency of BCMs toxicity (Ribeiro et al., 2018; Teixeira et al., 2021). The large biomass of morbid sea slugs, *Stilochelylus striatus*, observed during cyanobacterial blooms in Abrolhos reefs (Ribeiro et al., 2017) is in line with the toxic potential of Abrolhos BCMs observed here. This opisthobranch is a specialist consumer of BCMs and has been shown to sequester toxic compounds from the cyanobacteria, becoming unpalatable to predators (Pennings and Paul, 1993).

Our results indicate that *O. erythroflocculosa*, *M. bouillonii*, *Leptolyngbya* sp., and *A. turfae* are potential producers of toxic compounds within Abrolhos BCMs. *Moorena*, *Okeania*, and *Lyngbya* species have been reported to produce more than 40% of the cyanobacteria secondary metabolites described to date (Engene et al., 2011; Dittmann et al., 2015), many of which mediate interactions with grazers (Thacker et al., 1997) and competitors (Kuffner et al., 2006; Titlyanov and Titlyanova, 2009). The *Leptolyngbya* consortium produces cyanotoxins and is associated with BBD in corals (Gantar et al., 2009).

Field assays of susceptibility to herbivory showed that BCMs remained untouched by consumers, either in treatments with BCMs alone or mixed with *Halymenia* sp., a more palatable macroalgae. Even though the genus *Halymenia* produces ecologically relevant secondary metabolites (Malik et al., 2020), the presence of these types of chemicals did not seem to influence algal consumption by surgeon fish elsewhere (Wylie and Paul, 1988). Among consumers, both *A. bahianus* and *S. axillare* are detritivorous but consume macroalgae and have a scraper feeding mode. These species display high foraging plasticity and can explore different food items, including cyanobacteria, and also calcareous and sheet-like macroalgae (Ferreira and Gonçalves, 2006; Mendes et al., 2018). The angelfish *P. paru* has a varied, omnivorous diet, encompassing sessile invertebrates and several groups of macroalgae (Batista et al., 2012). These species readily consumed the macroalgae treatments in every trial, with just a few bites given their relatively large size and protruding feeding apparatus. Although reef fish may consume benthic cyanobacteria (Ferreira and Gonçalves, 2006; Cissell et al., 2019), our exploratory results suggest avoidance of BCMs by reef fishes in PAB. The limited sample size of our feeding trials requires caution in interpreting such results and making generalizations of this pattern to BCMs in general. Given that a significant discrepancy between feeding selectivity and nutrient content of cyanobacteria was reported for reef herbivores (Mendes et al., 2018), BCMs consumption and deterrence seem to be a trade-off between its composition, growth form, and chemistry (Cissell and McCoy, 2021; Ford et al., 2021). The bottom-up effects on reef herbivores remain elusive as contrasting evidence has recently emerged (Cissell et al., 2019; Ford et al., 2021). Consumers with different feeding strategies

and nutritional demands may then target high nutritious cyanobacteria (Clements and Choat, 2018), and also associated non-cyanobacterial components (Cissell and McCoy, 2021), or avoid less nutritious (Nagarkar et al., 2004) and toxic cyanobacteria (Ford et al., 2021).

Phase shifts from coral dominated to opportunistic BCMs taxa have been reported worldwide (de Bakker et al., 2017; Ford et al., 2018), and such expansion disturbs reef trophic dynamics (Ford et al., 2021) by reducing energy flow and ultimately promotes a trophic simplification of the ecosystem (Ullah et al., 2018). This trophic disruption brings reef communities to alternative unstable states (Teixeira et al., 2021), because cyanobacteria are highly efficient in nutrient uptake and grow fast (den Haan et al., 2016; Ford et al., 2018), boosting their capacity to outcompete other benthic species and change community structure (e.g., de Bakker et al., 2017; Ribeiro et al., 2018; Teixeira et al., 2021). In the Abrolhos reef complex, BCMs can be structured by chemically defended species of cyanobacteria that display allelopathic, grazing deterrent, and toxic properties. These ecologically significant, albeit antagonistic, interactions with reef organisms could promote deleterious cascading effects on the ecosystem and contribute to the homogenization of reef communities. The disturbance of such trophic relations is probably amplified during episodic blooms of benthic cyanobacteria.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found below: <https://www.ncbi.nlm.nih.gov/genbank/>, mw248552 to 248559 and mw520745 to 520751.

AUTHOR CONTRIBUTIONS

FR designed the study and experiments, performed fieldwork, analyzed data, wrote the manuscript, and revised the manuscript. TC performed taxonomical work, analyzed data, and wrote the manuscript. LV performed phylogenetic and molecular analyses and wrote the manuscript. MS and PH contributed in genetic sequencing and microscopy. GF and AC performed microalgae cultures and incubation experiments. GP-F analyzed macroalgae and cyanobacteria collection and revised the manuscript. RM contributed with study design and sampling, materials, resources and revised the manuscript. RP and PS designed the study and experiments and wrote the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work has been supported by the following Brazilian research agencies: CNPq (grants 30755/82019/3 and 430819/2018-8 to PS), and the Rio de Janeiro Research Foundation (FAPERJ)

(grant E-26/202.911/2018). Partial funding was also provided by the Long-Term Ecological Research Program (PELD-Abrolhos/CNPq) and Fundação RENOVA/FEST.

ACKNOWLEDGMENTS

We thank the three reviewers for their insightful comments and suggestions that helped us to improve the manuscript. PS, RP, and RM would like to thank the Brazilian National Council for Scientific and Technological Development (CNPq) for Research Productivity fellowships. FR would like to thank “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior” (CAPES) for DSc. fellowship. We would like to thank Prof. Inácio Domingos from UFRJ for microscopy support.

REFERENCES

- Ainsworth, T. D., Hoegh-Guldberg, O., Heron, S. F., Skirving, W. J., and Leggat, W. (2008). Early cellular changes are indicators of pre-bleaching thermal stress in the coral host. *J. Exp. Mar. Biol. Ecol.* 364, 63–71. doi: 10.1016/j.jembe.2008.06.032
- Albert, S., O’Neil, J. M., Udy, J. W., Ahern, K. S., O’Sullivan, C. M., and Dennison, W. C. (2005). Blooms of the cyanobacterium *Lyngbya majuscula* in coastal Queensland, Australia: disparate sites, common factors. *Mar. Poll. Bull.* 51, 428–437. doi: 10.1016/j.marpolbul.2004.10.016
- Batista, D., Muricy, G. R. D. S., Andréa, B. R., and Villaçã, R. C. (2012). High intraspecific variation in the diet of the french angelfish *Pomacanthus paru* in the south-western Atlantic. *Braz. J. Oceanogr.* 60, 449–454.
- Birkeland, C. (1997). *Life and Death of Coral Reefs*. Berlin: Springer Science & Business Media.
- Brocke, H. J., Polerecky, L., De Beer, D., Weber, M., Claudet, J., and Nugues, M. M. (2015). Organic matter degradation drives benthic cyanobacterial mat abundance on Caribbean coral reefs. *PLoS One* 10:e0125445. doi: 10.1371/journal.pone.0125445
- Brumley, D., Spencer, K. A., Gunasekera, S. P., Sauvage, T., Biggs, J., Paul, V. J., et al. (2018). Isolation and characterization of anaerobes A-C, alkylphenols from a filamentous cyanobacterium (*Hormosira* sp., Oscillatoriales). *J. Nat. Prod.* 81, 2716–2721. doi: 10.1021/acs.jnatprod.8b00650
- Caires, T. A., Lyra, G. M., Hentschke, G. S., Pedrini, A. G., Sant’Anna, C. L., and Nunes, J. M. C. (2018). *Neolyngbya* gen. nov. (Cyanobacteria): a new benthic marine taxon widely distributed on the Brazilian Coast. *Mol. Phylogenet. Evol.* 120, 196–211. doi: 10.1016/j.ympev.2017.12.009
- Capper, A., and Paul, V. J. (2008). Grazer interactions with four species of *Lyngbya* in southeast Florida. *Harmful Algae* 7, 717–728. doi: 10.1016/j.hal.2008.02.004
- Capper, A., Erickson, A. A., Ritson-Williams, R., Becerro, M. A., Arthur, K. A., and Paul, V. J. (2016). Palatability and chemical defences of benthic cyanobacteria to a suite of herbivores. *J. Exp. Mar. Biol. Ecol.* 474, 100–108.
- Carroll, A. R., Copp, B. R., Davis, R. A., Keyzers, R. A., and Prinsep, M. R. (2021). Marine natural products. *Nat. Prod. Rep.* 38, 362–413. doi: 10.1039/D0NP00089B
- Chadwick, N. E., and Morrow, K. M. (2011). “Competition among sessile organisms on coral reefs,” in *Coral Reefs: An Ecosystem in Transition*, eds Z. Dubinsky and N. Stambler (Heidelberg: Springer).
- Charpy, L., Alliod, R., Rodier, M., and Golubic, S. (2007). Benthic nitrogen fixation in the SW New Caledonia lagoon. *Aquat. Microb. Ecol.* 47, 73–81. doi: 10.3354/ame047073
- Cissell, E. C., and McCoy, S. J. (2021). Shotgun metagenomic sequencing reveals the full taxonomic, trophic, and functional diversity of a coral reef benthic cyanobacterial mat from Bonaire, Caribbean Netherlands. *Sci. Total Environ.* 755:142719. doi: 10.1016/j.scitotenv.2020.142719

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmars.2021.790277/full#supplementary-material>

Supplementary Figure 1 | Concentration–response curves after 48 h of incubation of *Artemia salina* nauplii exposed to cell-free filtrates of PAB_{BCM}. Response expressed as average% mortality in each exposure concentration. LC₅₀-48 h is shown for the three samples where this end point was calculable by the Trimmed Spearman–Kärber method. The other samples have LC₅₀-48 h higher than the highest exposure concentration. Whiskers: standard error.

Supplementary Table 1 | Summary of ANOVA type II results for comparisons between incubation treatments and Tukey’s HSD *post hoc* test to verify differences in *Symbiodinium* growth index μ (d-1), between treatments and respective controls (without exudates).

Supplementary Video 1 | Susceptibility of benthic cyanobacteria to herbivory in field experiments.

- Cissell, E. C., Manning, J. C., and McCoy, S. J. (2019). Consumption of benthic cyanobacterial mats on a Caribbean coral reef. *Sci. Rep.* 9:12693.
- Clements, K. D., and Choat, J. H. (2018). “Nutritional ecology of parrotfishes (Scarinae, Labridae),” in *The Biology and Ecology of Parrotfishes*, eds A. S. Hoey and R. M. Bonaldo (Boca Raton, FL: CRC).
- Connell, S. D., Foster, M. S., and Airoldi, L. (2014). What are algal turfs? Towards a better description of turfs. *Mar. Ecol. Prog. Ser.* 495, 299–307. doi: 10.3354/meps10513
- de Bakker, D. M., Van Duyl, F. C., Bak, R. P., Nugues, M. M., Nieuwland, G., and Meesters, E. H. (2017). 40 Years of benthic community change on the Caribbean reefs of Curaçao and Bonaire: the rise of slimy cyanobacterial mats. *Coral Reefs* 36, 355–367. doi: 10.1007/s00338-016-1534-1539
- den Haan, J., Huisman, J., Brocke, H. J., Goehlich, H., Latijnhouwers, K. R., Van Heeringen, S., et al. (2016). Nitrogen and phosphorus uptake rates of different species from a coral reef community after a nutrient pulse. *Sci. Rep.* 6: 28821.
- Dittmann, E., Gugger, M., Sivonen, K., and Fewer, D. P. (2015). Natural product biosynthetic diversity and comparative genomics of the cyanobacteria. *Trends Microbiol.* 23, 642–652. doi: 10.1016/j.tim.2015.07.008
- Doan, N. T., Rickards, R. W., Rothschild, J. M., and Smith, G. D. (2000). Allelopathic actions of the alkaloid 12-epi-hapalindole E isonitrile and calothrixin A from cyanobacteria of the genera *Fischerella* and *Calothrix*. *J. Appl. Phycol.* 12, 409–416. doi: 10.1023/A:1008170007044
- Echenique-Subiabre, L., Villeneuve, A., Golubic, S., Turquet, J., Humbert, J.-F., and Gugger, M. (2015). Influence of local and global environmental parameters on the composition of cyanobacterial mats in a tropical lagoon. *Microb. Ecol.* 69, 234–244. doi: 10.1007/s00248-014-0496-490
- Engene, N., Choi, H., Esquenazi, E., Rottacker, E. C., Ellisman, M. H., Dorrestein, P. C., et al. (2011). Underestimated biodiversity as a major explanation for the perceived rich secondary metabolite capacity of the cyanobacterial genus *Lyngbya*. *Environ. Microbiol.* 13, 1601–1610. doi: 10.1111/j.1462-2920.2011.02472.x
- Engene, N., Coates, R. C., and Gerwick, W. H. (2010). 16S rRNA gene heterogeneity in the filamentous marine cyanobacterial genus *Lyngbya*. *J. Phycol.* 46, 591–601. doi: 10.1111/j.1529-8817.2010.00840.x
- Engene, N., Gunasekera, S. P., Gerwick, W. H., and Paul, V. J. (2013a). Phylogenetic inferences reveal a large extent of novel biodiversity in chemically rich tropical marine cyanobacteria. *Appl. Environ. Microb.* 79, 1882–1888. doi: 10.1128/AEM.03793-3712
- Engene, N., Paul, V. J., Byrum, T., Gerwick, W. H., Thor, A., and Ellisman, M. H. (2013b). Five chemically rich species of tropical marine cyanobacteria of the genus *Okeania* gen. nov. (Oscillatoriales, Cyanoprokaryota). *J. Phycol.* 49, 1095–1106. doi: 10.1111/jpy.12115
- Engene, N., Rottacker, E. C., Kaštovský, J., Byrum, T., Choi, H., Ellisman, M. H., et al. (2012). *Moorea producens* gen. nov., sp. nov. and *Moorea bouillonii* comb.

- nov., tropical marine cyanobacteria rich in bioactive secondary metabolites. *Int. J. Syst. Evol. Microbiol.* 62, 1171–1178. doi: 10.1099/ijms.0.033761-33760
- Engene, N., Tronholm, A., and Paul, V. J. (2018). Uncovering cryptic diversity of *Lyngbya*: the new tropical marine cyanobacterial genus *Dapis* (Oscillatoriales). *J. Phycol.* 54, 435–446. doi: 10.1111/jpy.12752
- Ferreira, C. E. L., and Gonçalves, J. E. A. (2006). Community structure and diet of roving herbivorous reef fishes in the Abrolhos Archipelago, south-western Atlantic. *J. Fish Biol.* 69, 1533–1551. doi: 10.1111/j.1095-8649.2006.01220.x
- Fistarol, G. O., Hargreaves, P. L., Walter, J. M., Viana, T. V., Gomes, P. D. F., Lourenço, C. B., et al. (2018). Rapid isolation of culturable microalgae from a tropical shallow lake system. *J. Appl. Phycol.* 30, 1807–1819.
- Ford, A. K., Bejarano, S., Nugues, M. M., Visser, P. M., Albert, S., and Ferse, S. C. A. (2018). Reefs under Siege - the rise, putative drivers, and consequences of benthic cyanobacterial mats. *Front. Mar. Sci.* 5:18. doi: 10.3389/fmars.2018.00018
- Ford, A. K., Visser, P. M., van Herk, M. J., Jongepier, E., and Bonito, V. (2021). First insights into the impacts of benthic cyanobacterial mats on fish herbivory functions on a nearshore coral reef. *Sci. Rep.* 11:7147. doi: 10.1038/s41598-021-84016-z
- Gantar, M., Sekar, R., and Richardson, L. L. (2009). Cyanotoxins from black band disease of corals and from other coral reef environments. *Microb. Ecol.* 58, 856–864. doi: 10.1007/s00248-009-9540-x
- Gleason, F. K. (1990). The natural herbicide, cyanobacterin, specifically disrupts thylakoid membrane structure in *Euglena gracilis* strain. *FEMS Microbiol. Lett.* 68, 77–81.
- Graham, L. E., and Wilcox, L. W. (2000). *Algae*. Hoboken, NJ: Prentice Hall.
- Groendahl, S., and Fink, P. (2017). High dietary quality of non-toxic cyanobacteria for a benthic grazer and its implications for the control of cyanobacterial biofilms. *BMC Ecol.* 17:20. doi: 10.1186/s12898-017-0130-3
- Guillard, R. R. L. (1975). "Culture of phytoplankton for feeding marine invertebrates," in *Culture of Marine Invertebrate Animals*, eds W. L. Smith and M. H. Chanley (New York, NY: Plenum Press), 26–60.
- Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–332. doi: 10.1093/sysbio/syq010
- Hamilton, M. A., Russo, R. C., and Thurston, R. V. (1977). Trimmed Spearman-Kärber method for estimating median lethal concentrations in toxicity bioassays. *Environ. Sci. Tech.* 11, 714–719.
- Hay, M. E., Paul, V. J., Lewis, S. M., Gustafson, K., Tucker, J., and Trindell, R. N. (1988). Can tropical seaweeds reduce herbivory by growing at night? diel patterns of growth, nitrogen content, herbivory, and chemical versus morphological defenses. *Oecologia* 75, 233–245. doi: 10.1007/BF00378604
- Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q., and Vinh, L. S. (2018). UFBoot2: improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.* 35, 518–522. doi: 10.1093/molbev/msx281
- Jorissen, H., Skinner, C., Osinga, R., Beer, D., and Nugues, M. M. (2016). Evidence for water-mediated mechanisms in coral-algal interactions. *Proc. Biol. Sci.* 283, 1–10. doi: 10.1098/rspb.2016.1137
- Komárek, J., and Anagnostidis, K. (2005). "Cyanoprokaryota-2. Teil/2nd part: *Oscillatoriales*," in *Süsswasserflora von Mitteleuropa 19/778 2*, eds B. Büdel, G. Gärtner, L. Krienitz, and M. Schagerl (Heidelberg: Elsevier/Spektrum).
- Konstantinou, D., Voultziadou, E., Panteris, E., Zervou, S.-K., Hiskia, A., and Gkelis, S. (2019). Leptothoe, a new genus of marine cyanobacteria (Synechococcales) and three new species associated with sponges from the Aegean Sea. *J. Phycol.* 55, 882–897. doi: 10.1111/jpy.12866
- Kuffner, I. B., Walters, L. J., Becerro, M., Paul, V. J., Ritson-Williams, R., and Beach, K. (2006). Inhibition of coral recruitment by macroalgae and cyanobacteria. *Mar. Ecol. Prog. Ser.* 323, 107–117. doi: 10.3354/meps323107
- Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* 35, 1547–1549.
- Lane, D. J. (1991). "16S/23S rRNA sequencing," in *Nucleic Acid Techniques in Bacterial Systematics*, eds E. Stackebrandt and M. Goodfellow (New York, NY: John Wiley and Sons).
- Leão, P. N., Engene, N., Antunes, A., Gerwick, W. H., and Vasconcelos, V. (2012). The chemical ecology of cyanobacteria. *Nat. Prod. Rep.* 29, 372–391. doi: 10.1093/molbev/msy096
- Leão, P. N., Vasconcelos, M. T. S. D., and Vasconcelos, V. M. (2009). Allelopathic activity of cyanobacteria on green microalgae at low cell densities. *Eur. J. Phycol.* 44, 347–355. doi: 10.1080/09670260802652156
- Leão, Z. M., Kikuchi, R. K. P., Ferreira, B. P., Neves, E. G., Sovierzoski, H. H., Oliveira, M. D., et al. (2016). Brazilian coral reefs in a period of global change: a synthesis. *Braz. J. Oceanogr.* 64, 97–116.
- Malik, S. A. A., Bedoux, G., Robledo, D., García-Maldonado, J. Q., Frelle-Pelegrin, Y., and Bourgougnon, N. (2020). Chemical defense against microfouling by allelopathic active metabolites of *Halymenia foresii* (Rhodophyta). *J. Appl. Phycol.* 32, 2673–2687. doi: 10.1007/s10811-020-02094-2094
- Mendes, T. C., Ferreira, C. E. L., and Clements, K. D. (2018). Discordance between diet analysis and dietary macronutrient content in four nominally herbivorous fishes from the Southwestern Atlantic. *Mar. Biol.* 165:180. doi: 10.1007/s00227-018-3438-3434
- Morrow, K. M., Paul, V. J., Liles, M. R., and Chadwick, N. E. (2011). Allelochemicals produced by Caribbean macroalgae and cyanobacteria have species-specific effects on reef coral microorganisms. *Coral Reefs* 30, 309–320. doi: 10.1007/s00338-011-0747-741
- Nagarkar, S., Williams, G. A., Subramanian, G., and Saha, S. K. (2004). "Cyanobacteria-dominated biofilms: a high quality food resource for intertidal grazers," in *Proceedings of the Asian Pacific Phycology in the 21st Century: Prospects and Challenges* (Dordrecht: Springer).
- Nicholson, G. M., and Clements, K. D. (2020). Resolving resource partitioning in parrotfishes (Scarini) using microhistology of feeding substrata. *Coral Reefs* 39, 1313–1327.
- Nübel, U., Garcia-Pichel, F., and Muyzer, G. (1997). PCR primers to amplify 16S rRNA genes from cyanobacteria. *Appl. Environ. Microbiol.* 63, 3327–3332.
- Oakley, C. A., and Davy, S. K. (2018). "Cell biology of coral bleaching," in *Coral Bleaching. Ecological Studies (Analysis and Synthesis)*, eds M. Van Oppen and J. Lough (Cham: Springer).
- Patterson, M. R. (1992). A mass transfer explanation of metabolic scaling relations in some aquatic invertebrates and algae. *Science* 255, 1421–1423. doi: 10.1126/science.255.5050.1421
- Paul, V. J., Thacker, R. W., Banks, K., and Golubic, S. (2005). Benthic cyanobacterial bloom impacts the reefs of South Florida (Broward County, USA). *Coral Reefs* 24, 693–697. doi: 10.1007/s00338-005-0061-x
- Pennings, S. C., and Paul, V. J. (1993). Sequestration of dietary secondary metabolites by three species of sea hares: location, specificity and dynamics. *Mar. Biol.* 117, 535–546. doi: 10.1007/BF00349763
- Puglisi, M. P., Sneed, J. M., Ritson-Williams, R., and Young, R. (2019). Marine chemical ecology in benthic environments. *Nat. Prod. Rep.* 36, 410–429. doi: 10.1039/C8NP00061A
- Puyana, M., and Prato, J. (2013). Overgrowth of reef organisms by benthic cyanobacteria in the Colombian Caribbean. *Mutis* 3, 58–60. doi: 10.21789/22561498.885
- Puyana, M., Prato, J. A., Nieto, C. F., Ramos, F. A., Castellanos, L., Pinzón, P., et al. (2019). Experimental approaches for the evaluation of allelopathic interactions between hermatypic corals and marine benthic cyanobacteria in the Colombian Caribbean. *Acta Biol. Colomb.* 24, 243–254. doi: 10.15446/abc.v24n2.72706
- Rambaut, A. (2018). *FigTree, Version 1.4. 4. Program Distributed by the Author*. Institute of Evolutionary Biology: University of Edinburgh.
- R Core Team (2016). *R: a Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Ribeiro, F. V., Padula, V., Moura, R. L., Moraes, F. C., Salomon, P. S., Gibran, F. Z., et al. (2017). Massive opisthobranch aggregation in the largest coralline reefs in the South Atlantic Ocean: are mesoherbivores underestimated top-down players? *Bull. Mar. Sci.* 93, 915–916. doi: 10.5343/bms.2016.110
- Ribeiro, F. V., Sá, J. A., Fistarol, G. O., Salomon, P. S., Pereira, R. C., Souza, M. L. A. M., et al. (2018). Long-term effects of competition and environmental drivers on the growth of the endangered coral *Mussismilia braziliensis* (Verrill, 1867). *PeerJ* 6:e5419. doi: 10.7717/peerj.5419
- Ritson-Williams, R., Arnold, S. N., and Paul, V. J. (2020). The impact of macroalgae and cyanobacteria on larval survival and settlement of the scleractinian corals *Acropora palmata*, *A. cervicornis* and *Pseudodiploria strigosa*. *Mar. Biol.* 167:31.

- Ritson-Williams, R., Paul, V. J., and Bonito, V. (2005). Marine benthic cyanobacteria overgrow coral reef organisms. *Coral Reefs* 24:629. doi: 10.1007/s00338-005-0059-54
- Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., et al. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–542. doi: 10.1093/sysbio/sys029
- Silva-Lima, A. W., Walter, J. M., Garcia, G. D., Ramires, N., Ank, G., Meirelles, P. M., et al. (2015). Multiple *Symbiodinium* strains are hosted by the Brazilian endemic corals *Mussismilia* spp. *Microb. Ecol.* 70, 301–310. doi: 10.1007/s00248-015-0573-z
- Sneed, J. M., Meickle, T., Engene, N., Reed, S., Gunasekera, S., and Paul, V. J. (2017). Bloom dynamics and chemical defenses of benthic cyanobacteria in the Indian River Lagoon, Florida. *Harmful Algae* 69, 75–82. doi: 10.1016/j.hal.2017.10.002
- Suikkanen, S., Fistarol, G. O., and Granéli, E. (2004). Allelopathic effects of the Baltic cyanobacteria *Nodularia spumidigena*, *Aphanizomenon flosaquae* and *Anabaena lemmermannii* on algal monocultures. *J. Exp. Mar. Biol. Ecol.* 308, 85–101. doi: 10.1016/j.jembe.2004.02.012
- Teixeira, C. D., Chiroque-Solano, P. M., Ribeiro, F. V., Carlos-Júnior, L. A., Neves, L. M., Salomon, P. S., et al. (2021). Decadal (2006–2018) dynamics of Southwestern Atlantic's largest turbid zone reefs. *PLoS One* 16:e0247111. doi: 10.1371/journal.pone.0247111
- Teixeira, C. D., Leitão, R. L., Ribeiro, F. V., Moraes, F. C., Neves, L. M., Bastos, A. C., et al. (2019). Sustained mass coral bleaching (2016–2017) in Brazilian turbid-zone reefs: taxonomic, cross-shelf and habitat-related trends. *Coral Reefs* 38, 801–813. doi: 10.1007/s00338-019-01789-1786
- Thacker, R. W., Becerro, M. A., Lumbang, W. A., and Paul, V. J. (1998). Allelopathic interactions between sponges on a tropical reef. *Ecology* 79, 1740–1750. doi: 10.2307/176792
- Thacker, R. W., Nagle, D. G., and Paul, V. J. (1997). Effects of repeated exposures to marine cyanobacterial secondary metabolites on feeding by juvenile rabbitfish and parrotfish. *Mar. Ecol. Prog. Ser.* 147, 21–29.
- Titlyanov, E. A., and Titlyanova, T. V. (2009). The dynamics of the restoration of mechanical damage to colonies of the scleractinian coral *Porites lutea* under conditions of competition with algal settlers for substratum. *Russ. J. Mar. Biol.* 35, 230–235. doi: 10.1134/S1063074009030067
- Trifinopoulos, J., Nguyen, L. T., von Haeseler, A., and Minh, B. Q. (2016). W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 44, W232–W235. doi: 10.1093/nar/gkw256
- Tronholm, A., and Engene, N. (2019). *Moorena* gen. nov., a valid name for “*Moorea* Engene & al.” nom. inval. (Oscillatoriaceae, Cyanobacteria). *Not. Algar.* 122, 1–2.
- Ullah, H., Nagelkerken, I., Goldenberg, S. U., and Fordham, D. A. (2018). Climate change could drive marine food web collapse through altered trophic flows and cyanobacterial proliferation. *PLoS Biol.* 16:e2003446. doi: 10.1371/journal.pbio.2003446
- Walter, J. M., Coutinho, F. H., Leomil, L., Hargreaves, P. I., Campeão, M. E., Vieira, V. V., et al. (2020). Ecogenomics of the marine benthic filamentous cyanobacterium *Adonisia*. *Microb. Ecol.* 80, 249–265. doi: 10.1007/s00248-019-01480-x
- Walter, J. M., Tschoeke, D. A., Meirelles, P. M., de Oliveira, L., Leomil, L., Tenório, M., et al. (2016). Taxonomic and functional metagenomic signature of turfs in the Abrolhos reef system (Brazil). *PLoS One* 11:e0161168. doi: 10.1371/journal.pone.0161168
- Wangpraseurt, D., Weber, M., Røy, H., Polerecky, L., de Beer, D., Suharsono, et al. (2012). In situ oxygen dynamics in coral-algal interactions. *PLoS One* 7:e31192. doi: 10.1371/journal.pone.0031192
- Wylie, C. R., and Paul, V. J. (1988). Feeding preferences of the surgeonfish *Zebbrasoma flavescens* in relation to chemical defenses of tropical algae. *Mar. Ecol. Prog. Ser.* 45, 23–32. doi: 10.3354/meps045023
- Yamamoto, M. (1999). Importance of epiphytic cyanobacteria as food sources for heterotrophs in a tropical seagrass bed. *Coral Reefs* 18, 263–271.
- Yamazaki, S. S., Nakamura, T., Yuen, Y. S., and Yamasaki, H. (2008). “Reef-building coral *Goniastrea aspera* harbor a novel filamentous cyanobacterium in their skeleton,” in *Proceedings of the 11th International Coral Reef Symposium* (Ft. Lauderdale, FL).
- Yang, G. Y., and Dudgeon, D. (2010). Dietary variation and food selection by an algivorous loach (*Pseudogastromyzon myersi*: Balitoridae) in Hong Kong streams. *Mar. Freshwater Res.* 61, 49–56.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Ribeiro, Caires, Simões, Hargreaves, Villela, Fistarol, Cazalgrandi Junior, Pereira-Filho, Moura, Pereira and Salomon. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.