



OPEN ACCESS

EDITED AND REVIEWED BY
Eduardo Almansa,
Spanish Institute of Oceanography
(IEO), Spain

*CORRESPONDENCE
Qihui Yang
qihuiyang03@163.com

SPECIALTY SECTION
This article was submitted to
Marine Fisheries, Aquaculture and
Living Resources,
a section of the journal
Frontiers in Marine Science

RECEIVED 23 August 2022
ACCEPTED 15 September 2022
PUBLISHED 23 September 2022

CITATION
Lin H, Tan B, Yang Q, Chi S, Wei H,
Wu Y, Ray GW and Yohana MA (2022)
Corrigendum: Effects of dietary
glycerol monolaurate on growth,
antioxidant capacity and lipid
metabolism in cage-farmed pompano
(*Trachinotus ovatus*) juveniles.
Front. Mar. Sci. 9:1025794.
doi: 10.3389/fmars.2022.1025794

COPYRIGHT
© 2022 Lin, Tan, Yang, Chi, Wei, Wu,
Ray and Yohana. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Corrigendum: Effects of dietary glycerol monolaurate on growth, antioxidant capacity and lipid metabolism in cage-farmed pompano (*Trachinotus ovatus*) juveniles

Huaxing Lin ^{1,2,3}, Beiping Tan ^{1,2,3}, Qihui Yang ^{1,2,3*},
Shuyan Chi ^{1,2,3}, Hongjie Wei ^{1,2,3}, Yuancai Wu ^{1,2,3},
Gyan Watson Ray ^{1,2} and Mpwaga Alatwinusa Yohana ^{1,2}

¹College of Fisheries, Guangdong Ocean University, Zhanjiang, China, ²Aquatic Animals Precision Nutrition and High Efficiency Feed Engineering Research Center of Guangdong Province, College of Fisheries, Guangdong Ocean University, Zhanjiang, China, ³Guangdong Provincial Key Laboratory of Aquatic Animal Disease Control and Healthy Culture, College of Fisheries, Guangdong Ocean University, Zhanjiang, China

KEYWORDS

Trachinotus ovatus, glycerol monolaurate, growth performance, antioxidant capacity, lipid metabolism

A Corrigendum on

Effects of dietary glycerol monolaurate on growth, antioxidant capacity and lipid metabolism in cage-farmed pompano (*Trachinotus ovatus*) juveniles

by Lin H, Tan B, Yang Q, Chi S, Wei H, Wu Y, Ray GW and Yohana MA (2022) *Front. Mar. Sci.* 9:914134. doi: 10.3389/fmars.2022.914134

In the published article, there was an error in [Table 1](#) as published. Due to an accidental input error during the data entry process, the sum of each group did not equal 100. In [Table 1](#), Vitamin premix and Mineral premix should both be 0.50%. The corrected [Table 1](#) and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

TABLE 1 Composition and nutrient levels of the experimental diets (dry matter basis, %).

Ingredients	Diets					
	G0	G5	G10	G15	G20	G25
Fish meal	25.00	25.00	25.00	25.00	25.00	25.00
Chicken meal	8.00	8.00	8.00	8.00	8.00	8.00
Soybean meal	12.00	12.00	12.00	12.00	12.00	12.00
Peanut meal	8.00	8.00	8.00	8.00	8.00	8.00
Corn gluten meal	8.00	8.00	8.00	8.00	8.00	8.00
Wheat flour	20.00	20.00	20.00	20.00	20.00	20.00
CaH ₂ PO ₄	1.50	1.50	1.50	1.50	1.50	1.50
Vitamin C	0.05	0.05	0.05	0.05	0.05	0.05
Choline chloride	0.30	0.30	0.30	0.30	0.30	0.30
Soy lecithin	0.50	0.50	0.50	0.50	0.50	0.50
Fish oil	1.50	1.50	1.50	1.50	1.50	1.50
Soy oil	1.50	1.50	1.50	1.50	1.50	1.50
Vitamin premix ^a	0.50	0.50	0.50	0.50	0.50	0.50
Mineral premix ^b	0.50	0.50	0.50	0.50	0.50	0.50
DL-Methionine	0.46	0.46	0.46	0.46	0.46	0.46
L-Lysine	0.72	0.72	0.72	0.72	0.72	0.72
L-Threonine	0.24	0.24	0.24	0.24	0.24	0.24
L-Arginine	0.26	0.26	0.26	0.26	0.26	0.26
Microcrystalline cellulose	10.97	10.92	10.87	10.82	10.77	10.72
Glyceryl monolaurate (GML)	0.00	0.05	0.10	0.15	0.20	0.25
Total	100.00	100.00	100.00	100.00	100.00	100.00
Nutrient levels						
Crude protein ^c	40.67	40.73	39.86	40.52	40.55	39.93
Crude lipid ^c	12.18	12.56	12.26	12.35	12.41	12.37
Crude ash ^c	7.90	7.79	7.89	7.96	7.79	7.88
Nitrogen free extract ^c	21.91	21.94	22.86	21.78	21.87	22.28
Gross energy (MJ/kg) ^c	18.08	18.18	18.17	18.10	18.06	17.99

^aThe vitamin premix: vitamin A, 500,000 IU/kg; vitamin D3, 100,000 IU/kg; vitamin E, 4,000 mg/kg; vitamin K3, 1,000 mg/kg; vitamin B1, 500 mg/kg; vitamin B2, 1,000 mg/kg; vitamin B6, 1,000 mg/kg; vitamin B12, 2.0 mg/kg; nicotinic acid, 4,000 mg/kg; D-calcium pantothenate, 2,000 mg/kg; folic acid, 100 mg/kg; biotin, 10.0 mg/kg; vitamin C, 15,000 mg/kg;

^bMineral mixture: Fe, 10,000 mg/kg; Cu, 300 mg/kg; Zn, 5,000 mg/kg; Mn, 1,200 mg/kg; I, 80 mg/kg; Se, 30 mg/kg; Co, 20 mg/kg.

^cCrude protein, crude lipid, crude ash, nitrogen free extract and gross energy were measured values.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.