Check for updates

OPEN ACCESS

EDITED AND REVIEWED BY Lei Wang, Anhui Normal University, China

*CORRESPONDENCE Qiang Chen Chenqiang@zjou.edu.cn Tao Han goodhantao@gmail.com

RECEIVED 25 December 2024 ACCEPTED 06 January 2025 PUBLISHED 16 January 2025

CITATION

Chen S, Dai J, Chen Y, Chen Q, Dong F, Wang C, Sun Y, Wang J and Han T (2025) Corrigendum: Effects of *Bacillus subtilis*fermented soybean meal replacing fish meal on antioxidant activity, immunity, endoplasmic reticulum stress and hepatopancreas histology in Pacific white shrimp (*Litopenaeus vannamei*). *Front. Mar. Sci.* 12:1551241. doi: 10.3389/fmars.2025.1551241

COPYRIGHT

© 2025 Chen, Dai, Chen, Chen, Dong, Wang, Sun, Wang and Han. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms. Corrigendum: Effects of *Bacillus subtilis*-fermented soybean meal replacing fish meal on antioxidant activity, immunity, endoplasmic reticulum stress and hepatopancreas histology in Pacific white shrimp (*Litopenaeus vannamei*)

Songming Chen¹, Jieyu Dai¹, Yan Chen², Qiang Chen^{1*}, Fen Dong¹, Congcong Wang¹, Yulong Sun¹, Jiteng Wang¹ and Tao Han^{1*}

¹Department of Aquaculture, Zhejiang Ocean University, Zhoushan, China, ²Zhejiang Provincial Engineering Technology Research Center of Marine Biomedical Products, School of Food and Pharmacy, Zhejiang Ocean University, Zhoushan, China

KEYWORDS

antioxidant activity, endoplasmic reticulum stress, fermented soybean meal, immunity, Litopenaeus vannamei

A Corrigendum on

Effects of *Bacillus subtilis*-fermented soybean meal replacing fish meal on antioxidant activity, immunity, endoplasmic reticulum stress and hepatopancreas histology in Pacific white shrimp (*Litopenaeus vannamei*)

By Chen S, Dai J, Chen Y, Chen Q, Dong F, Wang C, Sun Y, Wang J and Han T (2024) *Front. Mar. Sci.* 11:1449066. doi: 10.3389/fmars.2024.1449066

In the published article, there was an error in Figure 2 and its legend as published. The Figure 2 in the published version lacks the content of Figures 2H–S. The corrected Figure 2 and its legend appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 2

The effects of FSBM substitution on the antioxidant indices in different tissues and the contents of T-CHO and TG in the hemolymph (n = 4). The results are presented as the mean \pm S. E. M. and were analyzed with Tukey's tests (bars bearing the same letters are not significantly different among treatments (P > 0.05)]. The same as below. (A) GPT (glutamic-pyruvic transaminase) activity in the hemolymph; (B) GOT (glutamic oxalacetic transaminase) activity in the hemolymph; (C) GSH (reduced glutathione) content in the hemolymph; (D) SOD (superoxide dismutase) activity in the hemolymph; (C) GSH (reduced glutathione) content in the hemolymph; (D) SOD (superoxide dismutase) activity in the hemolymph; (E) T-AOC (total antioxidant capacity) activity in the hemolymph; (F) T-CHO (total cholesterol) content in the hemolymph; (G) TG (triacylglycerol) content in the hemolymph; (H) SOD activity in the hepatopancreas; (I) MDA (malondialdehyde) content in the hepatopancreas; (J) GSH content in the hepatopancreas; (K) T-AOC activity in the hepatopancreas; (L) SOD activity in the muscle; (M) MDA content in the muscle; (N) GSH content in the intestinal; (G) T-AOC activity in the muscle; (P) SOD activity in the intestinal; (G) MDA content in the intestinal; (R) GSH content in the intestinal; (S) T-AOC activity in the intestinal. n. s., not significant.