



## OPEN ACCESS

## EDITED AND REVIEWED BY

Pavel Michalek,  
General University Hospital in  
Prague, Czechia

## \*CORRESPONDENCE

Michael Koeppen  
michael.koeppen@  
med.uni-tuebingen.de

## SPECIALTY SECTION

This article was submitted to  
Intensive Care Medicine and  
Anesthesiology,  
a section of the journal  
Frontiers in Medicine

RECEIVED 10 November 2022

ACCEPTED 14 November 2022

PUBLISHED 23 November 2022

## CITATION

Sari-Yavuz S, Heck-Swain K-L, Keller M,  
Magunia H, Feng Y-S, Haeberle HA,  
Wied P, Schlensak C, Rosenberger P  
and Koeppen M (2022) Corrigendum:  
Methylene blue dosing strategies in  
critically ill adults with shock—A  
retrospective cohort study.  
*Front. Med.* 9:1094735.  
doi: 10.3389/fmed.2022.1094735

## COPYRIGHT

© 2022 Sari-Yavuz, Heck-Swain, Keller,  
Magunia, Feng, Haeberle, Wied,  
Schlensak, Rosenberger and Koeppen.  
This is an open-access article  
distributed under the terms of the  
[Creative Commons Attribution License  
\(CC BY\)](#). The use, distribution or  
reproduction in other forums is  
permitted, provided the original  
author(s) and the copyright owner(s)  
are credited and that the original  
publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or  
reproduction is permitted which does  
not comply with these terms.

# Corrigendum: Methylene blue dosing strategies in critically ill adults with shock—A retrospective cohort study

Sibel Sari-Yavuz<sup>1</sup>, Ka-Lin Heck-Swain<sup>1</sup>, Marius Keller<sup>1</sup>,  
Harry Magunia<sup>1</sup>, You-Shan Feng<sup>2</sup>, Helene A. Haeberle<sup>1</sup>,  
Petra Wied<sup>1</sup>, Christian Schlensak<sup>3</sup>, Peter Rosenberger<sup>1</sup> and  
Michael Koeppen<sup>1\*</sup>

<sup>1</sup>Department of Anesthesiology and Intensive Care Medicine, University Hospital Tübingen, Tübingen, Germany, <sup>2</sup>Institute for Clinical Epidemiology and Applied Biostatistics (IKEaB), Eberhard-Karls-University Tübingen, Tübingen, Germany, <sup>3</sup>Department of Thoracic and Cardiovascular Surgery, University Hospital Tübingen, Tübingen, Germany

## KEYWORDS

shock, methylene blue, critical care, retrospective study, clinical trial

## A corrigendum on

### Methylene blue dosing strategies in critically ill adults with shock—A retrospective cohort study

by Sari-Yavuz, S., Heck-Swain, K.-L., Keller, M., Magunia, H., Feng, Y.-S., Haeberle, H. A., Wied, P., Schlensak, C., Rosenberger, P., and Koeppen, M. (2022). *Front. Med.* 9:1014276. doi: 10.3389/fmed.2022.1014276

In the published article, there was an error. The dosing of the continuous infusion of methylene blue is incorrect. Currently the text states “continuous infusion with 0.25 mg/kg/min” this should be changed to “continuous infusion with 0.25 mg/kg/h”.

A correction has been made to **Methods**, *Methylene blue therapy*, second paragraph. This sentence previously stated:

“Three different dosing strategies have been used in our institution in the past: bolus injection alone, continuous infusion without bolus, injection, or bolus injection followed by continuous infusion (bolus 2 mg/kg; continuous infusion with 0.25 mg/kg/min).”

The corrected sentence appears below:

“Three different dosing strategies have been used in our institution in the past: bolus injection alone, continuous infusion without bolus, injection, or bolus injection followed by continuous infusion (bolus 2 mg/kg; continuous infusion with 0.25 mg/kg/h).”

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.