



Protective host defense against disseminated candidiasis is impaired in mice expressing human interleukin-37

Frank L. van de Veerdonk^{1,2,3}*, Mark S. Gresnigt², Marije Oosting², Jos W. M. van der Meer², Leo A. B. Joosten^{2,3}, Mihai G. Netea^{2,3} and Charles A. Dinarello^{1,2}

¹ Department of Medicine, University of Colorado Denver, Denver, CO, USA

² Department of Medicine, Radboud University Nijmegen Medical Center, Nijmegen, Netherlands

³ Radboud Center for Infection, Nijmegen, Netherlands

Edited by:

Annelies Sophie Zinkernagel,
University Hospital Zurich—University
of Zurich, Switzerland

Reviewed by:

Thomas Tsaganos, University of
Athens Medical School, Greece
Stylianos Orfanos, University of
Athens Medical School, Greece

*Correspondence:

Frank L. van de Veerdonk, Department
of Medicine, Radboud University
Nijmegen Medical Center, Geert
Grooteplein Zuid 8, 6525 GA
Nijmegen, Netherlands
e-mail: f.veerdonk@aig.umcn.nl

The effect of the anti-inflammatory cytokine interleukin-37 (IL-37) on host defense against *Candida* infections remains unknown. We assessed the role of IL-37 in a murine model of disseminated candidiasis using mice transgenic for human IL-37 (hIL-37Tg). Upon exposure to *Candida albicans* pseudohyphae, macrophages from hIL-37Tg mice release 39% less TNF α compared to cells from wild-type (WT) mice ($p = 0.01$). *In vivo*, hIL-37Tg mice displayed a decreased capacity to recruit neutrophils to the site of infection. These defects were associated with increased mortality and organ fungal growth in hIL-37Tg compared to WT mice. We conclude that IL-37 interferes with the innate protective anti-*Candida* host response by reducing the production of proinflammatory cytokines and suppressing neutrophil recruitment in response to *Candida*, resulting in an increased susceptibility to disseminated candidiasis.

Keywords: IL-37, *Candida*, TNF α , inflammation, neutrophils, antifungal host defense, IL-17

INTRODUCTION

The interleukin-1 (IL-1) family includes inflammatory cytokines (such as IL-1 α , IL-1 β , IL-18, and IL-33) that activate target cells through a receptor-mediated mechanism; anti-inflammatory cytokine antagonists (such as IL-1Ra) block IL-1-dependent activation by competing for binding to the IL-1 receptors (Dinarello, 2011). Twelve years ago, using *in silico* searches, additional members of the IL-1 family namely IL-36, IL-37, and IL-38 were discovered, but only recently have their properties been studied (Dinarello, 2013). In the case of IL-37, this cytokine has emerged as fundamental inhibitor of innate inflammation using mice transgenic for human IL-37 as well as reducing endogenous IL-37 in human blood monocytes (Nold et al., 2010). For example, mice transgenic for human IL-37 are protected against systemic endotoxemia, chemical induced colitis, ConA hepatitis, and ischemic reperfusion injury, as reviewed in Dinarello and Bufler (2013). The mechanism by which IL-37 affords anti-inflammatory protection can be via a caspase-1-dependent nuclear translocation or by an extracellular mechanism engaging the IL-18 receptor (Bulau et al., 2014). It is important however, to recognize that anti-inflammatory strategies based upon IL-37 could have a negative impact on susceptibility to infections by inhibiting host defense. Since in models of live infection blocking inflammation mediated by cytokines such as IL-1 or TNF α can be either detrimental or beneficial for the outcome of the host, we therefore carried-out studies of disseminated candidiasis in mice expressing human IL-37.

MATERIALS AND METHODS

ANIMALS

Transgenic mice expressing human IL-37 (hIL-37Tg) have been previously described Nold et al. (2010). Control C57BL/6J (WT)

mice were purchased from the Jackson Laboratory and maintained under specific-pathogen free conditions. The mice were fed sterilized laboratory chow (Hope Farms, Woerden, The Netherlands) and water *ad libitum*. The experiments were approved by the Ethics Committee on Animal Experiments of Radboud University Nijmegen.

Candida albicans AND GROWTH CONDITIONS

Candida albicans ATCC MYA-3573 (UC 820), a strain well described elsewhere (Lehrer and Cline, 1969), was used in all experiments. *Candida* was grown overnight in Sabouraud broth at 37°C, cells were harvested by centrifugation, washed twice, and resuspended in culture medium (RPMI-1640 Dutch modification, ICN Biomedicals, Aurora, OH, USA; van der Graaf et al., 2005). To generate pseudohyphae, *C. albicans* blastoconidia were grown at 37°C in culture medium, adjusted to pH 6.4 by using hydrochloric acid. Pseudohyphae were killed at 100°C for 1 h and resuspended in culture medium to a hyphal inoculum size that originated from 10⁶ blastoconidia per ml (referred to as 10⁶ pseudohyphae per ml).

CYTOKINE RESPONSES OF *C. albicans*-STIMULATED MACROPHAGES AND SPLENCYTES

Resident peritoneal macrophages were harvested from groups of five hIL-37Tg and wild-type (WT) mice by injecting 4 ml of sterile phosphate-buffered saline (PBS) intraperitoneal containing 0.38% sodium citrate (Kullberg et al., 1993). After washing, the cells were resuspended in culture medium in 96-well microtiter plates (Greiner, Alphen, The Netherlands) at 10⁵ cells/well, in a final volume of 200 μ l. The cells were stimulated with either control medium or heat killed *C. albicans* at 1 \times 10⁷ microorganisms/ml. After 24 h of incubation at 37°C,

the plates were centrifuged (500 g, 10 min), and the supernatants were removed. The cells were lysed with three freeze-thaw cycles. Samples were stored at -80°C until cytokine assays were performed.

To assess cytokine production in splenocytes, spleen were gently squeezed into a sterile 200 mm filter chamber. The cells were washed and resuspended in RPMI 1640, counted in a Bürker chamber and adjusted to $5 \times 10^6/\text{ml}$. 200 μL of the cell suspension was stimulated with 1×10^7 heat-killed *C. albicans*/ml. Measurement of TNF α and IL-6 concentrations was performed in supernatants collected after 48 h of incubation at 37°C in 5% CO_2 in 48-well plate.

CYTOKINE ASSAYS

TNF α was determined by specific radioimmunoassay (detection limit 20 pg/ml), as previously described Netea et al. (1996). IL-6 concentrations were determined by a commercial ELISA (Biosource, Camarillo, CA, USA, detection limit 16 pg/ml) according to the instructions of the manufacturer.

C. albicans INFECTION MODEL

hIL-37Tg mice and WT mice were injected intravenously with *C. albicans* blastoconidia (1×10^6 CFU/mouse for survival and 5×10^5 CFU/mouse for testing fungal burden) in a 100 μl volume of sterile pyrogen-free PBS. Survival was assessed daily for 2 weeks. Subgroups of five animals were killed on days 3 and 7 of infection. To assess the tissue outgrowth of the microorganisms, the kidneys of the sacrificed animals were removed aseptically, weighed, and homogenized in sterile saline in a tissue grinder. The number of viable *Candida* cells in the tissues was determined by plating serial dilutions on Sabouraud dextrose agar plates. The CFU were counted after 24 h of incubation at 37°C , and expressed as log CFU/g tissue.

Candida OUTGROWTH IN PMN DEPLETED MICE

Sexually mature female C57bl/6 mice (8–12 weeks old, weights 20–25 g) were used. The animals were kept under routine laboratory conditions ($21\text{--}22^{\circ}\text{C}$, relative humidity 60% and a 12 h light–dark cycle), fed a standard commercial pellet diet (RHM, Hope Farms, The Netherlands), and given acidified tap water *ad libitum*. The mice were injected intraperitoneal with rat anti-mouse GR1 or isotype control (100 μg in 200 μl PBS; eBioscience) 4 days prior to the experiment to deplete the mice of PMNs. On day zero, the mice were injected intraperitoneal with 1×10^5 live *C. albicans* conidia. After 2 h the mice were anesthetized with isoflurane and blood samples were obtained for measurement of blood cells counts. Subsequently, the mice were sacrificed by cervical dislocation. The number of viable *Candida* cells in the peritoneal lavage was determined by plating serial dilutions on Sabouraud dextrose agar plates and the CFU were counted after 24 h of incubation at 37°C and expressed as log CFU.

IN VITRO CYTOKINE PRODUCTION BY PRIMED SPLENCYTES

To assess cytokine production during infection, primed spleen cells from mice on day 7 of infection with 2×10^5 CFU of *C. albicans* per mouse were stimulated *in vitro* with heat-killed *Candida* conidia or pseudohyphae (1×10^6 microorganisms/ml). Spleen

cells were washed and resuspended in RPMI1640, counted in a Bürker chamber and the number was adjusted to $5 \times 10^6/\text{ml}$. 200 μL of the cell suspension was stimulated with 1×10^6 heat killed *C. albicans*/ml. Measurement of IFN γ , IL-10, and IL-17 concentrations was performed in supernatants collected after 48 h of incubation at 37°C in 5% CO_2 in 48-well plate.

PHAGOCYTOSIS AND KILLING OF *C. albicans* BY MACROPHAGES

Phagocytosis and killing of *C. albicans* blastoconidia were assessed as described elsewhere (Vonk et al., 2006). Exudate peritoneal phagocytes from groups of five hIL-37Tg mice and WT mice were elicited by an i.p. injection of 10% proteose peptone. After 72 h, cells were collected in separate sterile tubes by washing the peritoneal cavity with 4 ml of ice-cold PBS that contained 50 U/ml heparin. Phagocytes were centrifuged (for 10 min at 2250 g), counted in a Bürker chamber, and resuspended in culture medium. 5×10^5 cells were dispensed into the wells of a 96-well flat bottom plate (Costar), allowed to adhere for 2 h, and washed to remove non-adherent cells. Subsequently, the cells were incubated with 1×10^4 CFU *C. albicans*, which were opsonized for 45 min at 24°C in modified Eagle's medium (MEM; Gibco Life Technologies) containing 2.5% fresh mouse serum (effector:target ratio, 40:1). After 15 min, supernatants were aspirated, and monolayers were gently washed with MEM to remove non-ingested microorganisms. The supernatant and well washings that contained the non-ingested *Candida* blastoconidia were plated in serial dilutions on Sabouraud agar plates. The percentage of phagocytosed microorganisms was defined as $[1 - (\text{number of uningested CFU}/\text{CFU at the start of incubation})] \times 100$. Killing of *C. albicans* by phagocytes was assessed in the same monolayers. After removal of the non-phagocytized *Candida* blastoconidia, 200 μl of culture medium, consisting of Sabouraud in MEM (50% vol/vol), was added to the monolayers. After 3 h of incubation at 37°C in air and 5% CO_2 , the wells were scraped gently with a plastic paddle and washed with 200 μl distilled H_2O to achieve lysis of phagocytes in three cycles, and 10-fold dilutions of each sample were spread on Sabouraud agar plates and incubated at 37°C for 24 h. The percentage of yeast killed by the phagocytes was determined as follows: $[1 - (\text{CFU after incubation}/\text{number of phagocytized CFU})]$ (Nold et al., 2010) $\times 100$. Phagocyte-free incubations of blastoconidia were included as a control for yeast viability. In earlier experiments we have shown that 90–97% of attached/internalized *C. albicans* are intracellular (Vonk et al., 2006).

STATISTICAL ANALYSIS

The differences between groups were analyzed by the Mann–Whitney *U*-test. The level of significance between groups was set at $p < 0.05$. The data are presented as cumulative results of all experiments performed.

RESULTS

hIL-37Tg MICE ARE MORE SUSCEPTIBLE TO DISSEMINATED *C. albicans* INFECTION

To investigate the role of IL-37 in the host defense against invasive *C. albicans* infection, hIL-37Tg mice, and WT mice were infected with *C. albicans* blastospores. The survival during disseminated candidiasis was 20% for WT-type mice after 2 weeks, whereas all

mice had died by day 10 in the hIL-37Tg group (**Figure 1A**). The fungal burden in the kidneys, the target organ of disseminated candidiasis in mice (Spellberg et al., 2005), was not significantly different at day 3 (mean \pm SD; WT: 3.6 ± 0.83 , IL-37Tg 3.8 ± 0.44 log CFU/g kidney), and 1-log higher in the hIL-37Tg mice than in WT mice on day 7 of infection ($p < 0.05$; **Figure 1B**). Histology revealed an accumulation of *Candida* in the pyelum of the kidneys of hIL-37Tg mice which was not observed in the WT mice (**Figure 1C**).

NEUTROPHIL RECRUITMENT AND PHAGOCYTOSIS AND KILLING OF *C. albicans*

In order to understand the enhanced lethality and greater fungal burden of the hIL-37Tg mice, we investigated the dynamics of the influx and function of neutrophils. Peritoneal cells were harvested and counted 4 h after intraperitoneal injection of heat-killed *Candida*. A significant lower influx of neutrophils was apparent 4 h after challenge with heat killed *Candida* in the hIL37Tg mice compared to WT mice (**Figure 2A**). Phagocytosis of *C. albicans* by hIL-37Tg cells was similar to that by cells of WT mice (37 vs. 39% phagocytized in 15 min; $p > 0.05$). hIL-37Tg cells killed 92% of phagocytized *Candida* blastoconidia after 3 h, which was not different from the killing activity of cells from WT cells (88%, $p > 0.05$, **Figure 2B**).

To assess whether the decrease of neutrophils within the first hours has a significant biological effect, we investigated the difference of fungal burden between neutrophil depleted mice and non-neutrophil depleted mice in a short *in vivo* *Candida* infection model. The lack of neutrophils at the site of infection resulted in a log increase in fungal growth within 2 h compared to mice in which neutrophil influx was normal (**Figure 2C**). These differences support the importance of early influx of neutrophils in restricting *Candida* growth *in vivo*.

IL-37 INHIBITS PROINFLAMMATORY CYTOKINE PRODUCTION BY MACROPHAGES AND SPLENCYTES STIMULATED WITH *C. albicans*

To investigate the inhibitory effect of IL-37 at the level of cytokine production, resident peritoneal macrophages, or naïve splenocytes


of hIL-37Tg and WT mice were exposed to heat-killed *C. albicans* blastoconidia and pseudohyphae *in vitro*. Cytokine production by unstimulated macrophages of each mouse strain was below the detection limit for TNF α and IL-6. However, after stimulation with *C. albicans* pseudohyphae, the production of TNF α was significantly lower in macrophages from the hIL-37Tg mice compared to cells from WT mice. There were no differences in the production of IL-6 in the same cultures (**Figure 3**).



hIL-37Tg MICE DISPLAY INCREASED Th17 AND IL-10 RESPONSES

Proinflammatory T helper responses such as Th1 and Th17 responses play an important role in fungal host defense (Gow et al., 2012). The Th17 response during disseminated fungal infection can be both protective and detrimental (Huang et al., 2004; Zelante et al., 2007; Lin et al., 2009). To study the impact of IL-37 on proinflammatory adaptive immune responses, we investigated the Th1 and Th17 characteristic cytokines IFN γ and IL-17. Splenocytes isolated from hIL-37Tg mice with a higher fungal burden at day 7 produced significant more IL-17 in response to *C. albicans* pseudohyphae (**Figure 4**). IFN γ production in response to *C. albicans* pseudohyphae was undetectable in WT and hIL-37Tg splenocytes at day 7 (**Figure 4**). Since IL-37 mainly has anti-inflammatory effects (Nold et al., 2010; Boraschi et al., 2011), we investigated levels of IL-10 in cells from IL-37Tg. There was no significant difference in IL-10 production in splenocytes from hIL-37Tg compared to WT cells in response to *C. albicans*, although hIL-37Tg mice showed a trend toward an increased IL-10 production (**Figure 4**).

DISCUSSION

In the present study, we demonstrate that overexpression of IL-37 is detrimental for the early host defense against *C. albicans* in a murine model of disseminated candidiasis. This conclusion is based on the increased fungal load in the kidney, the target organ of disseminated candidiasis, in hIL-37Tg mice compared to WT mice. The greater outgrowth of the fungus in the tissues is most probably due to impaired early influx of neutrophils into the infected tissues, which in turn may be due to lower production of neutrophil-inducing cytokines, such as TNF. Therefore, the






presence of IL-37 in the early stages of infection results in an impaired innate immune response that is essential to limit fungal dissemination.

Recently, several biological functions of the cytokine IL-37 have been described. IL-37 has a protective role in a murine model

of lethal endotoxemia (Nold et al., 2010; Bulau et al., 2011), and it has been reported that mice overexpressing IL-37 are protected from DSS colitis (McNamee et al., 2011). These reports point to significant anti-inflammatory properties of IL-37. In the present study we extend these findings to a live infection



model, the murine model of disseminated candidiasis, in which we assessed whether IL-37 hampers the innate immune defense against a fungal infection. We observed that mice overexpressing IL-37 were more susceptible to invasive candidiasis, due to the inability to control fungal growth in the kidney during infection. This effect is at least partly due to a defective neutrophil recruitment, a cell population central for the local antifungal host defense (Kullberg et al., 1990), and this supports the previously reported finding that leukocyte recruitment to the site of inflammation is decreased in hIL-37Tg mice (McNamee et al., 2011). The importance of neutrophils in the control of *Candida* outgrowth *in vivo* is further supported in this study by the observation that neutrophil depleted mice have a log increase within 2 h in fungal burden compared to mice that have a normal neutrophil influx. In addition, macrophages and splenocytes overexpressing IL-37 produced significantly less TNF α than WT macrophages when exposed to *C. albicans*. This is in line with other studies, showing that macrophages from hIL-37Tg mice produce less TNF and that IL-37 overexpression reduces TNF-dependent inflammation in a murine model of colitis (McNamee et al., 2011). An involvement of defective TNF production in the increased susceptibility of hIL-37Tg mice to disseminated candidiasis is also consistent with earlier studies showing that mice lacking TNF or the TNF receptor are highly susceptible to disseminated candidiasis (Netea et al., 1999).

Another observation of this study is that neutrophils from hIL-37Tg mice do not display an intrinsic defect in phagocytosis and killing, suggesting that IL-37 does not directly impair neutrophil function. The increased Th17 response in hIL-37Tg mice present on day 7 most likely reflects an increased cellular immune response due to higher fungal burden (antigen) exposure. Whether this increased Th17 response has directly contributed to the higher susceptibility of hIL-37Tg mice to disseminated candidiasis remains to be determined. Interestingly, hIL-37Tg splenocytes showed a consistent trend at day 7 toward an increased production of the anti-inflammatory cytokine IL-10 when exposed to *C. albicans*. This observation is consistent with the increased IL-10 production found in hIL-37Tg mice

with experimental colitis, and supports the concept that IL-37 can contribute to systemic anti-inflammatory effects (McNamee et al., 2011).

In conclusion, IL-37 reduces proinflammatory cytokine production induced by *Candida* in macrophages, and decreases neutrophil recruitment *in vivo* in response to *C. albicans*. These data highlight the potent anti-inflammatory effects of IL-37, and underline that the timing of IL-37 expression in the tissue at the site of inflammation is critical for the outcome of the host during inflammatory processes.

ACKNOWLEDGMENTS

This study was supported by the Niels Stensen Foundation and a Veni Grant of the Netherlands Foundation for Scientific Research to Frank L. van de Veerdonk. Charles A. Dinarello was supported by NIH grant AI-15614.

REFERENCES

- Boraschi, D., Luchesi, D., Hainzl, S., Leitner, M., Maier, E., Mangelberger, D., et al. (2011). IL-37: a new anti-inflammatory cytokine of the IL-1 family. *Eur. Cytokine Netw.* 22, 127–147. doi: 10.1684/ecn.2011.0288
- Bulau, A. M., Fink, M., Maucksch, C., Kappler, R., Mayr, D., Wagner, K., et al. (2011). In vivo expression of interleukin-37 reduces local and systemic inflammation in concanavalin a-induced hepatitis. *Sci. World J.* 11, 2480–2490. doi: 10.1100/2011/968479
- Bulau, A. M., Nold, M. F., Li, S., Nold-Petry, C. A., Fink, M., Mansell, A., et al. (2014). Role of caspase-1 in nuclear translocation of IL-37, release of the cytokine, and IL-37 inhibition of innate immune responses. *Proc. Natl. Acad. Sci. U.S.A.* 111, 2650–2655. doi: 10.1073/pnas.1324140111
- Dinarello, C. A. (2011). Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. *Blood* 117, 3720–3732. doi: 10.1182/blood-2010-07-273417
- Dinarello, C. A. (2013). Overview of the interleukin-1 family of ligands and receptors. *Semin. Immunol.* 25, 389–393. doi: 10.1016/j.smim.2013.10.001
- Dinarello, C. A., and Bufler, P. (2013). Interleukin-37. *Semin. Immunol.* 25, 466–468. doi: 10.1016/j.smim.2013.10.004
- Gow, N. A., van de Veerdonk, F. L., Brown, A. J., and Netea, M. G. (2012). *Candida albicans* morphogenesis and host defence: discriminating invasion from colonization. *Nat. Rev. Microbiol.* 10, 112–122. doi: 10.1038/nrmicro2711
- Huang, W., Na, L., Fidel, P. L., and Schwarzenberger, P. (2004). Requirement of interleukin-17A for systemic anti-*Candida albicans* host defense in mice. *J. Infect. Dis.* 190, 624–631. doi: 10.1086/422329
- Kullberg, B. J., van 't Wout, J. W., Hoogstraten, C., and Van Furth, R. (1993). Recombinant interferon- γ enhances resistance to acute disseminated *Candida albicans* infection in mice. *J. Infect. Dis.* 168, 436–443. doi: 10.1093/infdis/168.2.436
- Kullberg, B. J., van 't Wout, J. W., and Van Furth, R. (1990). Role of granulocytes in enhanced host resistance to *Candida albicans* induced by recombinant interleukin-1. *Infect. Immun.* 58, 3319–3324.
- Lehrer, R. I., and Cline, M. J. (1969). Interactions of *Candida albicans* with human leukocytes and serum. *J. Bacteriol.* 98, 996–1004.
- Lin, L., Ibrahim, A. S., Xu, X., Farber, J. M., Avanesian, V., Baquir, B., et al. (2009). Th1-Th17 cells mediate protective adaptive immunity against *Staphylococcus aureus* and *Candida albicans* infection in mice. *PLoS Pathog.* 5:e1000703. doi: 10.1371/journal.ppat.1000703
- McNamee, E. N., Masterson, J. C., Jedlicka, P., McManus, M., Grenz, A., Collins, C. B., et al. (2011). Interleukin 37 expression protects mice from colitis. *Proc. Natl. Acad. Sci. U.S.A.* 108, 16711–16716. doi: 10.1073/pnas.1111982108
- Netea, M. G., Demacker, P. N. M., Kullberg, B. J., Boerman, O. C., Verschuere, I., Stalenhoef, A. F., et al. (1996). Low-density-lipoprotein receptor deficient mice are protected against lethal endotoxemia and severe gram-negative infections. *J. Clin. Invest.* 97, 1366–1372. doi: 10.1172/JCI118556
- Netea, M. G., Van Tits, L. J. H., Curfs, J. H. A. J., Amiot, F., Meis, J. F. G. M., van der Meer, J. W. M., et al. (1999). The increased susceptibility of TNF α LT α

- double knock-out mice to systemic candidiasis is due to defective recruitment and phagocytosis by neutrophils. *J. Immunol.* 163, 1498–1505.
- Nold, M. F., Nold-Petry, C. A., Zepp, J. A., Palmer, B. E., Bufler, P., and Dinarello, C. A. (2010). IL-37 is a fundamental inhibitor of innate immunity. *Nat. Immunol.* 11, 1014–1022. doi: 10.1038/ni.1944
- Spellberg, B., Ibrahim, A. S., Edwards, J. E. Jr., and Filler, S. G. (2005). Mice with disseminated candidiasis die of progressive sepsis. *J. Infect. Dis.* 192, 336–343. doi: 10.1086/430952
- van der Graaf, C. A. A., Netea, M. G., Verschueren, I., van der Meer, J. W. M., and Kullberg, B. J. (2005). Differential cytokine production and Toll-like receptor signaling pathways by *Candida albicans* blastoconidia and hyphae. *Infect. Immun.* 73, 7458–7464. doi: 10.1128/IAI.73.11.7458-7464.2005
- Vonk, A. G., Netea, M. G., van Krieken, J. H., Iwakura, Y., van der Meer, J. W., and Kullberg, B. J. (2006). Endogenous interleukin (IL)-1 alpha and IL-1 beta are crucial for host defense against disseminated candidiasis. *J. Infect. Dis.* 193, 1419–1426. doi: 10.1086/503363
- Zelante, T., De Luca, A., Bonifazi, P., Montagnoli, C., Bozza, S., Moretti, S., et al. (2007). IL-23 and the Th17 pathway promote inflammation and impair antifungal immune resistance. *Eur. J. Immunol.* 37, 2695–2706. doi: 10.1002/eji.200737409
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 28 October 2014; accepted: 15 December 2014; published online: 07 January 2015.

Citation: van de Veerdonk FL, Gresnigt MS, Oosting M, van der Meer JWM, Joosten LAB, Netea MG and Dinarello CA (2015) Protective host defense against disseminated candidiasis is impaired in mice expressing human interleukin-37. *Front. Microbiol.* 5:762. doi: 10.3389/fmicb.2014.00762

This article was submitted to *Infectious Diseases*, a section of the journal *Frontiers in Microbiology*.

Copyright © 2015 van de Veerdonk, Gresnigt, Oosting, van der Meer, Joosten, Netea and Dinarello. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.