



# Cyanobacteria in Sulfidic Spring Microbial Mats Can Perform Oxygenic and Anoxygenic Photosynthesis Simultaneously during an Entire Diurnal Period

Judith M. Klatt<sup>1,2\*</sup>, Dirk de Beer<sup>1</sup>, Stefan Häusler<sup>1</sup> and Lubos Polerecky<sup>1,3</sup>

<sup>1</sup> Microsensor Group, Max-Planck-Institute for Marine Microbiology, Bremen, Germany, <sup>2</sup> Geomicrobiology Lab, Department of Earth and Environmental Sciences, University of Michigan, Ann Arbor, MI, USA, <sup>3</sup> Department of Earth Sciences – Geochemistry, Faculty of Geosciences, Utrecht University, Utrecht, Netherlands

## OPEN ACCESS

### Edited by:

Robert Eugene Blankenship,  
Washington University in St. Louis,  
USA

### Reviewed by:

Min Chen,  
University of Sydney, Australia  
Dmitry A. Los,  
Institute of Plant Physiology (RAS),  
Russia

### \*Correspondence:

Judith M. Klatt  
jmklatt@umich.edu

### Specialty section:

This article was submitted to  
Microbial Physiology and Metabolism,  
a section of the journal  
Frontiers in Microbiology

**Received:** 02 September 2016

**Accepted:** 24 November 2016

**Published:** 15 December 2016

### Citation:

Klatt JM, de Beer D, Häusler S and  
Polerecky L (2016) Cyanobacteria  
in Sulfidic Spring Microbial Mats Can  
Perform Oxygenic and Anoxygenic  
Photosynthesis Simultaneously during  
an Entire Diurnal Period.  
Front. Microbiol. 7:1973.  
doi: 10.3389/fmicb.2016.01973

We used microsensors to study the regulation of anoxygenic and oxygenic photosynthesis (AP and OP, respectively) by light and sulfide in a cyanobacterium dominating microbial mats from cold sulfidic springs. Both photosynthetic modes were performed simultaneously over all H<sub>2</sub>S concentrations (1–2200 μM) and irradiances (4–52 μmol photons m<sup>-2</sup> s<sup>-1</sup>) tested. AP increased with H<sub>2</sub>S concentration while the sum of oxygenic and anoxygenic photosynthetic rates was constant at each light intensity. Thus, the total photosynthetically driven electron transport rate was solely controlled by the irradiance level. The partitioning between the rates of these two photosynthetic modes was regulated by both light and H<sub>2</sub>S concentration. The plastoquinone pool (PQ) receives electrons from sulfide:quinone:reductase (SQR) in AP and from photosystem II (PSII) in OP. It is thus the link in the electron transport chain where both pathways intersect, and the compound that controls their partitioning. We fitted our data with a model of the photosynthetic electron transport that includes the kinetics of plastoquinone reduction and oxidation. The model results confirmed that the observed partitioning between photosynthetic modes can be explained by a simple kinetic control based on the affinity of SQR and PSII toward PQ. The SQR enzyme and PSII have similar affinities toward PQ, which explains the concurrent OP and AP over an astonishingly wide range of H<sub>2</sub>S concentrations and irradiances. The elegant kinetic control of activity makes the cyanobacterium successful in the fluctuating spring environment. We discuss how these specific regulation mechanisms may have played a role in ancient H<sub>2</sub>S-rich oceans.

**Keywords:** microbial mat, cyanobacteria, anoxygenic photosynthesis, sulfide:quinone:reductase, hydrogen sulfide, microsensors

## INTRODUCTION


Oxygenic photosynthesis (OP) is a process where light energy is used to extract electrons from water to reduce CO<sub>2</sub>. The evolution of this type of photosynthesis was predated by anoxygenic photosynthesis (AP), a process that uses another compound (e.g., H<sub>2</sub>S) as electron donor for the reduction of CO<sub>2</sub> (Blankenship, 2010). While the more ancient AP requires only one photosystem (PSI) to drive the electron flow, OP requires

the combined power of two photosystems (PSI+PSII), primarily because of the high energy demand in the water splitting reaction.

OP probably evolved in cyanobacteria (Mulkidjanian et al., 2006) inhabiting microbial mat-like structures. In these systems alternative electron donors for photosynthesis such as  $H_2S$  were likely abundant (Nisbet and Fowler, 1999; Buick, 2008) and remained available until the end of the Proterozoic (Canfield, 1998). Therefore, it has been hypothesized that AP performed by obligate anoxygenic phototrophs and cyanobacteria capable of both AP and OP (referred to as versatile cyanobacteria) was an important photosynthetic mode on a global scale before the complete oxygenation of the Earth's atmosphere and oceans during the Neoproterozoic oxidation event (Johnston et al., 2009). It is therefore intriguing how oxygenic phototrophs were finally so successful in oxygenating Earth despite the widespread availability of electron donors for anoxygenic phototrophs and the toxic effects of  $H_2S$  on the components of OP, especially considering that AP is a biochemically less complicated and energetically less demanding process than OP (Miller and Bebout, 2004; Klatt et al., 2015a).

New insights into possible mechanisms that allowed outcompetition of anoxygenic phototrophs by oxygenic phototrophs in the presence of  $H_2S$  can be gained by studying adaptations of extant cyanobacteria living in sulfidic environments. The cold, light-exposed sulfidic springs at Frasassi, Italy (Klatt et al., 2016), which harbor thin microbial mats inhabited by diverse anoxygenic, oxygenic and versatile phototrophs, are one example of such an environment. Our recent studies of cyanobacteria isolated from this system revealed two novel cyanobacterial adaptations to  $H_2S$ . The first adaptation mechanism, observed for a versatile cyanobacterium *Pseudoanabaena* str. FS39, included partitioning between AP and OP that is regulated kinetically by  $H_2S$  through the apparent affinity of the electron transport components involved in AP [sulfide:quinone:reductase (SQR)] and OP [photosystem II (PSII)] toward plastoquinone, which is where both electron transport pathways intersect (Figure 1; Klatt et al., 2016). OP in this cyanobacterial strain is active only when the electron transport chain is not fully used by AP, which occurs when  $H_2S$  is limiting. The second type of adaptation, observed for the obligatory oxygenic cyanobacterium *Planktothrix* str. FS34, included acceleration of the recovery of OP by  $H_2S$  after prolonged exposure to darkness and anoxia combined with an enhancement of OP rates by  $H_2S$  at low irradiance and a temporary resistance to  $H_2S$  toxicity (Klatt et al., 2015b).

The microbial mats in the Frasassi sulfidic springs are characterized by a microenvironment with rapidly fluctuating availability of light and  $H_2S$  (Klatt et al., 2016). Although both *Pseudoanabaena* str. FS39 and *Planktothrix* str. FS34 have adaptations that could make them successful in such environment, microscopic observations revealed that neither of them appeared to be abundant in the system. Instead, the mats were dominated by yet another, morphologically very distinct, cyanobacterium. The aim of this study was to understand the adaptations responsible for the success of this dominant cyanobacterium in the system. We hypothesized that the dominant cyanobacterium possesses a mechanism that



allows it to rapidly switch between AP and OP based on the instantaneous availability of light and  $H_2S$  and thus be adapted to the rapidly fluctuating microenvironmental conditions in the mats. However, we expected that the specific regulation by these parameters differs from that found in *Pseudoanabaena* str. FS39, given that the abundance of this strain in the mats is very low. To test this hypothesis, we quantified the combinatory effects of light and  $H_2S$  on the partitioning between AP and OP performed by the dominant cyanobacterium. Because our cultivation attempts were not successful, we performed our measurements in a natural mat sample dominated by the cyanobacterium.

## MATERIALS AND METHODS

### Microsensors

$O_2$ ,  $H_2S$ , and pH microsensors with a tip diameter of 10–50  $\mu m$  and response time of <2 s were built, calibrated and used for profiling and monitoring of local concentration dynamics as previously described (Revsbech, 1989; Jeroschewski et al., 1996; de Beer et al., 1997; Klatt et al., 2016). A fiber-optic microprobe for scalar irradiance (Lassen et al., 1992) was used to measure the locally available light in the mat. The microprobe was connected to a spectrometer (USB4000, Ocean Optics, USA) and calibrated against a PAR scalar irradiance sensor connected to a light meter (LI-250, both from LI-COR Biosciences, Lincoln, NE, USA) as previously described (Al-Najjar et al., 2010).


### Mat Sampling and Measurement Protocol

Thin cyanobacterial mat together with the underlying sediment were sampled from the Frasassi sulfidic springs (Klatt et al., 2016) in September 2009. Immediately after sampling it was placed in an aquarium and submersed in spring water. The mat was

left overnight in the dark to allow for diffusion of sulfide from the sediment into the water column and re-establishment of an anoxic sulfidic water column resembling *in situ* conditions. The conditions in the mat and water column were monitored by occasional microsensor depth profiling of O<sub>2</sub>, H<sub>2</sub>S, and pH. During measurements, the water column above the mat was covered with paraffin oil. A gentle stream of air onto the paraffin oil surface allowed for a slow, circular flow of the water column in the aquarium.

Net volumetric conversion rates of sulfide and O<sub>2</sub> were calculated from the measured steady-state depth profiles of concentration,  $c(z)$ , using Fick's law of diffusion,  $NP = -D \partial^2 c(z)/\partial z^2$ , where  $D$  is the corresponding molecular diffusion coefficient corrected for temperature and salinity (Gieseke and de Beer, 2004). Specifically, we used  $D = 2.10 \cdot 10^{-9} \text{ m}^2 \text{ s}^{-1}$  and  $D = 1.59 \cdot 10^{-9} \text{ m}^2 \text{ s}^{-1}$  as the diffusion coefficients for O<sub>2</sub> and sulfide, respectively. In our calculations we assumed porosity of 1, which is reasonable for a biofilm comprised primarily of microbial cells. Volumetric rates of gross photosynthesis in the mats were measured using the light–dark shift approach. This approach allows to differentiate between gross and net, and thus between photosynthetic and chemotrophic, sulfide and oxygen conversion rates, based on the dynamics of O<sub>2</sub> and S<sub>tot</sub> in the first few seconds after darkening the sample. The O<sub>2</sub>-based light–dark shift method is well-established and based on the dynamics of O<sub>2</sub> concentration after darkening (Revsbech and Jørgensen, 1983). Analogously, gross AP rates were determined as the rate of sulfide concentration change ( $-dS_{\text{tot}}/dt$ ) upon darkening where the total sulfide concentration ( $S_{\text{tot}} = \text{H}_2\text{S} + \text{HS}^- + \text{S}^{2-}$ ) was calculated from the measured H<sub>2</sub>S concentration and pH. As described previously (Klatt et al., 2016), the variation in pH with time during a light–dark shift measurement was ignored as it was substantially slower and insignificant compared with the variation in H<sub>2</sub>S. During the measurements the tips of the microsensors were separated by <50 μm.

First, the light–dark shift measurements were done at a constant incident irradiance ( $334 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) at various depths of the mat to identify the depth with the highest photosynthetic activity. At this depth gross OP and AP were subsequently determined at incident irradiance levels varying between 37 and  $490 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , adjusted in a random order using a cold halogen light source (KL2500, Schott, Germany). During these measurements H<sub>2</sub>S was gradually depleted in the mat (Figure 2A). Therefore, the mat was occasionally incubated in the dark for up to 1 h to allow sulfide diffuse from deeper layers into the photosynthetically active zone and for the re-establishment of the initial sulfidic conditions at the depth of measurements. Occasionally, complete concentration depth profiles were measured. After the light–dark shift measurements, a depth profile of scalar irradiance was measured in the same spot to determine the locally available light. To exclude a significant contribution of potentially abundant obligate anoxygenic phototrophs to the measured rates of AP, additional measurements were performed under illumination with light emitting diodes (emission maxima at 740 and 810 nm; H2A2 series, Roithner Lasertechnik, Austria) that specifically targeted bacteriochlorophyll in the mat. Finally, a sub-sample



**FIGURE 2 | Examples of transient depth profiles of O<sub>2</sub> and H<sub>2</sub>S concentration (A), and of a depth profile of scalar irradiance (B), measured in the cyanobacterial mat during illumination at incident irradiance of  $169 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . The profiles shown with filled symbols were measured about 2 h after those shown by open symbols. The scalar irradiance values in (B) were integrated over the wavelengths of photosynthetically active radiation (400–700 nm) and normalized to the value at the mat surface. Error bars represent the standard deviation of measurements in three replicate spots of the mat. Note that the filled and open symbols show the same profile in linear and logarithmic scale, respectively. The depth of highest photosynthetic activity (0.83 mm) is indicated by the green area.**


of the measured mat was examined by microscopy using an epifluorescence microscope (Axiophot, Zeiss, Germany).

## Calculations

Total volumetric photosynthetic electron transport rates were calculated as a sum of the estimated electron transport rates due to OP and AP (Klatt et al., 2015a). The former were calculated by multiplying the gross rates of O<sub>2</sub> production measured by the light–dark shift method with a factor of 4, as implied by the reaction  $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$ . Analogously, the latter were calculated by multiplying the gross rates of S<sub>tot</sub> depletion measured by the light–dark shift method with a factor of 2, as implied by the reaction  $\text{H}_2\text{S} \rightarrow \text{S}^0 + 2\text{H}^+ + 2\text{e}^-$ .

## Model of Photosynthetic Electron Transport Chain Reactions

We used a model of the intersecting anoxygenic and oxygenic electron transport chain (Figure 1) to fit our data. The model is described in detail in Klatt et al. (2015a). Briefly, the model focuses on the plastoquinone pool (PQ), the central component of both electron transport pathways. In the PQ pool both pathways intersect (Figure 1). The ratio of the reduction rate of PQ by SQR ( $k_{\text{anoxy}}$  in Figure 1) and the reduction rate of PQ by PSII ( $k_{\text{oxy}}$ ) directly translates into the partitioning between anoxygenic and oxygenic electron transport. The parameters used for fitting the observed partitioning were thus the apparent affinities of SQR and PSII toward PQ ( $K_{\text{SQR}}$  and  $K_{\text{PSII}}$ , respectively), and the affinity of SQR toward H<sub>2</sub>S ( $K_{\text{M}}$ ) and  $v_{\text{max}}$ . Other parameters are not effective for the partitioning



but determine the light dependency of the sum of oxygenic and anoxygenic electron transport rates.

## RESULTS

### Mat Structure and Microscopy


Along the flow path of the sulfidic water emerging from the Frasassi Cave system structurally and functionally diverse microbial mats form (Klatt et al., 2016). In this study, we sampled a mat that was characterized by a green cyanobacterial layer on top of a distinct, very thin layer of large sulfur oxidizing bacteria (SOB). After transfer to the aquarium the water column above the mat was temporary oxic and, when the mat was left in the dark, we observed migration of SOB filaments to the surface of the mat. Exposure to an anoxic and sulfidic water column overnight led to re-establishment of the initial mat structure.

The examined subsample of the cyanobacterial layer was dominated by a single filamentous cyanobacterium with a diameter of  $\sim 4 \mu\text{m}$  (Figure 3). Besides this cyanobacterial morphotype we only observed unicellular non-pigmented prokaryotes (data not shown).

### Microsensor Measurements


#### Oxygenic and Anoxygenic Photosynthesis

Upon illumination with visible light, total sulfide concentration in the cyanobacterial layer immediately decreased and  $\text{O}_2$  concentration increased, suggesting AP and OP, respectively. This was confirmed by detailed measurements of the local volumetric gross rates of AP and OP using the light-dark shift approach (Figure 4). An instantaneous increase of  $\text{H}_2\text{S}$  and decrease of  $\text{O}_2$  upon darkening (e.g., Figure 4C), synonymous with the simultaneously active AP and OP, was detectable until about 1.4 mm depth (Figure 4B). In contrast to visible light, light in the near infrared region of the spectrum, which specifically targets



bacteriochlorophylls of obligate anoxygenic phototrophs, did not induce sulfide consumption at any depth (Figure 5). Thus, all rates of sulfide-driven AP presented here were exclusively assigned to cyanobacteria.

The parallel light-dark shift measurements of  $\text{H}_2\text{S}$  and  $\text{O}_2$  concentration revealed that the gross rates of AP and OP were highest at 0.83 mm depth (Figure 4B). Scalar irradiance at this depth was about 10% of the surface value (Figure 2B). The detailed measurement of gross AP and OP at this depth of highest photosynthetic activity revealed that both types of photosynthesis (anoxygenic and oxygenic) were simultaneously performed over the complete range of light intensities and  $\text{H}_2\text{S}$  concentrations tested (Figure 6). At a given local irradiance, the gross rates of AP increased with increasing  $\text{H}_2\text{S}$  concentration, while the gross rates of OP decreased (Figure 6; open symbols). However,



the total photosynthetic electron transport rate, i.e., the sum of electron transport rates due to OP and AP, did not depend on H<sub>2</sub>S (Figure 6; closed symbols). Light intensity affected the partitioning between the photosynthetic modes, which was most apparent in the light dependency of the initial slopes of OP decrease and AP increase with H<sub>2</sub>S concentration (Figure 6) and the light-dependency of the threshold H<sub>2</sub>S concentration above which AP became dominant over OP (Figure 6). The total electron transport rate increased linearly in the studied range of the locally available light (Figure 6H). In summary, both H<sub>2</sub>S concentration and light intensity regulated the partitioning between OP and AP, while the total electron transport rate was dependent only on light. Importantly, even at very low local light intensities ( $\sim 4 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) and high H<sub>2</sub>S concentrations ( $> 2.4 \text{ mM}$ ) we always observed simultaneous OP and AP and never only AP (Figure 6).

We applied a model of the photosynthetic electron transport chain (Figure 1) to fit our data, assuming that the single morphotype shown in Figure 3 is one strain that is fully responsible for the observed rates. We found an excellent agreement between the measured and predicted rates of OP and AP and their light- and H<sub>2</sub>S-dependent partitioning (compare symbols with lines in Figures 6A–G). To fit the data we must assume that the apparent affinity of SQR to H<sub>2</sub>S is very low ( $K_M > 1 \text{ mM}$ ) and that the apparent affinities of SQR and PSII to oxidized PQ are very similar ( $K_{\text{PSII}}/K_{\text{SQR}} \approx 1$ ).

### Non-photosynthetic Sulfide Consumption

Microsensor depth profiles measured in the dark revealed that there were no significant non-photosynthetic sulfide sinks in the cyanobacterial layer of the mat. In the presence of oxygen in the water column, sulfide and oxygen showed opposing gradients in the uppermost layer (Figure 7A), suggesting chemosynthetic sulfide oxidation. However, the volumetric rates of sulfide consumption were very low and restricted to the top 0.1–0.2 mm, consistent with the slightly whitish appearance of the mat due to the low abundance of chemosynthetic large filamentous SOB. In contrast, O<sub>2</sub> consumption, although low, extended down

to  $\sim 0.8 \text{ mm}$  (Figure 7B), indicating that the main O<sub>2</sub> sink in the mat in darkness was aerobic respiration, e.g., by the cyanobacteria or heterotrophic bacteria. Upon turn of the system to anoxia, there was no detectable sulfide consumption at any depth (Figures 7C,D).


Upon illumination, the net volumetric rates of  $S_{\text{tot}}$  consumption in the cyanobacterial layer closely matched the gross rates of AP (Figure 4B), confirming that non-photosynthetic sulfide consumption was insignificant. Likewise, oxygen consumption rates in the cyanobacterial layer were below detection limit (Figure 4B). Instead, the oxygen produced by OP was consumed in the thin SOB layer underneath the photosynthetically active zone (below 1.4 mm depth; Figure 4B). In this layer also net sulfide consumption occurred (Figure 4B) while we did not observe instantaneous changes in sulfide concentration in response to abrupt changes in illumination (see Oxygenic and Anoxygenic Photosynthesis). This suggests that AP was insignificant while aerobic sulfide oxidation was the dominant sink of both sulfide and oxygen in the SOB layer during illumination.


## DISCUSSION

The cyanobacterial layer in the studied mat performed OP and AP simultaneously over a wide range of H<sub>2</sub>S concentrations and light intensities. The regulation of the total photosynthetic electron transport given by the sum of electron transports driven by AP and OP was astonishingly simple. Specifically, the total electron transport rate increased linearly with irradiance and did not depend on H<sub>2</sub>S at all (Figure 6H). The partitioning between oxygenic and anoxygenic modes seemed more complex as it was controlled by both H<sub>2</sub>S levels and light intensities (Figures 6A–G).

Using our numerical model of the electron transport reactions, we found that the observed activity patterns in this mat can be explained by kinetic controls. Thus, synthesis or degradation of cell components, such as pigments and photosystems, is not needed as an explanation for the complex light- and H<sub>2</sub>S-dependency of photosynthetic rates. We furthermore exclude changes in the abundance of cell components as an explanation for our data, because all our measurements were performed within a short time ( $\sim 6 \text{ h}$ ) and because the observed pattern emerged even though we did not follow a strict order (e.g., either gradually increasing or gradually decreasing) of light intensities and H<sub>2</sub>S concentrations. Also, we repeated the first measurements at the end of our experiments and showed that the physiologic properties of the mat, i.e., the overall photosynthetic rate and the ratio between AP and OP, had not significantly changed. We therefore conclude that the adjustment of rate and rate partitioning in response to the momentary microenvironmental light and H<sub>2</sub>S conditions were instantaneous and the result of kinetic control.

The specific H<sub>2</sub>S and light dependent partitioning between OP and AP can be explained by considering that the light-driven electron transport from PSII and the H<sub>2</sub>S-driven electron transport from SQR intersect in the plastoquinone (PQ) pool






(Figure 1). The electron transport rates are controlled by the affinities governing the rates of PQ reduction by PSII ( $k_{oxy}$  in Figure 1) and SQR ( $k_{anoxy}$ ). We determined that the apparent affinity of SQR to H<sub>2</sub>S has to be very low ( $K_M > 1$ ) (see notation in Figure 1). More importantly, the apparent affinities of SQR and PSII to PQ are in the same range. This means that PSII and SQR can compete equally well for oxidized PQ. As a consequence, OP is performed simultaneously with AP irrespective of the light intensity and H<sub>2</sub>S concentration (Figure 8A), even at H<sub>2</sub>S concentrations that are higher than ever observed *in situ* in the naturally occurring mats (Klatt et al., 2016).

We have observed such kinetic regulation mechanism previously in another cyanobacterium: *Pseudanabaena* str. FS39, a versatile cyanobacterium enriched from the same environment (Klatt et al., 2015a). As the apparent affinities of SQR and PSII toward PQ, however, fundamentally differ

from the cyanobacterium studied here, a completely different H<sub>2</sub>S- and light-dependent activity pattern emerges (Figure 8B). In *Pseudanabaena* str. FS39 the apparent affinity of SQR to H<sub>2</sub>S is very high, with the corresponding  $K_M$  at least two orders of magnitude lower than in the cyanobacterium studied here. Moreover, in *Pseudanabaena* str. FS39 the affinity of SQR to oxidized PQ is at least two orders of magnitude higher than that of PSII ( $K_{PSII}/K_{SQR} \geq 100$ ). In other words, OP is kinetically outcompeted by AP. This is manifested by the fact that OP is only performed in addition to AP when H<sub>2</sub>S is limiting SQR activity. As a consequence, *Pseudanabaena* str. FS39 switches from exclusive AP to simultaneous AP and OP at substantially lower H<sub>2</sub>S concentrations and/or higher light intensities than the cyanobacterium studied here (compare Figures 8A,B).

The striking differences between the affinities of SQR to PQ and H<sub>2</sub>S in *Pseudanabaena* str. FS39 and the dominant spring cyanobacterium suggest that their SQRs belong to different enzyme structure classes (Gregersen et al., 2011). The affinities of PSII to PQ also differ among the two cyanobacteria. Protein D1, which is a core component of the photosystem II reaction center, shapes the kinetics of quinone reduction. Different isoforms of this protein exist and are expressed dependent on oxygen concentration and light (Cardona et al., 2015). We therefore suggest that *Pseudanabaena* str. FS39 and the cyanobacterium studied here might employ fundamentally different types of D1. Thus, there might be a relationship between the type of D1 and regulatory mechanisms for AP, which might have far-reaching implications for research on the evolution of AP in cyanobacteria. The phylogeny of genes encoding for the well-conserved D1 proteins was successfully used to reconstruct their evolutionary history (Cardona et al., 2015). Such reconstruction is challenging when based only on SQR genes due to their history of intensive lateral gene transfer (Theissen et al., 2003).



Genes encoding for several isoforms of D1 have recently been detected in the genome of *Geitlerinema* sp. PCC 9228 (formerly *Oscillatoria limnetica*) (Grim and Dick, 2016), the most intensively studied photosynthetically versatile cyanobacterium, which further underlines the intriguing question if usage of different D1 types might be an important tool for cyanobacteria in the regulation of AP.

It is very unlikely that the observed activity pattern emerged from the simultaneous activity of a mix of several physiologically distinct cyanobacterial species or strains. The studied mats were dominated by a single cyanobacterial morphotype. The complex experimental data could be fitted by a simple kinetic model, thus by one kinetic expression per reaction step. Any residual uncertainty does, however, not invalidate the general conclusion that the most successful cyanobacterial population in the sulfidic spring can perform simultaneous OP and AP over a complete diurnal cycle unless H<sub>2</sub>S becomes entirely depleted within the photosynthetically active zone.

The range of H<sub>2</sub>S concentrations and irradiances over which OP and AP are performed concurrently in the studied mats is substantially wider than in *Pseudanabaena* FS39. In fact, it is even exceptional compared to other previously studied versatile cyanobacterial strains. It was shown that *Geitlerinema* PCC9228, for instance, performs simultaneous OP and AP only below a 80 μM H<sub>2</sub>S concentration threshold when exposed to 400 μmol photons m<sup>-2</sup> s<sup>-1</sup> (Cohen et al., 1986). Cohen et al. (1986) also report that in an *Oscillatoria* sp. from Stinky Hot Springs and in *Microcoleus chthonoplastes* (*Geitlerinema*) from Solar Lake, AP and OP occur concurrently below 550 μM H<sub>2</sub>S and 1000 μM H<sub>2</sub>S, respectively, at the same light intensity. For the case of the Frasassi sulfidic spring cyanobacterium, our model predicts that the OP:AP ratio would still be 0.9 at 1000 μM H<sub>2</sub>S and that exclusive AP is not possible at all at around 400 μmol photons m<sup>-2</sup> s<sup>-1</sup>.

This astonishing coverage of H<sub>2</sub>S concentrations and light intensities inspires an intriguing question: what is the advantage of simultaneous AP and OP in general, and specifically in the Frasassi sulfidic springs? As opposed to obligate anoxygenic phototrophs, both obligate oxygenic phototrophs and versatile cyanobacteria are generally never limited by electron donor availability. There is, however, a major difference between versatile and obligate oxygenic cyanobacteria: AP is driven by only one photosystem (PSI), whereas OP is driven by two photosystems (PSI+PSII). Thus, theoretically, the photon flux required to drive a certain electron transport and growth rate is twice as large for OP as for AP. Due to this lower energy demand of AP, one could expect that AP is favorable for cyanobacteria when H<sub>2</sub>S is not limiting. However, this expected advantage of AP seems not to necessarily hold true in versatile cyanobacteria. For the studied cyanobacterium and also for *Pseudanabaena* str. FS39 we have shown that the total electron transport rate driven by a specific photon flux is independent of H<sub>2</sub>S concentrations and the photosynthetic mode. A constant electron transport implies that the light energy harvested in PSII is basically wasted during AP. This is because only PSI is involved in anoxygenic electron transport and photons harvested in PSII are therefore not efficiently used to drive photochemical reactions.

AP would only be advantageous if the excitation energy were transferred from PSII to PSI or if the photosystem stoichiometry changed (Klatt et al., 2015a). This does not seem to occur in the cyanobacterium dominating the Frasassi sulfidic springs as the sum of photosynthetic electron transport rates is constant. Thus, AP does not appear to provide any energetic advantage to this phototroph.

Considering that quantum efficiency in cyanobacteria can be constant irrespective of the photosynthetic mode, cyanobacteria that perform only one mode of photosynthesis therefore can, from a thermodynamic perspective, theoretically be as successful as versatile cyanobacteria. However, obligate oxygenic phototrophs in the Frasassi sulfidic springs (e.g., *Planktothrix* str. FS34, Klatt et al., 2015b) do not seem to dominate the photosynthetic community. Similarly, cyanobacteria that would perform only AP for most of the diurnal cycle (e.g., *Pseudanabaena* str. FS39, Klatt et al., 2015a, **Figure 8B**) are not abundant. Instead, the key to success appears to lie in the simultaneity of both photosynthetic modes.

Production of oxygen in the presence of sulfide is tied to severe toxification risks by reactive oxygen species (ROS) generation (Latifi et al., 2009; Hoffman et al., 2012). A counterbalancing advantage of such adaptation strategy remains mysterious to us – especially in the modern world where potential competitors for space and resources [e.g., SOB and obligate anoxygenic phototrophs (Klatt et al., 2016)] as well as microorganisms that might affect this competition via feedback mechanisms on the S- and C-cycle (e.g., sulfate reducers) have developed sophisticated strategies to cope with fluctuating oxygen and sulfide concentrations (e.g., Blankenship and Matsuura, 2003; Baumgartner et al., 2006; Berghoff et al., 2011). We can therefore only speculate that the advantage of versatility might be understood by considering that cyanobacteria are not an independently operating entity but almost always rely on the interaction with other microbes, which is most apparent in the frequent failure of axenic cultivation of cyanobacteria. In this scenario a close beneficial interaction with aerobes and/or oxidizers of the products of cyanobacterial AP, i.e., intermediately reduced sulfur compounds (Cohen et al., 1975; de Wit and van Gemerden, 1987; Rabenstein et al., 1995), is plausible and represents a hypothesis that warrants testing.

While the advantage of continuous O<sub>2</sub> production and sulfide consumption at virtually any light intensity and H<sub>2</sub>S concentration in contemporary ecosystems remains to be understood, the capability to perform OP whenever there is a glimpse of light might have provided the ancestors of extant cyanobacteria with a crucial advantage in the microbial mats that covered Earth's costal oceans for much of the Proterozoic and likely already during the Archean. These mats represent the stage for the evolution and proliferation of OP (Buick, 2008; Lyons et al., 2014), a process which must have been a catastrophe for microbial life in an anaerobic world but also triggered the co-evolution of aerobic life. Thus, even the earliest aerobic life-style must have been able to thrive in the fluctuating redox conditions of microbial mats. Overall, our study emphasizes that the evolution of cyanobacterial adaptations strategies and



the resulting cyanobacterial activity, i.e., both the gross and net O<sub>2</sub> production rates in microbial mats, were tightly coupled to the evolution of Earth's redox environment (that is, the availability of alternative electron donors for photosynthesis), the energy supply in the form of light and chemical energy and likely the co-evolution of competitive and beneficial microbial interactions.

## AUTHOR CONTRIBUTIONS

JK, DB, SH, and LP made substantial contributions to the conception and the design of the study, the acquisition of data, and the analysis and interpretation of data; JK, DB, SH, and LP participated in drafting the article and revising it critically for important intellectual content; JK, DB, SH, and LP gave final approval of the version to be submitted. JK, DB, SH, and LP agreed to be accountable for all aspects of the work in ensuring

that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

## FUNDING

This work was financially supported by the Max-Planck-Society.

## ACKNOWLEDGMENTS

We thank the technicians of the microsensor group for the construction of microsensors, J. Macalady and D. S. Jones for their support during field campaigns and A. Montanari and P. Metallo for their hospitality during the sampling campaign at the Osservatorio Geologico di Coldigioco.

## REFERENCES

- Al-Najjar, M. A. A., de Beer, D., Jørgensen, B. B., Kühl, M., and Polerecky, L. (2010). Conversion and conservation of light energy in a photosynthetic microbial mat ecosystem. *ISME J.* 4, 440–449. doi: 10.1038/ismej.2009.121
- Baumgartner, L. K., Reid, R. P., Dupraz, C., Decho, A. W., Buckley, D. H., Spear, J. R., et al. (2006). Sulfate reducing bacteria in microbial mats: changing paradigms, new discoveries. *Sediment. Geol.* 185, 131–145. doi: 10.1016/j.sedgeo.2005.12.008
- Berghoff, B. A., Glaeser, J., Nuss, A. M., Zobawa, M., Lottspeich, F., and Klug, G. (2011). Anoxygenic photosynthesis and photooxidative stress: a particular challenge for *Roseobacter*. *Environ. Microbiol.* 13, 775–791. doi: 10.1111/j.1462-2920.2010.02381.x
- Blankenship, R. E. (2010). Early evolution of photosynthesis. *Plant Physiol.* 154, 434–438. doi: 10.1104/pp.110.161687
- Blankenship, R. E., and Matsuura, K. (2003). "Green photosynthetic bacteria," in *Light-Harvesting Antennas in Photosynthesis*, eds B. Green and W. W. Parson (Dordrecht: Springer Science & Business Media), 195–217.
- Buick, R. (2008). When did oxygenic photosynthesis evolve? *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 2731–2743. doi: 10.1098/rstb.2008.0041
- Canfield, D. E. (1998). A new model for Proterozoic ocean chemistry. *Nature* 396, 450–453. doi: 10.1038/24839
- Cardona, T., Murray, J. W., and Rutherford, A. W. (2015). Origin and evolution of water oxidation before the last common ancestor of the cyanobacteria. *Mol. Biol. Evol.* 32, 1310–1328. doi: 10.1093/molbev/msv024
- Cohen, Y., Jørgensen, B. B., Revsbech, N. P., and Poplawski, R. (1986). Adaptation to hydrogen sulfide of oxygenic and anoxygenic photosynthesis among cyanobacteria. *Appl. Environ. Microbiol.* 51, 398–407.
- Cohen, Y., Padan, E., and Shilo, M. (1975). Facultative anoxygenic photosynthesis in the cyanobacterium *Oscillatoria limnetica*. *J. Bacteriol.* 123, 855–861.
- de Beer, D., Schramm, A., Santegoeds, C. M., and Kühl, M. (1997). A nitrite microsensor for profiling environmental biofilms. *Appl. Environ. Microbiol.* 63, 973–977.
- de Wit, R., and van Gemerden, H. (1987). Oxidation of sulfide to thiosulfate by *Microcoleus* chthonoplastes. *FEMS Microbiol. Lett.* 45, 7–13. doi: 10.1111/j.1574-6968.1987.tb02332.x
- Gieseke, A., and de Beer, D. (2004). "Use of microelectrodes to measure in situ microbial activities in biofilms, sediments, and microbial mats," in *Molecular Microbial Ecology Manual*, eds A. Akkermans and D. van Elsas (Dordrecht: Springer), 1581–1612. doi: 10.1007/978-1-4020-2177-0\_802
- Gregersen, L. H., Bryant, D. A., and Frigaard, N.-U. (2011). Mechanisms and evolution of oxidative sulfur metabolism in green sulfur bacteria. *Front. Microbiol.* 2:116. doi: 10.3389/fmicb.2011.00116
- Grim, S., and Dick, G. (2016). Photosynthetic versatility in the genome of *Geitlerinema* sp. PCC 9228 (formerly *Oscillatoria* limnetica "Solar Lake"), a model anoxygenic photosynthetic cyanobacterium. *Front. Microbiol.* 7:1546. doi: 10.3389/fmicb.2016.01546
- Hoffman, M., Rajapakse, A., Shen, X., and Gates, K. S. (2012). Generation of DNA-damaging reactive oxygen species via the autoxidation of hydrogen sulfide under physiologically relevant conditions: chemistry relevant to both the genotoxic and cell signaling properties of H<sub>2</sub>S. *Chem. Res. Toxicol.* 25, 1609–1615. doi: 10.1021/tx300066z
- Jeroschewski, P., Steuckart, C., and Kühl, M. (1996). An amperometric microsensor for the determination of H<sub>2</sub>S in aquatic environments. *Anal. Chem.* 68, 4351–4357. doi: 10.1021/ac960091b
- Johnston, D. T., Wolfe-Simon, F., Pearson, A., and Knoll, A. H. (2009). Anoxygenic photosynthesis modulated Proterozoic oxygen and sustained Earth's middle age. *Proc. Natl. Acad. Sci. U.S.A.* 106, 16925–16929. doi: 10.1073/pnas.0909248106
- Klatt, J. M., Al-Najjar, M. A. A., Yilmaz, P., Lavik, G., de Beer, D., and Polerecky, L. (2015a). Anoxygenic photosynthesis controls oxygenic photosynthesis in a cyanobacterium from a sulfidic spring. *Appl. Environ. Microbiol.* 81, 2025–2031. doi: 10.1128/AEM.03579-14
- Klatt, J. M., Haas, S., Yilmaz, P., de Beer, D., and Polerecky, L. (2015b). Hydrogen sulfide can inhibit and enhance oxygenic photosynthesis in a cyanobacterium from sulphidic springs. *Environ. Microbiol.* 17, 3301–3313. doi: 10.1111/1462-2920.12791
- Klatt, J. M., Meyer, S., Häusler, S., Macalady, J. L., de Beer, D., and Polerecky, L. (2016). Structure and function of natural sulphide-oxidizing microbial mats under dynamic input of light and chemical energy. *ISME J.* 10, 921–933. doi: 10.1038/ismej.2015.167
- Lassen, C., Ploug, H., and Jørgensen, B. B. (1992). A fibre-optic scalar irradiance microsensor: application for spectral light measurements in sediments. *FEMS Microbiol. Lett.* 86, 247–254. doi: 10.1111/j.1574-6968.1992.tb04816.x
- Latifi, A., Ruiz, M., and Zhang, C.-C. (2009). Oxidative stress in cyanobacteria. *FEMS Microbiol. Rev.* 33, 258–278. doi: 10.1111/j.1574-6976.2008.00134.x
- Lyons, T. W., Reinhard, C. T., and Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and atmosphere. *Nature* 506, 307–315. doi: 10.1038/nature13068
- Miller, S. R., and Bebout, B. M. (2004). Variation in sulfide tolerance of photosystem II in phylogenetically diverse cyanobacteria from sulfidic habitats. *Appl. Environ. Microbiol.* 70, 736–744. doi: 10.1128/AEM.70.2.736-744.2004
- Mulkidjanian, A. Y., Koonin, E. V., Makarova, K. S., Mekhedov, S. L., Sorokin, A., Wolf, Y. I., et al. (2006). The cyanobacterial genome core and the origin of photosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* 103, 13126–13131. doi: 10.1073/pnas.0605709103
- Nisbet, E. G., and Fowler, C. M. R. (1999). Archaean metabolic evolution of microbial mats. *Proc. R. Soc. B Biol. Sci.* 266, 2375–2382. doi: 10.1098/rspb.1999.0934

- Rabenstein, A., Rethmeier, J., and Fischer, U. (1995). Sulphite as intermediate sulphur compound in anaerobic sulphide oxidation to thiosulphate by marine cyanobacteria. *Z. Naturforsch.* 50, 769–774.
- Revsbech, N. P. (1989). An oxygen microsensor with a guard cathode. *Limnol. Oceanogr.* 34, 474–478. doi: 10.4319/lo.1989.34.2.0474
- Revsbech, N. P., and Jørgensen, B. B. (1983). Photosynthesis of benthic microflora measured with high spatial resolution by the oxygen microprofile method: capabilities and limitations of the method. *Limnol. Oceanogr.* 28, 749–759. doi: 10.4319/lo.1983.28.4.0749
- Theissen, U., Hoffmeister, M., Grieshaber, M., and Martin, W. (2003). Single eubacterial origin of eukaryotic sulfide:quinone oxidoreductase, a mitochondrial enzyme conserved from the early evolution of eukaryotes during anoxic and sulfidic times. *Mol. Biol. Evol.* 20, 1564–1574. doi: 10.1093/molbev/msg174
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Klatt, de Beer, Häusler and Polerecky. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.