



Diversity of the Bacterial Microbiota of *Anopheles* Mosquitoes from Binh Phuoc Province, Vietnam

Chung T. Ngo^{1,2}, Sara Romano-Bertrand^{3,4*}, Sylvie Manguin¹ and Estelle Jumas-Bilak^{3,4}

¹ Institut de Recherche pour le Développement France, UMR-MD3, Faculté de Pharmacie, Montpellier, France, ² National Institute of Veterinary Research, Hanoi, Vietnam, ³ UMR 5569 Hydrosociétés, Equipe Pathogènes Hydriques, Santé et Environnements, Faculté de Pharmacie, Université de Montpellier, Montpellier, France, ⁴ Département d'Hygiène Hospitalière, Centre Hospitalier Universitaire de Montpellier, Montpellier, France

The naturally acquired microbiota of *Anopheles* can influence vector's susceptibility to *Plasmodium* and its capacity to transmit them. Microbiota modification is a new challenge to limit disease transmission but it still needs advanced knowledges on bacterial community in *Anopheles*, especially in wild and infected specimens from diverse origin and species. Bacterial culture and 16S rRNA gene-PCR associated to Temporal Temperature Gradient Electrophoresis (TTGE) were applied to explore the bacterial diversity in the abdomen of 100 wild specimens (eight *Anopheles* species) collected in the Binh Phuoc Province, Vietnam. Culture and PCR-TTGE were complementary. The bacterial richness of the mosquito collection encompassed 105 genera belonging to seven phyla, mostly *Proteobacteria*, *Firmicutes*, and *Actinobacteria*. *Staphylococcus*, *Clostridium*, and *Bacillus* in *Firmicutes* were the most prevalent genera. However, *Proteobacteria* represented by 57 genera was the most diversified phylum in *Anopheles* microbiota. The high overall of *Anopheles*-associated bacteria is confirmed with, to our knowledge, 51 genera described for the first time in *Anopheles* microbiota. However, the diversity per specimen was low with average diversity index and the average Shannon–Wiener score (H) of 4.843 and 5.569, respectively. The most represented bacterial genera were present in <30% of the specimens. Consequently, the core microbiota share by *Anopheles* from Binh Phuoc was very narrow, suggesting that *Anopheles* microbiota was greatly influenced by local environments. The repertory of bacterial genera in two specimens of *An. dirus* and *An. pampanai* naturally infected by *Plasmodium vivax* was also described as preliminary results. Finally, this study completed the repertory of bacteria associated to wild *Anopheles*. *Anopheles* associated-bacteria appeared specimen-dependent rather than mosquito species- or group-dependent. Their origin and the existence of *Anopheles*-specific bacterial taxa are discussed.

Keywords: *Anopheles* mosquitoes, abdomen bacterial microbiota, 16S rRNA PCR – TTGE, malaria, Binh Phuoc, Vietnam

OPEN ACCESS

Edited by:

Miguel Cacho Teixeira,
University of Lisbon, Portugal

Reviewed by:

Guy Smagghe,
Ghent University, Belgium
George Christophides,
Imperial College London, UK

*Correspondence:

Sara Romano-Bertrand
sara.romano-bertrand@umontpellier.fr

Specialty section:

This article was submitted to
Infectious Diseases,
a section of the journal
Frontiers in Microbiology

Received: 22 April 2016

Accepted: 12 December 2016

Published: 23 December 2016

Citation:

Ngo CT, Romano-Bertrand S,
Manguin S and Jumas-Bilak E (2016)
Diversity of the Bacterial Microbiota
of *Anopheles* Mosquitoes from Binh
Phuoc Province, Vietnam.
Front. Microbiol. 7:2095.
doi: 10.3389/fmicb.2016.02095

INTRODUCTION

Despite some success in controlling malaria, this disease continues to be a major health burden in many countries around the world with 438,000 deaths reported in 2015, particularly in sub-Saharan Africa recording 395,000 deaths (90%), but also in forested regions of Southeast Asia with 32,000 deaths (7%; WHO, 2015). Until now, vector control has been one of the key elements for controlling this disease based on the use of insecticides, such as DDT, pyrethroids, organophosphates, carbamates (Verhaeghen et al., 2009; Kabula et al., 2013). However, this strategy of insecticide use has many serious side-effects, in particular on human health and the environment (Dabrowski and Balderacchi, 2013; Singleton et al., 2013; Dabrowski et al., 2014), having also a direct impact on the increase of *Anopheles* resistance to insecticides which has now been reported in 64 countries (WHO, 2012; Chang et al., 2013; Kabula et al., 2013; Tangena et al., 2013). Therefore, new and innovative approaches to control *Anopheles* vectors for reducing malaria transmission on a more durable and safer manner for human health are, hence, needed.

In Vietnam, malaria remains the most important vector-borne parasitic disease with a strong prevalence in forested regions, in particular along the international borders with Cambodia, where Binh Phuoc Province is located. This province, situated in central-south Vietnam, is considered as having the highest malarial transmission of all (Abe et al., 2009; Tran et al., 2012). The local malaria situation reported both resistance of *Plasmodium* to anti-plasmodial drugs and *Anopheles* mosquitoes to the insecticide used in this province (Huong et al., 2001; Nguyen et al., 2003; Thanh et al., 2010; Verhaeghen et al., 2010; Tran et al., 2012). However, there is a lack of information on the distribution and vector competence of the local *Anopheles* vectors.

The impact of the microbiota on mosquito infection and more specifically on *Anopheles* resistance to malaria pathogens is showing great potential toward reducing the mosquito vector competence and blocking transmission of several infectious diseases (Dong et al., 2006, 2009, 2011; Blumberg et al., 2013). The influence of the *Anopheles* microbiota on its vectorial competence to transmit pathogens is a growing field of investigation as demonstrated by the recent increase in studies, to cite a few (Favia et al., 2007; Briones et al., 2008; Dong et al., 2009; Rodrigues et al., 2010; Cirimotich et al., 2011; Blumberg et al., 2013; Gendrin and Christophides, 2013; Dennison et al., 2014; Buck et al., 2016). A previous report on the biodiversity of the abdomen microbiota of 100 wild *Anopheles* specimens (five species) from Dak Nong Province showed a high taxonomic diversity, including species reported as implicated in the mosquito resistance to *Plasmodium* infection (Ngo et al., 2015). By the same approach, we present here the bacterial diversity detected in 100 *Anopheles* specimens of eight species collected from Binh Phuoc, a Province characterized by high malaria endemicity. The *Anopheles* studied herein included two specimens naturally infected by *Plasmodium vivax*.

MATERIALS AND METHODS

Anopheles Samples: Collection and Identification

One hundred *Anopheles* (out of 486 specimens collected from 4 sites located in Bu Gia Map District, Binh Phuoc Province in central-southern Vietnam (Ngo et al., 2014), which belonged to eight *Anopheles* species, including *Anopheles dirus*, *An. jeyporiensis*, *An. maculatus*, *An. minimus*, *An. pampanai*, *An. rampae*, *An. sawadwongporni*, and *An. scanloni*, were randomly chosen for the present study. These *Anopheles* specimens were collected between November and December, 2011 (rainy-beginning of the dry season) during 11 consecutive nights (17:00–20:00 h using methods, such as light trap capture and human landing catches indoor and outdoor (Ngo et al., 2014).

Anopheles mosquito identification was morphologically done in the field by sorting out each taxon. Specimens that belonged to the Dirus Complex or the Funestus and Maculatus Groups were individually identified to species level using the appropriate PCR-based assay as described by Walton et al. (1999, 2007) and Garros et al. (2005). Each individual was split into two pieces stored at -80°C until use: (1) head-thorax for species identification and detection of *Plasmodium* by molecular methods (Ngo et al., 2014); and (2) abdomen for bacterial analysis. One hundred abdomens of wild-caught *Anopheles* females were used for the subsequent bacterial study by both methods, culture and DNA fingerprint, as described below.

Bacterial Culture, DNA Extraction, and Strains Identification

Anopheles abdomens were surface rinsed twice in sterilized DNA-free water, and each abdomen was thoroughly disrupted using a sterilized tissue crusher device in 150 μl of sterile DNA-free water. Then, 10 μl of this suspension was spread on each prepared culture medium plate: blood sheep agar, R2A and *Acetobacter* agar. The detailed protocol followed the one published by Ngo et al. (2015). In order to identify strains, 16S rRNA gene was amplified using the Taq DNA polymerase (Go Taq Promega) and the universal primer pair (27f and 1492r) as previously described (Romano-Bertrand et al., 2012). PCR amplifications were checked by DNA electrophoresis in 1.5% agarose gels containing ethidium bromide and visualized under ultraviolet light. Sterile DNA-free water used for mosquito preparations was used in negative control for each series of DNA extraction and PCR. The successfully amplified products were sequenced with primer 27f, on an ABI 3730xl sequencer (Cogenics, Meylan, France). Each sequencing chromatograph was visually inspected and corrected.

Independent-Culture Analysis by 16S PCR-Temporal Temperature Gel Electrophoresis (PCR-TTGE)

Whole DNA was extracted from 100 μl of mosquito abdomen suspension using the Master Pure Gram Positive DNA

purification kit as recommended by the supplier (Epicentre Biotechnologies, Madison, WI, USA). The purified and raw DNAs were kept at -80°C before further analyses. Because of low bacterial load in mosquito's samples, a nested-PCR approach was necessary to obtain a sufficient PCR amplification (Romano-Bertrand et al., 2012). The first PCR amplified almost entirely the 16S rRNA gene using the 27f and 1492r primers as previously described (Romano-Bertrand et al., 2012). Then, products of the first PCR were used as template for the amplification of the V2–V3 hyper variable region using the FastStart high fidelity PCR system (Roche apply science) and the primers HDA1 with GC-clamp and HDA2 (Ogier et al., 2002; Roudiere et al., 2009). Again, sterile DNA-free water was used as negative control for each series of nested-PCR. If positive signal was observed, controls were analyzed in parallel of samples by TTGE migration. TTGE profiles of samples and controls were then compared. Positive controls, consisting on decreasing inoculums of *Elizabethkingia anophelis* (which is especially retrieved from mosquitos samples), were used to validate the analysis, from the DNA extraction and throughout the whole Temporal Temperature Gel Electrophoresis (TTGE) process. TTGE migration and TTGE bands sequencing were performed as described in Ngo et al. (2015).

Sequence Analysis, Species Identification, and OTU Affiliation

The sequences were analyzed by comparison with Genbank¹ and Ribosomal Databases Project 2 (RDPII)² using Basic Local Alignment Search Tool (BLAST) and Seqmatch programs, respectively. The sequence with the highest percentage was used for OTU affiliation as previously described by Ngo et al. (2015). Briefly, a sequence was affiliated to a species-level OTU when the percent of sequence similarity with the species type strain was above 99.0% (Drancourt et al., 2000). Below 99.0%, the sequence was affiliated to the genus of the reference sequence with the highest percentage. When several species reference sequences match equally, affiliation was done to the genus level or to a group of species if relevant. For example, sequence with 99.5% in similarity to both *Aeromonas caviae* and *A. hydrophila* was only assigned to the genus *Aeromonas*. The same rule was applied for the taxonomic level higher than the genus level if necessary. The 16S rRNA gene sequences from all cultured bacteria are available in GenBank (accession numbers KX449232–KX449311). Sequences from TTGE band are available by contacting corresponding author or by consulting the Supplementary Fasta File.

Diversity Index Calculation and Statistical Analysis

The microbiota diversity for each species of *Anopheles* was estimated by the calculation of crude diversity index (DI, number of different OTUs; Romano-Bertrand et al., 2012, 2014), Shannon–Wiener DI (H) and Simpson's index (D; Gafan et al., 2005; Ledder et al., 2007; Romano-Bertrand et al., 2014).

The 100 *Anopheles* specimens were classified into the Dirus Complex or into the Funestus and Maculatus Groups according to their species identification. The bacterial diversity scores (DI, H, and D) for the different *Anopheles* species, complex and groups were then calculated.

RESULTS

Anopheles Identification

Specific PCR assays identified eight *Anopheles* species among the 100 specimens collected in Binh Phuoc Province. Out of 100, 37 specimens belonged to the Dirus Complex including *An. dirus* ($n = 33$) and *An. scanloni* ($n = 4$), 43 to the Funestus Group with *An. minimus* ($n = 34$), *An. jeyporiensis* ($n = 5$), and *An. pampanai* ($n = 4$), and 20 to the Maculatus Group with *An. sawadwongporni* ($n = 12$), *An. rampae* ($n = 6$), and *An. maculatus* ($n = 2$; Figure 1A).

Bacterial Diversity in the Abdomen of *Anopheles* Adults

The abdomens of 100 *Anopheles* were analyzed using both bacterial culture and PCR-TTGE methods. Only a few negative controls presented weak amplification bands in agarose gel after nested-PCR but these bands were never observed after TTGE migration, permitting to validate the profiles for mosquitoes' samples. The regular positive controls highlighted a sensitivity of detection comprised between 10^1 and 10^2 bacterial inoculums throughout the TTGE process (data not shown). Bacteria were cultured or detected in 98% of the *Anopheles* specimens, the two negative specimens belonged to species *An. minimus* and *An. scanloni*. A total of 63 bacterial strains from 22 specimens positive by culture and 441 TTGE bands from 98 specimens positive by PCR were obtained. Distributions of strains and TTGE bands according to *Anopheles* species are shown in Figures 1B,C, respectively.

Both strains and 393 TTGE bands were further analyzed by 16S rRNA gene PCR and sequenced. The remaining 48 TTGE bands were identified by comparison with another co-migrating band on the same TTGE gel. Results were presented in Supplementary Tables. Strains and PCR-TTGE bands were affiliated to seven phyla, the most represented being *Proteobacteria*, *Firmicutes*, and *Actinobacteria* (Figure 2). Considering OTUs, 193 bacterial OTUs corresponding to 190 genera-level OTUs (including 85 OTUs for which identification was performed until species-level), one family-level and two phyla-level OTUs. The sequences obtained from 3 TTGE bands were not accurate enough to be affiliated to the genus level, probably due to mixed DNA amplified and sequenced from co-migrating bands; one of them was affiliated to the Phylum *Acidobacteria* and two others to the Family *Planctomycetaceae* (Phylum *Planctomycetes*). The OTU richness by phylum is summarized in Figure 3. *Proteobacteria* was the most diversified phylum with a richness corresponding to 57 genera-level OTUs, followed by *Actinobacteria* and *Firmicutes* with, respectively, 24 and 18 genera-level OTUs.

¹<http://www.ncbi.nlm.nih.gov/>

²<http://rdp.cme.msu.edu/>




FIGURE 1 | Distribution according to *Anopheles* species, complex and group of (A) mosquitoes specimens, (B) bacterial strains, and (C) TTGE bands.




FIGURE 2 | Distribution of TTGE bands and bacterial isolates according to phylum.

Only 32 from the 190 genera-level OTUs were retrieved in culture, mainly from *Actinobacteria* with 18 cultivable genera (56%). Out of these 32 cultivable OTUs, 15 (7.9%) were not retrieved by the molecular approach (genera underlined in Figure 3 and Supplementary Tables).

In this study, molecular approach allowed the detection of more bacteria than culture, although culture improved the description of bacterial richness in the *Anopheles* abdomen microbiota and should be considered as complementary to PCR-based methods. The *Actinobacteria* richness was particularly wide thanks to culture.

Bacterial Richness and Diversity of the Abdomen Microbiota According to the *Anopheles* Species

Bacterial diversity according to *Anopheles* species was displayed in Figure 4A. No obvious link was observed between *Anopheles* complex or groups and microbiota diversity. However, the number of specimens in several *Anopheles* species is too low for robust conclusions, only tendencies can be drawn. Amongst the three major phyla, *Firmicutes* and *Proteobacteria* were present in all *Anopheles* species, while *Actinobacteria* were absent from




FIGURE 3 | Repertory of genus-level OTUs according to phylum in adult *Anopheles* gut microbiota. *Cultivable genera; Genera only retrieved in culture are underlined; ¹Genera identified in *Anopheles* abdomen for the first time; Genera previously found in *Anopheles* from Vietnam in bold face (Ngo et al., 2015); In parenthesis, the percent (50% and higher) of *Anopheles* species carrying genera frequently found in Binh Phuoc *Anopheles* microbiota.

An. jeyporiensis, *An. scanloni*, and *An. rampae* (Figure 4). *An. minimus*, considered as one of the primary vector of *Plasmodium* in Southeast Asia, including Vietnam, was the sole species colonized by members of the seven bacterial phyla. Two other malaria vectors well represented in the collection, *An. dirus* and *An. sawadwongporni*, were each colonized by five bacterial phyla (Figure 4A). Despite the low number of specimens ($n = 6$) in this species, *An. rampae* presented a particular diversity profile with the replacement of *Actinobacteria* by *Bacteroidetes* regarding other *Anopheles* species. Nevertheless, the phylum richness appeared more dependent to the number of specimens in a species of *Anopheles* than to the *Anopheles* complex and groups (Figures 4A,B). Indeed, if *Anopheles* complex or group were considered, the phyla distribution in percent was very similar among them (Figure 4B).

However, at the OTU level the distribution varied according *Anopheles* complex and groups (Figure 4C). Among the 193 OTUs, 113, 109, and 57 OTUs were identified within the Dirus Complex, the Funestus and Maculatus Groups, respectively (Figure 4C). Out of them, 24 OTUs were shared by the three *Anopheles* taxa, 20 OTUs were shared by the Dirus Complex and the Funestus Group, and nine OTUs

were commonly identified either between the Funestus and Maculatus Groups, or the Dirus Complex and Funestus Group (Figure 4C).

Anopheles abdomen microbiota diversity quantified by the mean index and scores was presented in Table 1. The average DI per specimen reached 4.843. *An. maculatus* (DI = 7.0) and *An. jeyporiensis* (DI = 6.0) microbiota were the most diverse, whereas *An. minimus* displayed lowest diversity (DI = 4.429) (Table 1). Considering both bacterial richness and equitability of the OTUs within *Anopheles* species, the average Shannon–Wiener score (H) for all species was 5.569. *An. minimus* had the higher H score (H = 6.213) indicating that the numbers of OTUs are evenly distributed between all specimens (Table 1). The Simpson index (D) takes into account both bacterial species richness and an evenness of abundance among the bacterial species present, and is inversely proportional to the diversity (DI). For all *Anopheles* species, except *An. sawadwongporni* ($D = 0.308$), the D-values were low ranging between 0 and 0.082, suggesting high diversities (Table 1). However, these results should be taken with caution because of the low number of specimens for *An. scanloni* ($n = 3$), *An. jeyporiensis* ($n = 5$), *An. pampanai* ($n = 4$), *An. maculatus* ($n = 2$), and *An. rampae* ($n = 6$).




TABLE 1 | Number of OTUs and diversity indexes for each species of *Anopheles*.

Complex or group	Species	Number of different OTUs	DI (mean ± SD)	H (mean ± SD)	D (mean ± SD)
Dirus complex	<i>An. dirus</i>	110	5.176 (± 2.276)	4.446 (± 0.024)	0.011 (± 0.001)
	<i>An. scanloni</i>	13	4.667 (± 2.517)	2.656 (± 0.025)	0.013 (± 0.004)
	All	115	5.135 (± 2.263)	4.585 (± 0.022)	0.009 (± 0.000)
Funestus group	<i>An. minimus</i>	94	4.429 (± 1.852)	6.213 (± 0.039)	0.042 (± 0.005)
	<i>An. pampanai</i>	19	6.000 (± 2.708)	3.327 (± 0.042)	0.035 (± 0.004)
	<i>An. jeyporiensis</i>	19	5.000 (± 2.449)	3.327 (± 0.042)	0.035 (± 0.004)
	All	110	4.636 (± 2.001)	6.488 (± 0.028)	0.022 (± 0.001)
Maculatus group	<i>An. maculatus</i>	14	7.000 (± 1.414)	2.639 (± 0.000)	0 (± 0)
	<i>An. sawadwongporni</i>	41	4.462 (± 2.145)	4.552 (± 0.042)	0.308 (± 0.018)
	<i>An. rampae</i>	19	4.467 (± 1.506)	3.554 (± 0.058)	0.082 (± 0.009)
	All	60	4.762 (± 1.998)	4.798 (± 0.547)	0.012 (± 0.000)
	All	193	4.843 (± 2.098)	5.569 (± 0.011)	0.003 (± 0.000)

The Core Microbiota of Vietnam *Anopheles* Is Limited

Three *Firmicutes* genera, *Staphylococcus*, *Bacillus*, and *Clostridium*, were the most prevalent in the abdomen microbiota of Binh Phuoc *Anopheles*, respectively, identified in 29.6, 23.5, and 21.4% of the specimens. Besides, *Propionibacterium*, *Peptostreptococcus*, *Acinetobacter*, *Burkholderia*, *Comamonas*, *Delftia*, *Lysobacter*, and *Sphingomonas* were detected in more than 10% of the specimens, with frequencies ranging from 10.2 to

18.4%. The Family *Enterobacteriaceae* (*Enterobacter*, *Klebsiella*, *Pantoea*, and *Proteus*) was detected in 15.3% of the specimens. The other 90 genera were found at frequencies lower than 10%, including 81 genera detected in only 1–4 samples (<5%), certainly due to the important bacterial diversity in the mosquito microbiota (Supplementary Table 1).

The most represented genera in Binh Phuoc *Anopheles* microbiota were *Staphylococcus* and *Sphingomonas* (Supplementary Table 1 and Figure 3). Six other genera

retrieved at lower frequencies were newly identified in our study: *Agrococcus*, *Anoxybacillus*, *Peptostreptococcus*, *Lysobacter*, *Paracoccus*, and *Sphingopyxis* (Supplementary Table 1 and **Figure 3**). Finally, a total of 21 genera were detected in more than 4 of 8 *Anopheles* species but these genera were present in <30% of the mosquito specimens confirming that the high bacterial diversity in Binh Phuoc *Anopheles* impaired the definition a core microbiota. Nevertheless, the presence of some frequent genera in Binh Phuoc *Anopheles* microbiota that were not yet described in *Anopheles* pan-microbiota, suggested the existence of specific microbiota according to the geographic region probably due to a strong influence of local environmental conditions.

Concerning the pan-microbiota of the whole genus *Anopheles*, meta-analysis of recent studies (Gendrin and Christophides, 2013; Manguin et al., 2013; Minard et al., 2013; Sharma et al., 2014; Villegas and Pimenta, 2014; Ngo et al., 2015) and bibliography databases survey for older publications, indicated that 51 bacterial genera (48.6%) were probably detected herein for the first time in *Anopheles* (1 in **Figure 3** and Supplementary Table 1). These newly described genera could be slightly overestimated due to the meta-analysis approach and the comparison with non-fully exhaustive data but it clearly suggested that the overall richness of *Anopheles* microbiome is far from being fully explored. This is particularly true in Southeast Asia where very little work has been done on this topic (Gendrin and Christophides, 2013; Ngo et al., 2015).

In comparison with a previous study on the microbiota diversity in specimens from Dak Nong Province, Vietnam (Ngo et al., 2015), 17 genera (in bold in **Figure 3**) were common in *Anopheles* species from both provinces. Most of these genera were found at frequencies below 10% in the *Anopheles* specimens from Binh Phuoc Province, except four genera: *Sphingomonas* (13.3%), *Acinetobacter* (18.4%), *Bacillus* (23.5%), and *Staphylococcus* (29.6%). The genus *Acinetobacter* was detected in more than half *Anopheles* species in both Binh Phuoc and Dak Nong *Anopheles* microbiota. It could be considered as the sole genus forming the core microbiota of Vietnam *Anopheles*.

Characterization of the Abdomen Microbiota of *Plasmodium* Naturally Infected *Anopheles* Specimens

Two specimens were naturally infected by *P. vivax*, out of the 486 *Anopheles* specimens collected in Binh Phuoc Province (Ngo et al., 2014). They belonged to the species *An. dirus* and *An. pampanai*. The abdomen bacterial microbiota of these two specimens was described by PCR-TTGE only, because cultured isolates were not obtained from these two infected mosquitoes. The sequence analysis of seven TTGE bands showed the presence of six bacterial taxa: *Acinetobacter* sp., *Geobacillus* sp., *Luteimonas* sp., *Methylobacterium* sp., *Rubellimicrobium* sp., and *Sphingomonas* sp., among which five (83.3%) belonged to *Proteobacteria* and only one, *Geobacillus* sp., to *Firmicutes*. The genus *Acinetobacter*, *Geobacillus*, *Methylobacter*, and *Sphingomonas* belonged to the core^{50%} microbiota and were also retrieved from other non-infected specimens of *Anopheles*, whereas *Luteimonas* was only retrieved from the infected

An. pampanai specimen, and *Rubellimicrobium* from both the infected *An. dirus* and one specimen of *An. rampae* (Supplementary Table 1).

DISCUSSION

The description of microbiota in vector arthropods presented increasing interest because of potential role in vector competence. Protective role of *Anopheles* bacterial microbiota against *Plasmodium* infections has been demonstrated because clearing midgut microbiota by antibiotic treatment resulted in enhanced *Plasmodium* infections rate (Beier et al., 1994; Favia et al., 2007; Dong et al., 2009; Rodrigues et al., 2010). Similarly but in semi-natural settings, antibiotics in ingested blood enhanced the susceptibility of *An. gambiae* to *Plasmodium* infection and increasing mosquito fitness through higher survival and fecundity. Moreover, the presence of antibiotics in the blood of malaria-infected people was identified as a risk for increased disease transmission (Gendrin et al., 2015). In addition, the experimental co-infections of bacteria with *Plasmodium* have showed the reduction of the number of developing oocysts in the mosquito midgut, both in laboratory and field conditions (Pumpuni et al., 1993; Lowenberger et al., 1999; Gonzalez-Ceron et al., 2003; Dong et al., 2009; Meister et al., 2009; Cirimotich et al., 2011). More recently, it was observed that expansion of midgut microbiota by negative regulation of *An. gambiae* immune response increased susceptibility to *Plasmodium* as observed before for microbiota depletion (Dennison et al., 2015). Finally, it's probably the microbiota equilibrium that confers resistance to *Plasmodium* infection. Then, the development of new approaches in malaria control needs the description of the microbiota of wild *Anopheles* species and its variations (Rani et al., 2009; Boissière et al., 2012; Chavshin et al., 2012; Osei-Poku et al., 2012; Gendrin and Christophides, 2013; Manguin et al., 2013; Ngo et al., 2015).

As previously done (Ngo et al., 2015), we present herein a combined approach associating molecular detection and culture that improve the performance of each separate method. About one third of the bacterial genera detected in this study were obtained by culture, thus providing a valuable collection of isolates for further experiments such as experimental challenges (Ramirez et al., 2014) and in paratransgenesis projects (Chavshin et al., 2014). Due to the low resolution of TTGE fingerprints when the number of band is high, PCR-TTGE is only suitable for the description of microbiota with relative low richness as demonstrated for human neonatal microbiota (Jacquot et al., 2011) and surgical microbiota (Romano-Bertrand et al., 2014), but also for the description of mosquito microbiota (Manguin et al., 2013; Ngo et al., 2015). By this mean, we previously described most of the known repertory of bacterial taxa found in Asian *Anopheles* as shown by Villegas and Pimenta (2014). TTGE fingerprinting gave a good description of majority bacterial populations in mosquito microbiota, with a rather good sensitivity, as demonstrated by the extent of the taxonomic richness described herein compared to other studies even those which used Next Generation Sequencing (NGS; Villegas and

Pimenta, 2014). Moreover, the use of PCR-TTGE in this study allowed a comparison with previous studies in Vietnam (Ngo et al., 2015), but also in Thailand (Manguin et al., 2013). NGS is now becoming a reference method for microbiota description and knowledge of mosquito-associated bacteria would likely benefit from studies by NGS. Because NGS provide huge amounts of data sometimes partially exploited, preliminary and comparative studies by TTGE or other community fingerprinting approaches allow asking scientific questions to be further in-depth explored by NGS.

The *Anopheles* collection of 100 wild specimens collected in Binh Phuoc Province was colonized by a great diversity of bacteria with, to our knowledge, 51 genera not yet described in anophelines. Thereby, this study widens the description of the *Anopheles* pan-microbiota. The huge diversity of *Anopheles* associated bacteria contrasts the low richness observed per specimen. This result is similar to that shown by Osei-Poku et al. (2012) in different mosquito genera, including *Anopheles*, in which bacterial richness per specimen is very low, nearly always dominated by a small number of taxa. This low richness could explain that only 78% of the specimens are positive in culture. Other reasons could be the trapping of some bacteria in the *Anopheles* tissues despite the preparative crushing step, the presence of viable but non-cultivable bacteria and the limited panel of culture medium used in the study. Culturomics approaches (Tandina et al., 2016) with the use of large panel of culture media and conditions could enhance the culture yield in next studies.

Within the bacteria detected in *Anopheles* specimens from Binh Phuoc, approximately half of the genera have previously been identified in the intra-abdominal microbiota of *Anopheles* populations from Thailand (Manguin et al., 2013), Vietnam (Manguin et al., 2013; Ngo et al., 2015), Cameroon (Boissière et al., 2012), Iran (Chavshin et al., 2012), India (Rani et al., 2009), or in reviews (Gendrin and Christophides, 2013; Minard et al., 2013; Hughes et al., 2014; Villegas and Pimenta, 2014).

Our results also showed that there is no link between bacterial diversity and host species or group of host species collected in Binh Phuoc Province. This result is in accordance with previous reports that rather showed the influence of the local environment of the sampling site, environment that gives a specific bacterial profile to mosquito specimens (Boissière et al., 2012; Manguin et al., 2013; Buck et al., 2016). The microbiota diversity of host species will also be significantly influenced by the origin of specimens like wild *versus* laboratory reared *Anopheles* mosquitoes (Chavshin et al., 2012). For instance, the selective pressure of laboratory conditions may limit bacteria acquisition at both larval and adult stages resulting in great reduction or nil cultivable bacteria in mosquito midgut (Riehle and Jacobs-Lorena, 2005; Favia et al., 2007; Gusmão et al., 2010). Studies on microbiota of *An. gambiae* larvae and pupae showed that only subset of bacteria from the aquatic habitat were able to inhabit the mosquito gut (Wang et al., 2011). The same influence of environment on *Anopheles* microbiota is likely to occur during the terrestrial life of adults. Considering the repertory of bacterial genera presented in **Figure 3**, the presence of soil- and plant-associated bacteria in *Anopheles* microbiota

is probable. For instance, *Agrococcus*, *Janibacter*, *Streptomyces*, *Bacillus* (and related genera), and *Lysobacter* are classical soil inhabitants, whereas *Asaia*, *Burkholderia*, *Comamonas*, *Delftia*, and *Sphingomonas* and *Methylobacterium* are frequently associated with plants. In addition, several prevalent bacterial genera are associated with animal or human microbiota, mainly to cutaneous microbiota: *Propionibacterium*, *Staphylococcus*, *Peptostreptococcus*, *Clostridium*, and *Acinetobacter*. These results suggest that bacteria from diverse origins could colonize adult *Anopheles*.

Interestingly, the *Anopheles* species from Binh Phuoc Province contained members of the genus *Enterobacter* sp. recently isolated from wild *An. gambiae* from Zambia. *Enterobacter* sp. received lot of attention due its anti-*Plasmodium* effect through the production of reactive oxygen species that directly target *Plasmodium* parasites in the midgut of *An. gambiae* (Cirimotich et al., 2011). More recently, another bacteria, *Chromobacterium Csp_P*, has been described as responsible of vectors refractoriness to *P. falciparum* and dengue virus (Ramirez et al., 2014). Then, further studies on mosquito microbiota should provide new insights on pathogens transmission and might open new ways to prevent vector-borne diseases.

Recently published studies have described new bacterial species, first isolated from *Anopheles* mosquitoes, such as *E. anophelis* (Kampfer et al., 2011), *Thorsellia anophelis* (Kampfer et al., 2006a), and *Janibacter anophelis* (Kampfer et al., 2006b). The latter species was accurately identified herein after bacterial culture and the species *J. terrae* described in soil was also found in Binh Phuoc *Anopheles*. Bacterial identification to the species level or below, defining variants within a bacterial species (genovar, genomovar, biovar, etc.), is needed to determine whether strictly identical bacterial populations are shared between environment, host and vector, or if *Anopheles*-associated bacteria are specific.

As a tendency to be confirmed, we also described herein the microbiota of two *Anopheles* (*An. dirus* and *An. pampanai*) naturally infected by *P. vivax*. The genus *Acinetobacter* solely was common to the two infected specimens but it was also one of the most prevalent bacterial species forming *Anopheles* microbiota. Indeed, it was frequently identified in the microbiota of both Asian and African anopheline species (Cameroon, Iran, India, Kenya, Mali, and Vietnam; Straif et al., 1998; Rani et al., 2009; Chavshin et al., 2012). Nevertheless, there is no available report on the implication of this genus for the development of *Plasmodium* in *Anopheles* and the interaction between this bacterial genus and malarial parasites in *Anopheles* might be studied.

The huge diversity of the microbial world even within a single species urges to overpass genus or species level for microbiota descriptions. Indeed, it is likely that the informative taxonomic level for host and microbes interactions is the subspecies level. Such a precise characterization needs availability of bacterial strains, highlighting again the interest of culture in microbiota studies. Moreover, the phenotype of bacterial strains could be of great interest to understand mechanisms involved in bacteria/*Plasmodium*/*Anopheles* interactions. This study provides 51 strains available for further experiments. Moreover, to the niche level, functional metagenomic and

metabolomic approaches should be undertaken to determine microbiota conditions that favor or not vector competence.

AUTHOR CONTRIBUTIONS

CN made the bacterial analyses (bacterial cultures and molecular analyses), and results' interpretation. SR-B helped for DNA sequencing, bacterial identification, and results' interpretation. SM contributed to the study design, the specimens collection and the results' interpretation. EJ-B supervised the study design, the bacterial analyses, and the results' interpretation. All the authors contributed to the manuscript redaction.

REFERENCES

Abe, T., Honda, S., Nakazawa, S., Tuong, T. D., Thieu, N. Q., Hung, L. X., et al. (2009). Risk factors for malaria infection among ethnic minorities in Binh Phuoc, Vietnam. *Southeast Asian J. Trop. Med. Public Health* 40, 18–29.

Beier, M. S., Pumpuni, C. B., Beier, J. C., and Davis, J. R. (1994). Effects of para-aminobenzoic acid, insulin, and gentamicin on *Plasmodium falciparum* development in anopheline mosquitoes (Diptera: Culicidae). *J. Med. Entomol.* 31, 561–565. doi: 10.1093/jmedent/31.4.561

Blumberg, B. J., Trop, S., Das, S., and Dimopoulos, G. (2013). Bacteria- and IMD pathway-independent immune defenses against *Plasmodium falciparum* in *Anopheles gambiae*. *PLoS ONE* 8:e72130. doi: 10.1371/journal.pone.0072130

Boissière, A., Tchioffo, M. T., Bachar, D., Abate, L., Marie, A., Nsango, S. E., et al. (2012). Midgut microbiota of the malaria mosquito vector *Anopheles gambiae* and interactions with *Plasmodium falciparum* infection. *PLoS Pathog* 8:e1002742. doi: 10.1371/journal.ppat.1002742

Briones, A. M., Shililu, J., Githure, J., Novak, R., and Raskin, L. (2008). *Thorsellia anophelis* is the dominant bacterium in a Kenyan population of adult *Anopheles gambiae* mosquitoes. *ISME J.* 2, 74–82. doi: 10.1038/ismej.2007.95

Buck, M., Nilsson, L. K., Brunius, C., Dabire, R. K., Hopkins, R., and Terenius, O. (2016). Bacterial associations reveal spatial population dynamics in *Anopheles gambiae* mosquitoes. *Sci. Rep.* 6, 22806. doi: 10.1038/srep22806

Chang, X. L., Xue, Y. Q., Zhang, A. D., Zhu, G. D., and Fang, Q. (2013). [Deltamethrin resistance, metabolic detoxification enzyme and kdr mutation in *Anopheles sinensis* in region along Huaihe River in Anhui Province]. *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi* 25, 263–267.

Chavshin, A. R., Oshaghi, M. A., Vatandoost, H., Pourmand, M. R., Raeisi, A., Enayati, A. A., et al. (2012). Identification of bacterial microflora in the midgut of the larvae and adult of wild caught *Anopheles stephensi*: a step toward finding suitable paratransgenesis candidates. *Acta Trop.* 121, 129–134. doi: 10.1016/j.actatropica.2011.10.015

Chavshin, A. R., Oshaghi, M. A., Vatandoost, H., Pourmand, M. R., Raeisi, A., and Terenius, O. (2014). Isolation and identification of culturable bacteria from wild *Anopheles culicifacies*, a first step in a paratransgenesis approach. *Parasit Vectors* 7, 419. doi: 10.1186/1756-3305-7-419

Cirimotich, C. M., Dong, Y., Clayton, A. M., Sandiford, S. L., Souza-Neto, J. A., Mulenga, M., et al. (2011). Natural microbe-mediated refractoriness to *Plasmodium* infection in *Anopheles gambiae*. *Science* 332, 855–858. doi: 10.1126/science.1201618

Dabrowski, J. M., and Balderacchi, M. (2013). Development and field validation of an indicator to assess the relative mobility and risk of pesticides in the Lourens River catchment, South Africa. *Chemosphere* 93, 2433–2443. doi: 10.1016/j.chemosphere.2013.08.070

Dabrowski, J. M., Shadung, J. M., and Wepener, V. (2014). Prioritizing agricultural pesticides used in South Africa based on their environmental mobility and potential human health effects. *Environ. Int.* 62, 31–40. doi: 10.1016/j.envint.2013.10.001

Dennison, N. J., BenMarzouk-Hidalgo, O. J., and Dimopoulos, G. (2015). MicroRNA-regulation of *Anopheles gambiae* immunity to *Plasmodium falciparum* infection and midgut microbiota. *Dev. Comp. Immunol.* 49, 170–178. doi: 10.1016/j.dci.2014.10.016

ACKNOWLEDGMENTS

This study was made possible through a PhD allowance provided to the CN by the Institute of Research for Development (IRD), France. This work was also supported by the association ADEREMPHA, Sauzet, France.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.02095/full#supplementary-material>

Dennison, N. J., Jupatanakul, N., and Dimopoulos, G. (2014). The mosquito microbiota influences vector competence for human pathogens. *Curr. Opin. Insect Sci.* 3, 6–13. doi: 10.1016/j.cois.2014.07.004

Dong, Y., Aguilar, R., Xi, Z., Warr, E., Mongin, E., and Dimopoulos, G. (2006). *Anopheles gambiae* immune responses to human and rodent *Plasmodium* parasite species. *PLoS Pathog* 2:e52. doi: 10.1371/journal.ppat.0020052

Dong, Y., Das, S., Cirimotich, C., Souza-Neto, J. A., McLean, K. J., and Dimopoulos, G. (2011). Engineered anopheles immunity to *Plasmodium* infection. *PLoS Pathog* 7:e1002458. doi: 10.1371/journal.ppat.1002458

Dong, Y., Manfredini, F., and Dimopoulos, G. (2009). Implication of the mosquito midgut microbiota in the defense against malaria parasites. *PLoS Pathog* 5:e1000423. doi: 10.1371/journal.ppat.1000423

Drancourt, M., Bollet, C., Carlioz, A., Martelin, R., Gayral, J. P., and Raoult, D. (2000). 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J. Clin. Microbiol.* 38, 3623–3630.

Favia, G., Ricci, I., Damiani, C., Raddadi, N., Crotti, E., Marzorati, M., et al. (2007). Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proc. Natl. Acad. Sci. U.S.A.* 104, 9047–9051. doi: 10.1073/pnas.0610451104

Gafan, G. P., Lucas, V. S., Roberts, G. J., Petrie, A., Wilson, M., and Spratt, D. A. (2005). Statistical analyses of complex denaturing gradient gel electrophoresis profiles. *J. Clin. Microbiol.* 43, 3971–3978. doi: 10.1128/JCM.43.8.3971-3978.2005

Garros, C., Harbach, R. E., and Manguin, S. (2005). Morphological assessment and molecular phylogenetics of the *Funestus* and *Minimus* groups of *Anopheles* (Cellia). *J. Med. Entomol.* 42, 522–536. doi: 10.1603/0022-25852005042[0522:MAAMPO]2.0.CO;2

Gendrin, M., and Christophides, G. K. (2013). “The *Anopheles* mosquitoes microbiota and their impact on pathogen transmission,” in *Anopheles mosquitoes - New insights into malaria vectors*, ed. S. Manguin (Rijeka: InTech Open Access), 525–548.

Gendrin, M., Rodgers, F. H., Yerbanga, R. S., Ouedraogo, J. B., Basanez, M. G., Cohuet, A., et al. (2015). Antibiotics in ingested human blood affect the mosquito microbiota and capacity to transmit malaria. *Nat. Commun.* 6, 5921. doi: 10.1038/ncomms6921

Gonzalez-Ceron, L., Santillan, F., Rodriguez, M. H., Mendez, D., and Hernandez-Avila, J. E. (2003). Bacteria in midguts of field-collected *Anopheles albimanus* block *Plasmodium vivax* sporogonic development. *J. Med. Entomol.* 40, 371–374. doi: 10.1603/0022-2585-40.3.371

Gusmão, D. S., Santos, A. V., Marini, D. C., Bacci, M. Jr., Berbert-Molina, M. A., and Lemos, F. J. (2010). Culture-dependent and culture-independent characterization of microorganisms associated with *Aedes aegypti* (Diptera: Culicidae) (L.) and dynamics of bacterial colonization in the midgut. *Acta Trop* 115, 275–281. doi: 10.1016/j.actatropica.2010.04.011

Hughes, G. L., Dodson, B. L., Johnson, R. M., Murdock, C. C., Tsujimoto, H., Suzuki, Y., et al. (2014). Native microbiome impedes vertical transmission of *Wolbachia* in *Anopheles* mosquitoes. *Proc. Natl. Acad. Sci. U.S.A.* 111, 12498–12503. doi: 10.1073/pnas.1408881111

Huong, N. M., Hewitt, S., Davis, T. M., Dao, L. D., Toan, T. Q., Kim, T. B., et al. (2001). Resistance of *Plasmodium falciparum* to antimalarial drugs in

- a highly endemic area of southern Viet Nam: a study in vivo and in vitro. *Trans. R. Soc. Trop. Med. Hyg.* 95, 325–329. doi: 10.1016/S0035-9203(01)90254-8
- Jacquot, A., Neveu, D., Aujoulat, F., Mercier, G., Marchandin, H., Jumas-Bilak, E., et al. (2011). Dynamics and clinical evolution of bacterial gut microflora in extremely premature patients. *J. Pediatr.* 158, 390–396. doi: 10.1016/j.jpeds.2010.09.007
- Kabula, B., Tungu, P., Malima, R., Rowland, M., Minja, J., Wililo, R., et al. (2013). Distribution and spread of pyrethroid and DDT resistance among the *Anopheles gambiae* complex in Tanzania. *Med. Vet. Entomol.* 28, 244–252. doi: 10.1111/mve.12036
- Kampfer, P., Lindh, J. M., Terenius, O., Haghdoost, S., Falsen, E., Busse, H. J., et al. (2006a). *Thorsellia anophelis* gen. nov., sp. nov., a new member of the *Gammaproteobacteria*. *Int. J. Syst. Evol. Microbiol.* 56, 335–338. doi: 10.1099/ijs.0.63999-0
- Kampfer, P., Terenius, O., Lindh, J. M., and Faye, I. (2006b). *Janibacter anophelis* sp. nov., isolated from the midgut of *Anopheles arabiensis*. *Int. J. Syst. Evol. Microbiol.* 56, 389–392. doi: 10.1099/ijs.0.63905-0
- Kampfer, P., Matthews, H., Glaeser, S. P., Martin, K., Lidders, N., and Faye, I. (2011). *Elizabethkingia anophelis* sp. nov., isolated from the midgut of the mosquito *Anopheles gambiae*. *Int. J. Syst. Evol. Microbiol.* 61, 2670–2675. doi: 10.1099/ijs.0.026393-0
- Ledder, R. G., Gilbert, P., Huws, S. A., Aarons, L., Ashley, M. P., Hull, P. S., et al. (2007). Molecular analysis of the subgingival microbiota in health and disease. *Appl. Environ. Microbiol.* 73, 516–523. doi: 10.1128/AEM.01419-06
- Lowenberger, C. A., Kamal, S., Chiles, J., Paskewitz, S., Bulet, P., Hoffmann, J. A., et al. (1999). Mosquito-Plasmodium interactions in response to immune activation of the vector. *Exp. Parasitol.* 91, 59–69. doi: 10.1006/expr.1999.4350
- Manguin, S., Ngo, C. T., Tainchum, K., Juntarajumnong, W., Chareonviriyaphap, T., Michon, A. L., et al. (2013). “Bacterial biodiversity in midguts of *Anopheles* mosquitoes, malaria vectors in Southeast Asia,” in *Anopheles mosquitoes - New insights into malaria vectors*, ed. S. Manguin (Rijeka: InTech Open Access), 549–576.
- Meister, S., Agianian, B., Turlure, F., Relogio, A., Morlais, I., Kafatos, F. C., et al. (2009). *Anopheles gambiae* PGRPLC-mediated defense against bacteria modulates infections with malaria parasites. *PLoS Pathog* 5:e1000542. doi: 10.1371/journal.ppat.1000542
- Minard, G., Mavingui, P., and Moro, C. V. (2013). Diversity and function of bacterial microbiota in the mosquito holobiont. *Parasit Vectors* 6, 146. doi: 10.1186/1756-3305-6-146
- Ngo, C. T., Aujoulat, F., Veas, F., Jumas-Bilak, E., and Manguin, S. (2015). Bacterial diversity associated with wild caught *Anopheles* mosquitoes from Dak Nong Province, Vietnam using culture and DNA fingerprint. *PLoS ONE* 10:e0118634. doi: 10.1371/journal.pone.0118634
- Ngo, C. T., Dubois, G., Sinou, V., Parzy, D., Le, H. Q., Harbach, R. E., et al. (2014). Diversity of *Anopheles* mosquitoes in Binh Phuoc and Dak Nong Provinces of Vietnam and their relation to disease. *Parasit Vectors* 7, 316. doi: 10.1186/1756-3305-7-316
- Nguyen, M. H., Davis, T. M., Cox-Singh, J., Hewitt, S., Tran, Q. T., Tran, B. K., et al. (2003). Treatment of uncomplicated falciparum malaria in southern Vietnam: can chloroquine or sulfadoxine-pyrimethamine be reintroduced in combination with artesunate? *Clin. Infect. Dis.* 37, 1461–1466. doi: 10.1086/379323
- Ogier, J. C., Son, O., Gruss, A., Tailliez, P., and Delacroix-Buchet, A. (2002). Identification of the bacterial microflora in dairy products by temporal temperature gradient gel electrophoresis. *Appl. Environ. Microbiol.* 68, 3691–3701. doi: 10.1128/AEM.68.8.3691-3701.2002
- Osei-Poku, J., Mbogo, C. M., Palmer, W. J., and Jiggins, F. M. (2012). Deep sequencing reveals extensive variation in the gut microbiota of wild mosquitoes from Kenya. *Mol. Ecol.* 21, 5138–5150. doi: 10.1111/j.1365-294X.2012.05759.x
- Pumpuni, C. B., Beier, M. S., Nataro, J. P., Guers, L. D., and Davis, J. R. (1993). *Plasmodium falciparum*: inhibition of sporogonic development in *Anopheles stephensi* by gram-negative bacteria. *Exp. Parasitol.* 77, 195–199. doi: 10.1006/expr.1993.1076
- Ramirez, J. L., Short, S. M., Bahia, A. C., Saraiva, R. G., Dong, Y., Kang, S., et al. (2014). *Chromobacterium Csp_P* reduces malaria and dengue infection in vector mosquitoes and has entomopathogenic and in vitro anti-pathogen activities. *PLoS Pathog* 10:e1004398. doi: 10.1371/journal.ppat.1004398
- Rani, A., Sharma, A., Rajagopal, R., Adak, T., and Bhatnagar, R. K. (2009). Bacterial diversity analysis of larvae and adult midgut microflora using culture-dependent and culture-independent methods in lab-reared and field-collected *Anopheles stephensi*-an Asian malarial vector. *BMC Microbiol.* 9:96. doi: 10.1186/1471-2180-9-96
- Riehle, M. A., and Jacobs-Lorena, M. (2005). Using bacteria to express and display anti-parasite molecules in mosquitoes: current and future strategies. *Insect Biochem. Mol. Biol.* 35, 699–707. doi: 10.1016/j.ibmb.2005.02.008
- Rodrigues, J., Brayner, F. A., Alves, L. C., Dixit, R., and Barillas-Mury, C. (2010). Hemocyte differentiation mediates innate immune memory in *Anopheles gambiae* mosquitoes. *Science* 329, 1353–1355. doi: 10.1126/science.1190689
- Romano-Bertrand, S., Frapier, J. M., Calvet, B., Colson, P., Albat, B., Parer, S., et al. (2014). Dynamics of the surgical microbiota along the cardiothoracic surgery pathway. *Front. Microbiol.* 5:787. doi: 10.3389/fmicb.2014.00787
- Romano-Bertrand, S., Parer, S., Lotthé, A., Colson, P., Albat, B., and Jumas-Bilak, E. (2012). “Temporal temperature gel electrophoresis to survey pathogenic bacterial communities: the case of surgical site infections,” in *Gel Electrophoresis - Advanced Techniques*, ed. S. Magdeldin (Rijeka: InTech Open Access), 500.
- Roudiere, L., Jacquot, A., Marchandin, H., Aujoulat, F., Devine, R., Zorogniotti, I., et al. (2009). Optimized PCR-Temporal Temperature Gel Electrophoresis compared to cultivation to assess diversity of gut microbiota in neonates. *J. Microbiol. Methods* 79, 156–165. doi: 10.1016/j.mimet.2009.08.005
- Sharma, P., Sharma, S., Maurya, R. K., Das De, T., Thomas, T., Lata, S., et al. (2014). Salivary glands harbor more diverse microbial communities than gut in *Anopheles culicifacies*. *Parasit. Vectors* 7:235. doi: 10.1186/1756-3305-7-235
- Singleton, S. T., Lein, P. J., Farahat, F. M., Farahat, T., Bonner, M. R., Knaak, J. B., et al. (2013). Characterization of alpha-cypermethrin exposure in Egyptian agricultural workers. *Int. J. Hyg. Environ. Health* 217, 538–545. doi: 10.1016/j.ijheh.2013.10.003
- Straif, S. C., Mbogo, C. N., Toure, A. M., Walker, E. D., Kaufman, M., Toure, Y. T., et al. (1998). Midgut bacteria in *Anopheles gambiae* and *An. funestus* (Diptera: Culicidae) from Kenya and Mali. *J. Med. Entomol.* 35, 222–226. doi: 10.1093/jmedent/35.3.222
- Tandina, F., Almeras, L., Kone, A. K., Doumbo, O. K., Raout, D., and Parola, P. (2016). Use of MALDI-TOF MS and culturomics to identify mosquitoes and their midgut microbiota. *Parasit Vectors* 9, 495. doi: 10.1186/s13071-016-1776-y
- Tangena, J. A., Adiamoh, M., D'Alessandro, U., Jarju, L., Jawara, M., Jeffries, D., et al. (2013). Alternative Treatments for Indoor Residual Spraying for Malaria Control in a Village with Pyrethroid- and DDT-Resistant Vectors in The Gambia. *PLoS ONE* 8:e74351. doi: 10.1371/journal.pone.0074351
- Thanh, N. V., Toan, T. Q., Cowman, A. F., Casey, G. J., Phuc, B. Q., Tien, N. T., et al. (2010). Monitoring for *Plasmodium falciparum* drug resistance to artemisinin and artesunate in Binh Phuoc Province, Vietnam: 1998–2009. *Malar. J.* 9, 181. doi: 10.1186/1475-2875-9-181
- Tran, T. H., Nguyen, T. T., Nguyen, H. P., Boni, M. F., Ngo, V. T., Nguyen, T. N., et al. (2012). In vivo susceptibility of *Plasmodium falciparum* to artesunate in Binh Phuoc Province, Vietnam. *Malar. J.* 11, 355. doi: 10.1186/1475-2875-11-355
- Verhaeghen, K., Van Bortel, W., Trung, H. D., Sochantha, T., and Coosemans, M. (2009). Absence of knockdown resistance suggests metabolic resistance in the main malaria vectors of the Mekong region. *Malar. J.* 8, 84. doi: 10.1186/1475-2875-8-84
- Verhaeghen, K., Van Bortel, W., Trung, H. D., Sochantha, T., Keokenchanh, K., and Coosemans, M. (2010). Knockdown resistance in *Anopheles vagus*, *An. sinensis*, *An. paraliae* and *An. peditaeniatus* populations of the Mekong region. *Parasit. Vectors* 3, 59. doi: 10.1186/1756-3305-3-59
- Villegas, L. M., and Pimenta, P. F. (2014). Metagenomics, paratransgenesis and the *Anopheles* microbiome: a portrait of the geographical distribution of the anopheline microbiota based on a meta-analysis of reported taxa. *Mem. Inst. Oswaldo Cruz* 109, 672–684. doi: 10.1590/0074-0276140194
- Walton, C., Handley, J. M., Kuvangkadilok, C., Collins, F. H., Harbach, R. E., Baimai, V., et al. (1999). Identification of five species of the *Anopheles dirus* complex from Thailand, using allele-specific polymerase chain reaction. *Med. Vet. Entomol.* 13, 24–32. doi: 10.1046/j.1365-2915.1999.00142.x

- Walton, C., Somboon, P., O'Loughlin, S. M., Zhang, S., Harbach, R. E., Linton, Y. M., et al. (2007). Genetic diversity and molecular identification of mosquito species in the *Anopheles maculatus* group using the ITS2 region of rDNA. *Infect. Genet. Evol.* 7, 93–102. doi: 10.1016/j.meegid.2006.05.001
- Wang, Y., Gilbreath, T. M. III, Kukutla, P., Yan, G., and Xu, J. (2011). Dynamic gut microbiome across life history of the malaria mosquito *Anopheles gambiae* in Kenya. *PLoS ONE* 6:e24767. doi: 10.1371/journal.pone.0024767
- WHO (2012). *World Malaria Report*. Geneva: WHO Press.
- WHO (2015). *World Malaria Report*. Geneva: World Health Organization.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Ngo, Romano-Bertrand, Manguin and Jumas-Bilak. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.