



Corrigendum: Thymoquinone Inhibits Virulence Related Traits of *Cronobacter sakazakii* ATCC 29544 and Has Anti-biofilm Formation Potential

Chao Shi¹, Chunhong Yan^{2,3}, Yue Sui¹, Yi Sun¹, Du Guo¹, Yifei Chen¹, Tong Jin¹, Xiaoli Peng¹, Linlin Ma⁴ and Xiaodong Xia^{1*}

¹ College of Food Science and Engineering, Northwest A&F University, Yangling, China, ² College of Animal Science and Technology, Northwest A&F University, Yangling, China, ³ School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, China, ⁴ Xi'an Yurun Agricultural Products Global Sourcing Co., LTD., Xi'an, China

Keywords: *Cronobacter sakazakii*, Thymoquinone, sub-inhibitory concentration, HT-29 cells, Virulence Factors

A corrigendum on

Thymoquinone Inhibits Virulence Related Traits of *Cronobacter sakazakii* ATCC 29544 and Has Anti-biofilm Formation Potential

by Shi, C., Yan, C., Sui, Y., Sun, Y., Guo, D., Chen, Y., et al. (2017). *Front. Microbiol.* 8:2220. doi: 10.3389/fmicb.2017.02220

OPEN ACCESS

Edited and reviewed by:

Lanming Chen,
Shanghai Ocean University, China

*Correspondence:

Xiaodong Xia
foodscixiaodong@yahoo.com

Specialty section:

This article was submitted to
Food Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 05 January 2018

Accepted: 07 February 2018

Published: 20 February 2018

Citation:

Shi C, Yan C, Sui Y, Sun Y, Guo D, Chen Y, Jin T, Peng X, Ma L and Xia X (2018) Corrigendum: Thymoquinone Inhibits Virulence Related Traits of *Cronobacter sakazakii* ATCC 29544 and Has Anti-biofilm Formation Potential. *Front. Microbiol.* 9:290. doi: 10.3389/fmicb.2018.00290

In the first paragraph in the “Introduction” of this article, the number of *Cronobacter* species was incorrectly described as ten. Actually, *Cronobacter pulveris*, *Cronobacter helveticus* and *Cronobacter turicensis* were removed from the genus in 2014.

The original sentence should be corrected as follows: *Cronobacter* spp. is currently considered to consist of 7 species: *Cronobacter sakazakii*, *C. malonaticus*, *C. universalis*, *C. dublinensis*, *C. muytjensii*, *C. condiment*, and *C. zurichensis* (Stephan et al., 2014), among which *C. sakazakii* is one of the two group 1 clinically relevant species that form the majority of the clinical isolates. The authors regret this error.

The original article has been updated.

REFERENCES

- Stephan, R., Grim, C. J., Gopinath, G. R., Mammel, M. K., Sathyamoorthy, V., Trach, L. H., et al. (2014). Re-examination of the taxonomic status of *Enterobacter helveticus*, *Enterobacter pulveris* and *Enterobacter turicensis* as members of the genus *Cronobacter* and their reclassification in the genera *Franconibacter* gen. nov. and *Siccibacter* gen. nov. as *Franconibacter helveticus* comb. nov., *Franconibacter pulveris* comb. nov. and *Siccibacter turicensis* comb. nov., respectively. *Int. J. Syst. Evol. Microbiol.* 64, 3402–3410. doi: 10.1099/ijs.0.059832-0

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Shi, Yan, Sui, Sun, Guo, Chen, Jin, Peng, Ma and Xia. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.