



Corrigendum: TaqMan qPCR for Quantification of *Clonostachys rosea* Used as a Biological Control Agent Against *Fusarium graminearum*

OPEN ACCESS

Approved by:
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

***Correspondence:**
Arja Laitila
arja.laitila@outlook.com
Susanne Vogelgsang
susanne.vogelgsang@
agroscope.admin.ch

Specialty section:
This article was submitted to
Fungi and Their Interactions,
a section of the journal
Frontiers in Microbiology

Received: 24 July 2019

Accepted: 25 July 2019

Published: 13 August 2019

Citation:
Gimeno A, Sohlberg E, Pakula T,
Limnell J, Keller B, Laitila A and
Vogelgsang S (2019) Corrigendum:
TaqMan qPCR for Quantification of
Clonostachys rosea Used as a
Biological Control Agent Against
Fusarium graminearum.
Front. Microbiol. 10:1839.
doi: 10.3389/fmicb.2019.01839

Alejandro Gimeno^{1,2}, Elina Sohlberg³, Tiina Pakula³, Jenni Limnell³, Beat Keller²,
Arja Laitila^{3*} and Susanne Vogelgsang^{1*}

¹ Ecological Plant Protection in Arable Crops, Research Division Plant Protection, Agroscope, Zurich, Switzerland, ² Molecular Plant Biology and Phytopathology, Department of Plant and Microbial Biology, University of Zurich, Zurich, Switzerland, ³ VTT Technical Research Centre of Finland Ltd., Espoo, Finland

Keywords: qPCR, *Clonostachys rosea*, biological control agent, *Fusarium graminearum*, *Fusarium* head blight, MycoKey

A Corrigendum on

TaqMan qPCR for Quantification of *Clonostachys rosea* Used as a Biological Control Agent Against *Fusarium graminearum*

by Gimeno, A., Sohlberg, E., Pakula, T., Limnell, J., Keller, B., Laitila, A., et al. (2019). *Front. Microbiol.* 10:1627. doi: 10.3389/fmicb.2019.01627

In the original article, there was a mistake in **Table 2** as published. The primer sequence (5'-3') of *VTact*—reverse contained a false nucleobase. The corrected **Table 2** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2019 Gimeno, Sohlberg, Pakula, Limnell, Keller, Laitila and Vogelgsang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 2 | Primer and hydrolysis probe sequences (5'–3') for the specific amplification of *Clonostachys rosea* with TaqMan qPCR.

Primer/probe	Sequence (5' to 3')
<i>VTTact</i> -forward	GGCCAGAGATTGTGTTGATGA
<i>VTTact</i> -reverse	ACAGGTTAGGCTCAATGCTC
<i>VTTact</i> probe	GAGGCTGGCAAGAGAGGTCAGTCAC