



Distribution of HCV Genotypes Among People Who Inject Drugs in Tunisia: New Evidence for Scaling Up Prevention and Treatment Toward National Elimination Goal

OPEN ACCESS

Edited by:

Anna Kramvis,
University of the Witwatersrand,
South Africa

Reviewed by:

Xavier López-Labrador,
Fundación para el Fomento de la
Investigación Sanitaria y Biomédica
de la Comunitat Valenciana, Spain

Jordi Gómez,
Instituto de Parasitología y
Biomedicina López-Neyra, Spain

*Correspondence:

Anissa Chouikha
chouikhaanissa@gmail.com
orcid.org/0000-0001-5616-0204
Amine Ghrabi
dr.ghrabi@gmail.com
orcid.org/0000-0003-0454-3339

Specialty section:

This article was submitted to
Virology,
a section of the journal
Frontiers in Microbiology

Received: 20 April 2021

Accepted: 24 June 2021

Published: 27 July 2021

Citation:

Chouikha A, Ghrabi A,
Ghodbane A, Hammemi W,
Khedhiri M, Sadraoui A, Touzi H,
Hassine HB, Maatoug S,
Bensaoud C, Abdelhak S,
Bouarrouj S, Gdoura M, Chaouachi H
and Triki H (2021) Distribution of HCV
Genotypes Among People Who Inject
Drugs in Tunisia: New Evidence
for Scaling Up Prevention
and Treatment Toward National
Elimination Goal.
Front. Microbiol. 12:697859.
doi: 10.3389/fmicb.2021.697859

Anissa Chouikha^{1*}, Amine Ghrabi^{2*}, Amira Ghodbane¹, Walid Hammemi¹,
Marwa Khedhiri¹, Amel Sadraoui¹, Henda Touzi¹, Hichem Ben Hassine³,
Sonia Maatoug³, Chaima Bensaoud^{3,4}, Sonia Abdelhak³, Samir Bouarrouj²,
Mariem Gdoura¹, Hedia Chaouachi² and Henda Triki¹

¹ Laboratory of Clinical Virology, WHO Reference Laboratory for Poliomyelitis and Measles in the Eastern Mediterranean Region, Pasteur Institute of Tunis, Tunis El Manar University, Tunis, Tunisia, ² Association Tunisienne d'Information et d'Orientation sur le SIDA et la Toxicomanie (Tunisian Association for Information and Orientation on HIV/AIDS and Toxicomania – ATIOST), Tunis, Tunisia, ³ Science Shop, Institut Pasteur de Tunis, Tunis, Tunisia, ⁴ Institute of Parasitology, Biology Centre, Czech Academy of Sciences, Ceske Budejovice, Czechia

Little is known about the distribution of hepatitis C virus (HCV) genotypes among people who inject drugs (PWID) in North African countries, including Tunisia. This study aims to describe HCV genotypes circulating among Tunisian PWID. A cross-sectional study was conducted, and 128 HCV-positive PWID were recruited between 2018 and 2019 from community-based harm reduction centers. After informed consent, sociodemographic characteristics and risk behavior data were obtained using an interviewer-administrated questionnaire. Blood samples were collected for further serological and molecular testing. Overall, five women and 123 men were included. The median age was 39.5 years. The majority of PWID (56.3%) had less than a secondary level of education, were single (57%), were unemployed (65.6%), were incarcerated at least once (93.0%), and had a history of residency in at least one foreign country (50.8%). During the previous 12 months, 82.0% reported having reused syringes at least once, 43.8% shared syringes at least once, while 56.2% had at least one unprotected sexual relation, and 28.1% had more than two different sexual partners. Tattooing was reported among 60.2%. All positive results for HCV-infection by rapid testing were confirmed by enzyme-linked immunosorbent assay (ELISA). HCV-RNA was detectable in 79.7%. Genotyping showed a predominance of genotype 1 (52%) followed by genotype 3 (34%) and genotype 4 (10%). Four patients (4%) had an intergenotype mixed infection. Subtyping showed the presence of six different HCV subtypes as follows: 1a (53.2%), 1b (6.4%), 3a (33.0%), 4a (3.2%), and 4d (4.3%). This is the first study describing circulating HCV genotypes among PWID in Tunisia. The distribution of HCV genotypes is distinct from

the general population with a predominance of subtypes 1a and 3a. These findings can be used to guide national efforts aiming to optimize the access of PWID to relevant HCV prevention and treatment measures including pangenotypic regimens for patients infected with HCV genotype 3.

Keywords: hepatitis C virus, injecting drug users, PWID, prevention, treatment, substance abuse, Tunisia

INTRODUCTION

Infection with hepatitis C virus (HCV) remains a major global health problem with significant morbidity and mortality due to cirrhosis and hepatocellular carcinoma. The World Health Organization (WHO) estimates that 71.1 million people are living with HCV worldwide, out of which the majority remain untreated (World Health Organization, 2018b). In 2015, 23% of newly diagnosed HCV infections were attributed to injecting drug use (IDU) (World Health Organization, 2017). Due to sharing of injection equipment, people who inject drugs (PWID) are disproportionately affected by blood-borne viruses including HCV. About 52.3% of the global PWID population (15.6 million) had been exposed to HCV (Degenhardt et al., 2017) and PWID populations accounted for 8.5% of all HCV infections globally (Grebely et al., 2019).

The Global Health Sector Strategy on Viral Hepatitis 2016–2021 of the WHO recognizes harm reduction services for PWID and HCV treatment as key interventions for the elimination of HCV as a global public health threat by 2030 (World Health Organization, 2016). Harm reduction services includes testing for human immunodeficiency virus (HIV), HCV, and hepatitis B virus (HBV) infections (Martin et al., 2013), as well as syringe services programs (SSPs), which provide PWID with sterile syringes to minimize unsafe injecting risk behaviors (Palmateer et al., 2010; MacArthur et al., 2014). The recent advent of highly efficacious direct-acting antiviral (DAA) therapies (Ara and Paul, 2015) and their associated cure and prevention benefits (Martin et al., 2013) provide an unprecedented opportunity to reduce HCV-related morbidity and mortality among PWID (Grebely et al., 2017a).

The Middle East and North Africa (MENA) region has been identified as the most affected region by HCV infection worldwide (World Health Organization, 2017). There are significant variations for national HCV prevalence levels between MENA countries (Chaabna et al., 2018). While 15% of the adults were infected by HCV in Egypt, the overall HCV prevalence across the Maghreb subregion was about 1% with IDU identified as a significant contributor to HCV transmission in these countries (Fadlalla et al., 2015). Notably, in Tunisia, high levels of HCV prevalence among local groups of PWID were estimated, ranging from 21.7 to 29.1% (Fadlalla et al., 2015; Chaabna et al., 2018; Grebely et al., 2019).

In addition, it has been established that HCV genotypes' distribution varies significantly between PWID and the general population, with IDU as a key vector for the diversification of circulating viral genotypes (Ruta and Cernescu, 2015). A systematic review published in 2016 concluded that HCV genotypes 1a and 3 were the most common among populations

of PWID globally (Robaey et al., 2016). While the available estimations for the MENA region were limited to five countries only, this trend was confirmed in neighboring Libya and Morocco where genotypes 1 and 3 accounted for 86.6 and 90.5% of PWID, respectively (Alashek and Altagdi, 2008; Trimbilas et al., 2014). In Tunisia, genotype 1b had been identified as the most common genotype in the general population (79.5–82.6%) followed by genotype 2 (10.1–13.3%) (Djebbi et al., 2004; Mejri et al., 2005; Gower et al., 2014; Chouikha et al., 2021). However, to our knowledge, there are no published studies describing the distribution of HCV genotypes among PWID up to now. The absence of such studies suggests that the distribution of HCV genotypes in Tunisia might be underestimated especially among PWID groups. Therefore, the aim of this work is to describe the HCV genotypes circulating among a Tunisian population of PWID attending a harm reduction program managed by the Tunisian Association for Information and Orientation on HIV/AIDS and Toxicomania (ATIOST).

MATERIALS AND METHODS

Study Population and Design

Participants were recruited through word of mouth and self-referral at a harm reduction program managed by the community-based organization ATIOST. This program provides peer educators and active PWID with free and confidential access to sterile injection equipment as well as HIV/HCV screening across four harm reduction centers covering the North Eastern regions of Greater Tunis and Bizerta. All participants who (1) provided signed informed consent, (2) had an available positive rapid test for HCV antibody detection, (3) completed an interviewer-based questionnaire, and (4) had an available blood sample collected at one of ATIOST harm reduction centers between February 19, 2018 and January 24, 2019 were considered eligible for inclusion. Sociodemographic characteristics and risk behaviors data were collected at enrollment using an interviewer-administrated questionnaire. Further serologic and molecular testing were performed at the Laboratory of Clinical Virology of IPT.

Screening for HCV, HIV, and HBV Infection

Hepatitis C virus and HIV antibody detection was performed at enrollment using the appropriate available rapid test (SD Bioline HCV, Labman® HCV Test, or Vikia® Anti-HCV, and Turklab® Anti-HIV 1/2 Test or Alere Determine® HIV-1/2 Ag/Ab Combo). Confirmation of HCV positivity was carried out at the laboratory by commercial third-generation enzyme-linked

immunosorbent assay (ELISA) from Murex-Diagnostics-France (Murex anti-HCV) according to the manufacturers' instructions. Serologic detection of HBV antigen (AgHBs) and HBV antibodies (anti-HBc) was also carried out for confirmed HCV-positive samples using commercial ELISA from BIORAD-Diagnostics according to the manufacturers' instructions.

Hepatitis C Virus RNA Quantification and Genotyping

The detection and quantification of HCV RNA was conducted on plasma samples using real-time RT-PCR test from Roche Diagnostics (COBAS® AmpliPrep/COBAS® TaqMan®) with a limit of detection of 15 IU/ml [1 International Unit (IU) = 5.81 genome copies]. For samples with detectable HCV RNA, genotyping kits of HCV real-time RT-PCR Cobas 4800 HCV were used based on a set of primers that targets core, NS5b, and 5'UTR regions of the HCV genome for genotype identification. It allows the detection of genotypes 1a, 1b, 2, 3, 4, 5, and 6 as well as intergenotype mixed infections. In a next step, a selected subset of samples including non-determined genotypes and/or subtypes was selected to be assessed by nucleotide sequencing in the NS5b gene as described previously (Chouikha et al., 2021). Obtained sequences were submitted to GenBank database under accession numbers MW221786–MW221821 and MW678731–MW678756. Sequence alignment was performed by Clustal W within the software “Molecular Evolutionary Genetic Analyses” (MEGA) software version 7.0.26 and phylogenetic tree was also constructed using by the maximum likelihood method and the kimura-2 parameter model using MEGA software (Kumar et al., 2016). Topology was supported by 1,000 bootstrap replicates.

RESULTS

In total, 153 participants signed the informed consent form, out of which 128 had a positive rapid anti-HCV test result at enrollment, completed the interviewer-based questionnaire, and had an available blood sample for further laboratory serological and molecular testing (Figure 1).

Sociodemographic Characteristics and Risk Behaviors

Sociodemographic characteristics of the study population are described in Table 1. Among the 128 included PWID, there was a total of five women and 123 men. The age ranged from 23 to 61 years with a median value of 39.5 years (25–75 IQR = 34–46). The majority of PWID had less than a secondary level of education ($n = 72$; 56.3%), were single ($n = 73$; 57%), were unemployed ($n = 84$; 65.6%), and were incarcerated at least once ($n = 119$; 93.0%). Moreover, 85.9% ($n = 110$) were living with their family members, while 14.1% ($n = 18$) were living alone. Among 65 participants (50.8%) who had a history of residency in at least one foreign country, specific information regarding countries of residency was obtained for 57 participants. Most of these PWID lived in European countries ($n = 54$), especially Italy ($n = 47$) and/or France ($n = 27$). Only seven PWID had a history

of residency at least in one country from the Maghreb region, while five PWID lived in at least one country from the broader MENA region. A history of tattooing was reported among 60.2% ($n = 77$). Regarding injection risk behaviors during the last 12 months, 82.0% ($n = 105$) reported having reused syringes at least once, and 43.8% ($n = 56$) reported having shared syringes at least once. For sexual risk behaviors during the last 12 months, 56.2% ($n = 72$) reported having at least one unprotected sexual relation and 28.1% ($n = 36$) had more than two different sexual partners. Overall, 10.2% ($n = 13$) of HCV-positive PWID had a coinfection with HIV, whereas 3.9% ($n = 5$) were positive for HBsAg, and 83.6% ($n = 107$) were positive for anti-HBc. Based on these results, a previous exposure to all of HIV, HBV, and HCV infections was observed among 9.4% ($n = 12$).

Hepatitis C Virus Serological and Molecular Analysis


All positive results for HCV antibody detection obtained by rapid testing ($n = 128$; 100%) were confirmed at the laboratory using ELISA test. Viral HCV RNA was detectable in 79.7% ($n = 102$), with viral load ranging from 25 to 80,200,000 UI/mL. In the first step, genotyping by real-time RT PCR was performed for all 102 samples with detectable HCV RNA. This method allowed the genotype identification for 97 samples, while five samples only could not be genotyped. Samples with identified genotypes included genotype 1a ($n = 41$), 1b ($n = 4$), 1 ($n = 5$), 3 ($n = 33$), and 4 ($n = 10$), in addition to four samples with intergenotype mixed infections with genotypes 1a and 3 ($n = 2$) or 1b and 3 ($n = 2$). In the next step, a subset of 64 ($n = 64/102$; 62.7%) samples was selected for sequencing in the NS5b region. This selection included 46 samples with non-identified genotypes and/or subtypes as well as 18 samples with identified genotype and subtype. Among this selection, 62 HCV sequences were obtained for genotype and subtype identification or confirmation. A phylogenetic tree comparing the obtained HCV sequences with reference strains is shown in Figure 2.

Out of a total of 102 samples with detectable HCV RNA, 100 (98%) were successfully genotyped, whereas two (2%) samples could not be genotyped. Among HCV-infected PWID with identified genotypes, the predominant genotype was 1 (52%) followed by genotype 3 (34%) and genotype 4 (10%). Four patients (4%) had an intergenotype mixed infection with genotypes 1 and 3. Overall, six different HCV subtypes had been identified among 94 PWID. The HCV subtype distribution was as follows: subtypes 1a was identified in 50 (53.2%), 1b in 6 (6.4%), 3a in 31 (33.0%), 4a in 3 (3.2%), and 4d in 4 (4.3%) (Figure 3).

DISCUSSION

The present study reports the sociodemographic features, risk behaviors, and the distribution of HCV genotypes in a population of PWID recruited between 2018 and 2019 at a community-based SSP program in the North Eastern region of Tunisia.

One important point to note is that there was a low proportion of women included in the study. This might be explained



partly by the fact that women consuming drugs are less likely to openly report their IDU behavior because of significant social and cultural stigma faced by female PWID across North African countries. The difficulty of recruiting female PWID participants was previously reported in Tunisia (Mosbah et al., 2018), Libya (Mirzoyan et al., 2013), and Morocco (Trimbilas et al., 2014). Additionally, this study shows that HCV-infected PWID attending an SSP program were somewhat younger than larger groups of HCV-infected patients from the general Tunisian population (Bettaieb et al., 2019; Chouikha et al., 2021). The age of the studied PWID was close the age of previously studied populations of injection and non-injection drug users (Sellami et al., 2016; Mosbah et al., 2018).

Furthermore, significant social and economic vulnerability was observed in this study. The majority of PWID had a maximum of primary schooling, were unemployed, and had been previously incarcerated at least once. Low education levels and high unemployment rates were previously reported in PWID groups from other countries (Omland et al., 2013; Edmunds et al., 2019). This social and economic vulnerability may enhance exposure to high-risk environments where harmful behaviors can occur. In fact, high prevalence levels of HCV infection have been reported in prison settings where various risky behaviors, such as use of non-sterile injecting equipment, unsafe tattooing, and unprotected sex, remain important drivers of HCV transmission (Larney et al., 2013; Heijnen et al., 2016). In this light, the

TABLE 1 | Sociodemographic characteristics and risk behaviors of hepatitis C virus (HCV)-positive people who inject drugs (PWID) attending harm reduction centers in the North Eastern region of Tunisia.

		Number (n)	Percentage (%)
Gender	Male	123	96.1
	Female	5	3.9
Age group in years	20–29	8	6.3
	30–39	56	43.8
	40–49	40	31.3
	50–59	22	17.2
	> 60	2	1.6
Education level	Primary	72	56.3
	High school	52	40.6
	University	4	3.1
Civil status	Single	73	57
	Married	33	25.8
	Divorced/widowed	22	17.2
Employment	Employed	44	34.4
	Unemployed	84	65.6
History of incarceration	yes	119	93.0
	no	9	7.0
Housing	Living alone	18	14.1
	Living with family members	110	85.9
Residency in a foreign country	Yes	65	50.8
	No	63	49.2
History of tattooing	Yes	77	60.2
Syringe reuse in the last 12 months	No	51	39.8
	Yes	105	82.0
Syringe sharing in the last 12 months	No	23	18.0
	Yes	56	43.8
Unprotected sexual relations	No	72	56.2
	yes	72	56.2
More than one sexual partner	No	56	43.8
	Yes	36	28.1
	No	92	71.9

access of PWID to adequate prevention and treatment services should be strengthened in prison settings to reduce harmful behaviors and to improve the overall health status of PWID groups (Kamarulzaman et al., 2016; Rich et al., 2016).

In the present work, sharing of injection equipment and syringe reuse during the past 12 months was also frequently reported among our PWID. In reference to the international literature, the main factors associated with transmission and acquisition of HCV infection include sharing of injection equipment, early initiation of IDU, and frequent injecting (Zhou et al., 2019). While this study reported significant levels of risky injection behaviors, little is known about the coverage of existing harm reduction interventions in Tunisia. Although SSPs are provided in Tunisia since 2009 (World Health Organization,

2012), these programs have been implemented on a small scale, in a limited number of community-based centers. Thus, data on intervention coverage is not readily available. There is also a low coverage of interventions for HIV and HCV prevention in the MENA region, with less than four syringes are distributed annually per PWID through harm reduction programs (Larney et al., 2017). Furthermore, access to opioid substitution therapies (OSTs) is not provided in many MENA countries including Tunisia. While the WHO and UNAIDS identified SSP and OST as high-priority interventions for PWID, greater investments in evidence-based interventions for HCV prevention are urgently needed in Tunisia.

In addition to HCV, it is estimated that PWID are highly exposed to other blood-borne viruses such as HIV and HBV. A global systematic review estimated that 13% of PWID are living with HCV–HIV coinfection and that 3% are HCV–HBV coinfecting (Rashti et al., 2020). Results from the present study showed similar trends regarding HCV–HIV and HCV–HBV coinfections. Thus, HIV and HBV prevention and treatment among PWID should be improved through the development of specific recommendations that are specific to HCV-infected PWID. This includes enhancing systematic screening for HBV and HIV coinfections and increasing HBV vaccination rates in non-immunized PWID groups.

From the virological side, patients included in this study were recruited based on HCV testing using rapid tests at the SSP facility level. Positive results for HCV rapid test were confirmed by ELISA testing for all participants. This is in concordance with previous findings showing a high performance of rapid tests in screening of HCV infection especially in high-risk persons (Shivkumar et al., 2012; Poiteau et al., 2016; Barbosa et al., 2017; Mane et al., 2019; Sharafi et al., 2019). Rapid tests represent an interesting alternative for HCV screening among high-risk populations given the advantage of simplicity, quick performance at room temperature, and the limited need of equipment and training for health professionals. Furthermore, point-of-care HCV RNA viral load tests are becoming increasingly available, allowing quick and reliable detection of active HCV infection from venipuncture and finger-stick capillary whole-blood samples without requiring special laboratory infrastructure (Wang et al., 2011; Cooper, 2017; Grebely et al., 2017b; Saludes et al., 2020). In particular, these point-of-care diagnostics for HCV antibodies and HCV RNA are interesting quick and easy testing tools that should be used as part of the national elimination strategy in Tunisia to expand access to HCV diagnosis and treatment among hard-to-reach groups such as PWID. In the present study, 80% had a detectable HCV-RNA, reflecting a high rate of active viral replication within Tunisian PWID. While little is known about rates of chronic HCV infection in Tunisia, especially among high-risk groups, a recent study showed a detectable viral RNA rate of 56.8% in a Center – western city (Thala) in Tunisia (Bettaieb et al., 2019). Higher rates (79%) were also observed among groups of Tunisian hemophiliacs (Djebbi et al., 2008). Furthermore, a systematic review conducted in 2019 highlighted the lack of data for the MENA region and the need to improve country-level estimates for viremic HCV prevalence among PWID (Grebely et al., 2019).




FIGURE 2 | Phylogenetic tree of HCV genotypes identified among HCV-positive PWID attending an SSP in North Eastern region of Tunisia in comparison with 45 reference strains. The tree was performed using the maximum likelihood method and the kimura-2 parameter model. Topology was supported by 1,000 bootstrap replicates. Bootstrap values lower than 50 were not indicated. The sequences reported in this study were identified by the accession numbers.




FIGURE 3 | Distribution of HCV genotypes (A) and subtypes (B) among PWID attending syringe services programs (SSPs) in the North Eastern region of Tunisia.

Generally, high rates of chronic HCV infection are associated with PWID populations because of recurrent exposure to HCV infection (Midgard et al., 2016). HCV chronicity levels among PWID in 12 European countries ranged between 53 and 97% (Wiessing et al., 2014).

Hepatitis C virus genotype distribution among PWID is completely different from that observed in the Tunisian

general population. HCV genotypes 1 and 3 were the most prevalent genotypes (85%) with a predominance of subtype 1a overall. In contrast, previous studies conducted in the general population, either in Tunisia or in some neighboring countries from the Maghreb region, have shown a predominance of subtype 1b followed by subtype 2c (Djebbi et al., 2003, 2004; Mejri et al., 2005; Ezzikouri et al., 2013; Bettaieb et al., 2019;

Chouikha et al., 2021). In this study, there were very few cases of subtype 1b detected among PWID, while genotype 2 was not detected at all. High prevalence of genotypes 1a and 3a among PWID was reported globally (Ruta and Cernescu, 2015; Robaey et al., 2016; Grassi and Ballardini, 2017). This difference in genotype distribution could be explained by the fact that the majority of participants had a history of residency in a foreign country, mainly in Italy and France, where subtypes 1a and 3a were widely detected in PWID groups (Bourlière et al., 2002; Sereno et al., 2009; Trimbilas et al., 2014; Bielen et al., 2017). Thus, it is likely that PWID, with their history of residency in foreign countries, contribute to the introduction of additional HCV subtypes in Tunisia.

We also found in our series that four participants (4%) had an intergenotype mixed infections with genotypes 1 and 3. In fact, PWID are also likely to be exposed to multiple HCV infections due to recurrent risky injection behaviors. The prevalence of intergenotype mixed infections ranged from 14 to 39% (Cunningham et al., 2015).

In addition, sequencing-based methods are increasingly used to identify subtypes when RT-PCR or hybridization tests give non-conclusive results, to identify already new genotypes/subtypes that may emerge and to differentiate between genetically different variants within HCV subtypes. In the present work, sequencing and phylogenetic analysis allowed us to genotype three of the five not typed ones by qRT-PCR, and to determine subtypes within genotypes 1, 3, and 4 by comparison with published reference sequences. The association of multiple techniques for HCV genotyping especially for the high-risk populations could be of great interest to better assess the right genotype distribution and to guide therapeutic approach.

It is noteworthy that national guidelines for the introduction of the DAA regimens in Tunisia were developed in 2015 based on sofosbuvir/ledipasvir as a preferred first-line regimen (Société Tunisienne de Pathologie Infectieuse and Société Tunisienne de Gastro-entérologie., 2015). This is a non-pangenotypic regimen, which is commonly used in countries where only a single HCV genotype like genotype 1 is widely isolated (Zoratti et al., 2020). In the current DAA era, HCV genotype 3 infection has emerged as relatively difficult to treat when compared with other HCV genotypes, causing lower sustained virologic response rates and longer treatment duration (Ampuero et al., 2014). Patients with HCV genotype 3 infection had an accelerated fibrosis progression and higher incidence of hepatocellular carcinoma (Bochud et al., 2009; Nkontchou et al., 2011; Kanwal et al., 2014; McMahon et al., 2017). As HCV genotyping remains a key component of pretherapeutic assessment prior to treatment initiation, national guidelines recommend treating HCV genotype 3 infection with sofosbuvir-ribavirin or with sofosbuvir/daclatasvir for 24 weeks among patients without cirrhosis and with sofosbuvir/daclatasvir-ribavirin for 24 weeks among patients with compensated or decompensated cirrhosis even though daclatasvir is not available in Tunisia. The WHO guidelines for HCV treatment were updated twice since 2016 to include pangenotypic DAA regimens reducing the need for genotyping to guide treatment decisions

and recommending sofosbuvir/daclatasvir combination for genotype 3 infection during 12 weeks for patients without cirrhosis and 24 weeks for patients with compensated cirrhosis (World Health Organization, 2018a). Current European recommendations advice for the use of pangenotypic regimens of sofosbuvir/velpatasvir for 12 weeks or glecaprevir/pibrentasvir for 8 weeks for patients infected with HCV genotype 3 and without cirrhosis, and with glecaprevir/pibrentasvir for 12 weeks for patients with compensated cirrhosis (European Association for the Study of the Liver, 2018). Due to the potential suboptimal treatment outcomes associated with currently available DAA regimens, in addition to the high rate of genotype 3 reported in the present study, the introduction of pangenotypic treatments, such as sofosbuvir–daclatasvir, sofosbuvir/velpatasvir, and glecaprevir/pibrentasvir, would be highly beneficial to maximize access to more efficacious and shorter HCV treatments in Tunisia especially for PWID with HCV genotype 3 infection.

While this study illuminates significant knowledge gaps regarding HCV genotype distribution among PWID, these findings cannot be generalized to the overall population of HCV-infected PWID in Tunisia for several reasons. First, the study was limited to the North Eastern geographic region of the country, and some subgroups of PWID such as youth and women were not adequately represented. In the future, this limitation could be addressed by initiating research partnerships with community-based organizations that provide gender-sensitive harm reduction services, which can be also respondent to adolescents and young people's specific needs with reduced stigma and discrimination toward drug users. In addition, the study is potentially subject to selection bias because included participants were exclusively recruited within a local SSP where participants were more likely to have benefited from diverse prevention and treatment interventions. On the other hand, interview responses might be affected by information bias since IDU is severely stigmatized and criminalized in Tunisia. To reduce stigma and discrimination toward PWID and minimize information bias, all interviews were administrated by trained medical doctors in close cooperation with peer educators and community representatives. In light of these different strengths and limitations, this study provides the first exploration about HCV genotype diversity among PWID in Tunisia. In conclusion, findings from this study provided the first exploration about the distribution of HCV genotypes among a population of Tunisian PWID, which can be used to improve national efforts aiming to achieve national HCV elimination goal. Given the predominance of HCV subtypes 1a and 3a among PWID, this distribution of HCV genotypes was distinct from the general population. While further genotype surveillance and monitoring studies among vulnerable populations are highly needed to improve the current knowledge of HCV epidemiology in Tunisia, community-based prevention and harm reduction efforts for PWID should be strengthened and expanded, and current national guidelines for HCV treatment should be updated to include the use of new pangenotypic DAA regimens and to ensure access to better treatment outcomes especially for patients infected with HCV genotype 3.

DATA AVAILABILITY STATEMENT

The datasets presented in this study may be found in online repositories. The names of the repositories and accession numbers can be found in the article.

ETHICS STATEMENT

This study was conducted in accordance with the Declaration of Helsinki and the study protocol was reviewed and approved by the Biomedical Ethics Committee of the Institute Pasteur de Tunis (reference number: 2018/04/I/LVCIPT/V2). All participants provided their written informed consent before enrolment in the study.

AUTHOR CONTRIBUTIONS

AC and AGhr contributed to the conception and design of the study, and wrote the first draft of the manuscript. AGho, WH, MK, AS, and HTo contributed in the investigation. HH, SM, CB, and SA contributed in the data curation and resources. SB and MG contributed in the data curation and investigation.

REFERENCES

- Alashek, W. A., and Altagdi, M. (2008). Risk factors and genotypes of hepatitis C virus infection in libyan patients. *Libyan J. Med.* 3, 162–165. doi: 10.4176/080425
- Ampuero, J., Romero-Gómez, M., and Reddy, K. R. (2014). Review article: HCV Genotype 3 – the new treatment challenge. *Aliment. Pharmacol. Ther.* 39, 686–698. doi: 10.1111/apt.12646
- Ara, A. K., and Paul, J. P. (2015). New direct-acting antiviral therapies for treatment of chronic hepatitis C virus infection. *Gastroenterol. Hepatol.* 11, 458–466.
- Barbosa, J. R., Colares, J. K. B., Flores, G. L., Cortes, V. F., Miguel, J. C., and Portilho, M. M. (2017). Performance of rapid diagnostic tests for detection of Hepatitis B and C markers in HIV infected patients. *J. Virol. Methods* 248, 244–249. doi: 10.1016/j.jvromet.2017.08.001
- Bettaieb, J., Chouikha, A., Khedhiri, M., Kharroubi, G., Badreddine, M., Bel Hadj, et al. (2019). Hepatitis C virus epidemiology in central-west tunisia: a population-based cross-sectional study. *Arch. Virol.* 164, 2243–2253. doi: 10.1007/s00705-019-04308-8
- Bielen, R., Moreno, C., Van Vlierberghe, H., Bourgeois, S., Mulkay, J. P., and Vanwolleghem, T. (2017). Belgian experience with direct acting antivirals in people who inject drugs. *Drug Alcohol. Depend.* 177, 214–220. doi: 10.1016/j.drugalcdep.2017.04.003
- Bochud, P. Y., Cai, T., Overbeck, K., Bochud, M., Dufour, J. F., and Müllhaupt, B. (2009). Genotype 3 is associated with accelerated fibrosis progression in chronic Hepatitis C. *J. Hepatol.* 51, 655–666. doi: 10.1016/j.jhep.2009.05.016
- Bourlière, M., Barberin, J. M., Rotily, M., Guagliardo, V., Portal, I., and Lecomte, L. (2002). Epidemiological changes in hepatitis C virus genotypes in france: evidence in intravenous drug users. *J. Viral Hepat.* 9, 62–70. doi: 10.1046/j.1365-2893.2002.00319.x
- Chaabna, K., Cheema, S., Abraham, A., Alrouh, H., Lowenfels, A. B., and Maisonneuve, P. (2018). Systematic overview of Hepatitis C infection in the middle East and North Africa. *World J. Gastroenterol.* 24, 3038–3054. doi: 10.3748/wjg.v24.i27.3038
- Chouikha, A., Khedhiri, M., Triki, H., Hammemi, W., Sadraoui, A., and Touzi, H. (2021). Focus on Hepatitis C virus genotype distribution in tunisia prior to elimination: a 16-year retrospective study. *Arch. Virol.* 166, 501–510. doi: 10.1007/s00705-020-04918-7

HTr contributed in the reviewing, editing the manuscript, and supervising the work. All authors contributed to the manuscript revision, and read and approved the submitted version.

FUNDING

This study received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No. 741677. It was also jointly supported by the Tunisian Association for Information on HIV/AIDS and Toxicomania (ATIOST), the Ministry of Higher Education (Research Laboratory LR20IPT02: "Virus, Vectors and Hosts: One Health approach and technological innovation for a better health") and the Clinical Investigation Center (CIC).

ACKNOWLEDGMENTS

The authors gratefully acknowledge the efforts made by all participants who took part in this study in addition to all peer educators and staff members from both the Tunisian Association for Information on HIV/AIDS and Toxicomania (ATIOST) and "la boutique des sciences" of the Institut Pasteur de Tunis (IPT).

- Cooper, C. (2017). Rapid HCV RNA testing: removing the final obstacle to elimination. *Lancet Gastroenterol. Hepatol.* 2, 468–469. doi: 10.1016/s2468-1253(17)30086-9
- Cunningham, E. B., Applegate, T. L., Lloyd, A. R., Dore, G. J., and Grebely, J. (2015). Mixed HCV infection and reinfection in people who inject drugs—impact on therapy. *Nat. Rev. Gastroenterol. Hepatol.* 12, 218–230. doi: 10.1038/nrgastro.2015.36
- Degenhardt, L., Peacock, A., Colledge, S., Leung, J., Grebely, J., and Vickerman, P. (2017). Global prevalence of injecting drug use and sociodemographic characteristics and prevalence of HIV, HBV, and HCV in people who inject drugs: a multistage systematic review. *Lancet Global Health* 5, e1192–e1207.
- Djebbi, A., Bahri, O., Langar, H., Sadraoui, A., Mejri, S., and Triki, H. (2008). Genetic variability of Genotype 1 Hepatitis C virus isolates from Tunisian haemophiliacs. *New Microbiol.* 31, 473–480.
- Djebbi, A., Mejri, S., Thiers, V., and Triki, H. (2004). Phylogenetic analysis of Hepatitis C virus isolates from Tunisian patients. *Eur. J. Epidemiol.* 19, 555–562. doi: 10.1023/b:ejep.0000032348.83087.01
- Djebbi, A., Triki, H., Bahri, O., Cheikh, I., Sadraoui, A., Ben Ammar, A., et al. (2003). Genotypes of Hepatitis C virus circulating in Tunisia. *Epidemiol. Infect.* 130, 501–505. doi: 10.1017/s095026880300846x
- Edmunds, B. L., Miller, E. R., and Tsourtos, G. (2019). The distribution and socioeconomic burden of Hepatitis C Virus in South Australia: a cross-sectional study 2010–2016. *BMC Public Health* 19:527.
- European Association for the Study of the Liver (2018). EASL Recommendations on treatment of Hepatitis C 2018. *J. Hepatol.* 69, 461–511. doi: 10.1016/j.jhep.2018.03.026
- Ezzikouri, S., Pineau, P., and Benjelloun, S. (2013). Hepatitis C virus Infection in the maghreb region. *J. Med. Virol.* 85, 1542–1549. doi: 10.1002/jmv.23643
- Fadlalla, F. A., Mohamoud, Y. A., Mumtaz, G. R., and Abu-Raddad, L. J. (2015). The epidemiology of Hepatitis C Virus in the maghreb region: systematic review and meta-analyses. *PLoS One* 10:e0121873. doi: 10.1371/journal.pone.0121873
- Gower, E., Estes, C., Blach, S., Razavi-Shearer, K., and Razavi, H. (2014). Global epidemiology and genotype distribution of the Hepatitis C virus infection. *J. Hepatol.* 61(Suppl.), S45–S57.
- Grassi, A., and Ballardini, G. (2017). Hepatitis C in injection drug users: it is time to treat. *World J. Gastroenterol.* 23, 3569–3571. doi: 10.3748/wjg.v23.i20.3569

- Grebely, J., Dore, G. J., Morin, S., Rockstroh, J. K., and Klein, M. B. (2017a). Elimination of HCV as a public health concern among people who inject drugs by 2030 - what will it take to get there? *J. Int. AIDS Soc.* 20:22146. doi: 10.7448/IAS.20.1.22146
- Grebely, J., Lamoury, F. M. J., Hajarizadeh, B., Mowat, Y., Marshall, A. D., and Bajis, S. (2017b). Evaluation of the Xpert HCV viral load point-of-care assay from venepuncture-collected and finger-stick capillary whole-blood samples: a cohort study. *Lancet Gastroenterol. Hepatol.* 2, 514–520. doi: 10.1016/s2468-1253(17)30075-4
- Grebely, J., Larney, S., Peacock, A., Colledge, S., Leung, J., and Hickman, M. (2019). Global, regional, and country-level estimates of Hepatitis C infection among people who have recently injected drugs. *Addiction* 114, 150–166. doi: 10.1111/add.14393
- Heijnen, M., Mumtaz, G. R., and Abu-Raddad, L. J. (2016). Status of HIV and Hepatitis C virus infections among prisoners in the middle East and North Africa: review and synthesis. *J. Int. AIDS Soc.* 19:20873. doi: 10.7448/ias.19.1.20873
- Kamarulzaman, A., Reid, S. E., Schwitters, A., Wiessing, L., El-Bassel, N., and Dolan, K. (2016). Prevention of transmission of HIV, Hepatitis B Virus, Hepatitis C Virus, and Tuberculosis in prisoners. *Lancet* 388, 1115–1126. doi: 10.1016/s0140-6736(16)30769-3
- Kanwal, F., Kramer, J. R., Ilyas, J., Duan, Z., and El-Serag, H. B. (2014). HCV Genotype 3 is associated with an increased risk of cirrhosis and hepatocellular cancer in a national sample of U.S. Veterans with HCV. *Hepatology* 60, 98–105. doi: 10.1002/hep.27095
- Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: molecular evolutionary genetics analysis Version 7.0 for bigger datasets. *Mol. Biol. Evol.* 33, 1870–1874. doi: 10.1093/molbev/msw054
- Larney, S., Kopinski, H., Beckwith, C. G., Zaller, N. D., Jarlais, D. D., and Hagan, H. (2013). Incidence and prevalence of Hepatitis C in prisons and other closed settings: results of a systematic review and meta-analysis. *Hepatology* 58, 1215–1224. doi: 10.1002/hep.26387
- Larney, S., Peacock, A., Leung, J., Colledge, S., Hickman, M., and Vickerman, P. (2017). Global, regional, and country-level coverage of interventions to prevent and manage HIV and Hepatitis C among people who inject drugs: a systematic review. *Lancet Global Health* 5, e1208–e1220.
- MacArthur, G. J., van Velzen, E., Palmateer, N., Kimber, J., Pharris, A., and Hope, V. (2014). Interventions to prevent HIV and Hepatitis C in people who inject drugs: a review of reviews to assess evidence of effectiveness. *Int. J. Drug Policy* 25, 34–52. doi: 10.1016/j.drugpo.2013.07.001
- Mane, A., Sacks, J., Sharma, S., Singh, H., Tejada-Strop, A., and Kamili, S. (2019). Evaluation of five rapid diagnostic tests for detection of antibodies to Hepatitis C Virus (HCV): a step towards scale-up of HCV screening efforts in India. *PLoS One* 14:e0210556. doi: 10.1371/journal.pone.0210556
- Martin, N. K., Hickman, M., Hutchinson, S. J., Goldberg, D. J., and Vickerman, P. (2013). Combination interventions to prevent HCV transmission among people who inject drugs: modeling the impact of antiviral treatment, needle and syringe programs, and opiate substitution therapy. *Clin. Infect. Dis.* 57(Suppl. 2), S39–S45.
- McMahon, B. J., Bruden, D., Townshend-Bulson, L., Simons, B., Spradling, P., and Livingston, S. (2017). Infection with Hepatitis C Virus Genotype 3 is an independent risk factor for end-stage liver disease, Hepatocellular Carcinoma, and liver-related death. *Clin. Gastroenterol. Hepatol.* 15, 431–437.e2.
- Mejri, S., Ben Salah, A., Triki, H., Ben Alaya, N., Djebbi, A., and Dellagi, K. (2005). Contrasting patterns of Hepatitis C Virus infection in two regions from Tunisia. *J. Med. Virol.* 76, 185–193. doi: 10.1002/jmv.20342
- Midgard, H., Weir, A., Palmateer, N., Lo, R., Pineda, J. A., et al. (2016). HCV epidemiology in high-risk groups and the risk of reinfection. *J. Hepatol.* 65(Suppl.), S33–S45.
- Mirzoyan, L., Berendes, S., Jeffery, C., Thomson, J., Ben Othman, H., and Danon, L. (2013). New evidence on the HIV epidemic in Libya: why countries must implement prevention programs among people who inject drugs. *J. Acquir. Immune Defic. Syndr.* 62, 577–583. doi: 10.1097/qai.0b013e318284714a
- Mosbah, H. B., Ghorbel, A., and Hedhili, A. (2018). Illicit use of high-dosage buprenorphine in Tunisia. *Afr. J. Drug Alcohol Stud.* 17, 39–51.
- Nkontchou, G., Ziou, M., Aout, M., Lhabadie, M., Baazia, Y., Mahmoudi, A., et al. (2011). HCV genotype 3 is associated with a higher hepatocellular carcinoma incidence in patients with ongoing viral C cirrhosis. *J. Viral. Hepat.* 18, e516–e522. doi: 10.1111/j.1365-2893.2011.01441.x
- Omland, L. H., Osler, M., Jepsen, P., Krarup, H., Weis, N., and Christensen, P. B. (2013). Socioeconomic status in HCV infected patients - risk and prognosis. *Clin. Epidemiol.* 5, 163–172. doi: 10.2147/cep.s43926
- Palmateer, N., Kimber, J., Hickman, M., Hutchinson, S., Rhodes, T., Goldberg, D. (2010). Evidence for the effectiveness of sterile injecting equipment provision in preventing hepatitis C and human immunodeficiency virus transmission among injecting drug users: a review of reviews. *Addiction*. 105, 844–859. doi: 10.1111/j.1360-0443.2009.02888.x
- Poiteau, L., Soulier, A., Rosa, I., Roudot-Thoraval, F., Hézode, C., and Pawlotsky, J. M. (2016). Performance of rapid diagnostic tests for the detection of antibodies to Hepatitis C Virus in whole blood collected on dried blood spots. *J. Viral. Hepat.* 23, 399–401. doi: 10.1111/jvh.12501
- Rashti, R., Sharafi, H., Alavian, S. M., Moradi, Y., Mohamadi Bolbanabad, A., and Moradi, G. (2020). Systematic review and meta-analysis of global prevalence of HBsAg and HIV and HCV antibodies among people who inject drugs and female sex workers. *Pathogens* 9:432. doi: 10.3390/pathogens9060432
- Rich, J. D., Beckwith, C. G., Macmadu, A., Marshall, B. D. L., and Brinkley-Rubinstein, L. (2016). Clinical care of incarcerated people with HIV, viral Hepatitis, or Tuberculosis. *Lancet* 388, 1103–1114. doi: 10.1016/s0140-6736(16)30379-8
- Robaey, G., Bielen, R., Azar, D. G., Razavi, H., and Nevens, F. (2016). Global Genotype Distribution of Hepatitis C Viral Infection among People Who Inject Drugs. *J. Hepatol.* 65, 1094–1103. doi: 10.1016/j.jhep.2016.07.042
- Ruta, S., and Cernescu, C. (2015). Injecting drug use: a vector for the introduction of new Hepatitis C virus genotypes. *World J. Gastroenterol.* 21, 10811–10823. doi: 10.3748/wjg.v21.i38.10811
- Saludes, V., Antuori, A., Lazarus, J. V., Folch, C., González-Gómez, S., and González, N. (2020). Evaluation of the Xpert HCV VL fingerstick point-of-care assay and dried blood spot HCV-RNA testing as simplified diagnostic strategies among people who inject drugs in Catalonia, Spain. *Int. J. Drug Policy* 80:102734. doi: 10.1016/j.drugpo.2020.102734
- Sellami, R., Feki, I., Zahaf, A., and Masmoudi, J. (2016). The profile of drug users in Tunisia: implications for prevention. *La Tunisie Médicale* 94, 531–534.
- Sereno, S., Perinelli, P., and Laghi, V. (2009). Changes in the prevalence of Hepatitis C virus genotype among Italian injection drug users—relation to period of injection started. *J. Clin. Virol.* 45, 354–357. doi: 10.1016/j.jcv.2009.04.022
- Sharafi, H., Poustchi, H., Azimian, F., Tamadoni, B., Ramezani, R., and Gouya, M. M. (2019). Performance of a rapid diagnostic test for screening of Hepatitis C in a real-life prison setting. *J. Clin. Virol.* 113, 20–23. doi: 10.1016/j.jcv.2019.02.005
- Shivkumar, S., Peeling, R., Jafari, Y., Joseph, L., and Pant Pai, N. (2012). Accuracy of rapid and point-of-care screening tests for Hepatitis C: a systematic review and meta-analysis. *Ann. Intern. Med.* 157, 558–566. doi: 10.7326/0003-4819-157-8-201210160-00006
- Société Tunisienne de Pathologie Infectieuse and Société Tunisienne de Gastro-entérologie. (2015). *Actualisation des Recommandations du Traitement de l'hépatite Virale C en Collaboration Entre la STGE et la STPI*. Available online at: https://infectiologie.org.tn/pdf_ppt_docs/recommandations/1536342624.pdf
- Trimbilas, R.-D., Serghini, F. Z., Lazaar, F., Baha, W., Foullous, A., and Essalhi, M. (2014). The 'hidden' epidemic: a snapshot of Moroccan intravenous drug users. *Virol. J.* 11:43. doi: 10.1186/1743-422x-11-43
- Wang, Q. Q., Zhang, J., Hu, J. S., Chen, H. T., Du, L., and Wu, L. Q. (2011). Rapid detection of Hepatitis C Virus RNA by a reverse transcription loop-mediated isothermal amplification assay. *FEMS Immunol. Med. Microbiol.* 63, 144–147.
- Wiessing, L., Ferri, M., Grady, B., Kantzanou, M., Sperle, I., and Cullen, K. J. (2014). Hepatitis C virus infection epidemiology among people who inject drugs in Europe: a systematic review of data for scaling up treatment and prevention. *PLoS One* 9:e103345. doi: 10.1371/journal.pone.0103345
- World Health Organization (2012). *MENAHRA—The Middle East and North Africa Harm Reduction Association Best Practices in Strengthening Civil Society's Role in Delivering Harm Reduction Services*. Geneva: WHO.

- World Health Organization (2016). *Global Health Sector Strategy on Viral Hepatitis 2016-2021. Towards Ending Viral Hepatitis*. Geneva: World Health Organization.
- World Health Organization (2017). *Global Hepatitis Report 2017*. Geneva: World Health Organization.
- World Health Organization (2018a). *Guidelines for the Care and Treatment of Persons Diagnosed with Chronic Hepatitis C Virus Infection*. Geneva: WHO.
- World Health Organization (2018b). *Progress Report on Access to Hepatitis C Treatment: Focus on Overcoming Barriers in Low-and Middle-Income Countries*. Geneva: World Health Organization.
- Zhou, B., Cai, G. F., Lv, H. K., Xu, S. F., Wang, Z. T., and Jiang, Z. G. (2019). Factors correlating to the development of Hepatitis C virus infection among drug users—findings from a systematic review and meta-analysis. *Int. J. Environ. Res. Public Health* 16:2345. doi: 10.3390/ijerph16132345
- Zoratti, M. J., Siddiqua, A., Morassut, R. E., Zeraatkar, D., Chou, R., and van Holten, J. (2020). Pangenotypic direct acting antivirals for the treatment of chronic Hepatitis C virus infection: a systematic literature review and meta-analysis. *EClinicalMedicine* 18:100237. doi: 10.1016/j.eclinm.2019.12.007
- Conflict of Interest:** The authors declare that Prochidia Laboratory provided genotyping kits for free to the Laboratory of Clinical Virology to conduct the present work.
- Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Chouikha, Ghrabi, Ghodbane, Hammemi, Khedhiri, Sadraoui, Touzi, Hassine, Maatoug, Bensaoud, Abdelhak, Bouarrouj, Gdoura, Chaouachi and Triki. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.