Check for updates

OPEN ACCESS

EDITED AND REVIEWED BY Rustam Aminov, University of Aberdeen, United Kingdom

*CORRESPONDENCE María Guembe ⊠ mariaguembe@hotmail.com

RECEIVED 09 January 2024 ACCEPTED 24 January 2024 PUBLISHED 02 February 2024

CITATION

Benjumea A, Diaz-Navarro M, Hafian R, Cercenado E, Sánchez-Somolinos M, Vaquero J, Chana F, Muñoz P and Guembe M (2024) Corrigendum: Tranexamic acid in combination with vancomycin or gentamicin has a synergistic effect against staphylococci. *Front. Microbiol.* 15:1355087. doi: 10.3389/fmicb.2024.1355087

COPYRIGHT

© 2024 Benjumea, Díaz-Navarro, Hafian, Cercenado, Sánchez-Somolinos, Vaquero, Chana, Muñoz and Guembe. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Tranexamic acid in combination with vancomycin or gentamicin has a synergistic effect against staphylococci

Antonio Benjumea¹, Marta Díaz-Navarro^{2,3}, Rama Hafian⁴, Emilia Cercenado^{2,5}, Mar Sánchez-Somolinos², Javier Vaquero^{1,3,6}, Francisco Chana¹, Patricia Muñoz^{2,3,5,6} and María Guembe^{2,3*}

¹Department of Orthopaedic Surgery and Traumatology, Hospital General Universitario Gregorio Marañón, Madrid, Spain, ²Department of Clinical Microbiology and Infectious Diseases, Hospital General Universitario Gregorio Marañón, Madrid, Spain, ³Instituto de Investigación Sanitaria Gregorio Marañón, Madrid, Spain, ⁴School of Biology, Universidad Complutense de Madrid, Madrid, Spain, ⁵CIBER Enfermedades Respiratorias-CIBERES (CB06/06/0058), Madrid, Spain, ⁶School of Medicine, Universidad Complutense de Madrid, Madrid, Spain

KEYWORDS

tranexamic acid, bacterial growth, antimicrobial effect, synergy, in vitro model

A corrigendum on

Tranexamic acid in combination with vancomycin or gentamicin has a synergistic effect against staphylococci

by Benjumea, A., Díaz-Navarro, M., Hafian, R., Cercenado, E., Sánchez-Somolinos, M., Vaquero, J., Chana, F., Muñoz, P., and Guembe, M. (2022). *Front. Microbiol.* 13:935646. doi: 10.3389/fmicb.2022.935646

In the published article, there were errors in Table 2 as published. The *S. aureus* MIC_{50} fold reductions for gentamicin+TXA 50 mg/ml appeared as 5. This has been corrected to 1. The *S. epidermidis* MIC_{50} fold reductions for gentamicin+TXA 10mg/ml appeared as 9. This has been corrected to 8. The corrected Table 2 and its caption appear below.

The authors apologize for these errors and state that they do not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Treatment	<i>S. aureus</i> MIC ₅₀ (ranges) mg/L	MIC ₅₀ fold reductions (no.)	S. epidermidis MIC ₅₀ (ranges) mg/L	MIC ₅₀ fold reductions (no.)	Overall MIC ₅₀ (ranges) mg/L	MIC ₅₀ fold reductions (no.)
Vancomycin	4 (0.12–16)		16 (16–16)		8 (0.12–16)	
Vancomycin+TXA 10 mg/ml	0.03 (0.03-0.06)	8	0.03 (0.03-0.03)	8	0.03 (0.03-0.06)	8
Vancomycin+TXA 50 mg/ml	0.12 (0.03-0.25)	1	8 (0.03–16)	1	4 (0.03–16)	1
Gentamicin	16 (0.06-32)		32 (32–32)		32 (0.06-32)	
Gentamicin+TXA 10 mg/ml	0.06 (0.06-0.12)	7	0.12 (0.06-0.25)	8	0.12 (0.06-0.25)	8
Gentamicin+TXA 50 mg/ml	8 (0.06-32)	1	16 (0.06–32)	1	16 (0.06–32)	1

TABLE 2 MIC₅₀ for vancomycin and gentamicin alone and in combination with TXA using the automatized turbidity method.

TXA, tranexamic acid; MIC, minimal inhibitory concentration.