



# Polymorphism of Alcohol Metabolizing Gene ADH3 Predisposes to Development of Alcoholic Pancreatitis in North Indian Population

OPEN ACCESS

**Edited by:**

Arun Bhardwaj,  
University of South Alabama Mitchell  
Cancer Institute, USA

**Reviewed by:**

Sandeep Kumar,  
State University of New York College  
of Optometry, USA  
Surendra Kumar Shukla,  
University of Nebraska Medical  
Center, USA  
Vinay Kumar Mittal,  
Thermo Fisher Scientific, USA  
Shinjini Singh,  
University of Texas MD Anderson  
Cancer Center, USA

**\*Correspondence:**

Ghanshyam Upadhyay  
upadhyayitr@gmail.com;  
Gourdas Choudhuri  
choudhuri.gour@gmail.com

**Specialty section:**

This article was submitted to  
Molecular Diagnostics,  
a section of the journal  
Frontiers in Molecular Biosciences

**Received:** 01 September 2015

**Accepted:** 13 November 2015

**Published:** 16 December 2015

**Citation:**

Singh D, Negi TS, Upadhyay G and  
Choudhuri G (2015) Polymorphism of  
Alcohol Metabolizing Gene ADH3  
Predisposes to Development of  
Alcoholic Pancreatitis in North Indian  
Population. *Front. Mol. Biosci.* 2:67.  
doi: 10.3389/fmolb.2015.00067

Divya Singh<sup>1,2</sup>, Tajwar S. Negi<sup>2</sup>, Ghanshyam Upadhyay<sup>1\*</sup> and Gourdas Choudhuri<sup>2,3\*</sup>

<sup>1</sup> Department of Biology, City College of New York, New York, NY, USA, <sup>2</sup> Department of Gastroenterology, Sanjay Gandhi Post Graduate Institute of Medical Sciences, Lucknow, India, <sup>3</sup> Department of Gastroenterology and Hepatobiliary Sciences, Fortis Memorial Research Institute, Gurgaon, India

**Background and aim:** Genetic factors regulating alcohol metabolism could predispose in developing alcoholic pancreatitis (ACP). Studies revealed that alcohol could be metabolized by both ways, oxidative and non-oxidative. The main oxidative pathway includes alcohol dehydrogenase (ADH), aldehyde dehydrogenase (ALDH), and cytochrome P450 enzyme. We investigated the association of polymorphisms in these enzymes with the alcoholic pancreatitis in the north Indian population.

**Method:** Patients with alcoholic pancreatitis (ACP;  $n = 72$ ), tropical calcific pancreatitis (TCP;  $n = 75$ ), alcoholic controls (AC;  $n = 40$ ), and healthy controls (HC;  $n = 100$ ) were included in the study. Blood samples were collected from the subjects in EDTA coated vials. DNA was extracted and genotyping for ADH3, ALDH2, and CYP2E1 was done by PCR-RFLP (polymerase chain reaction—restriction fragment length polymorphism). The products were analyzed by gel electrophoresis.

**Result:** The frequency distribution of ADH3\*1/\*1 genotype was significantly higher in ACP group (59.7%) compared with TCP (38.7%), HC (42%), and AC (37.5%) and was found to be associated with increased risk of alcoholic pancreatitis. There was no statistically significant difference between the frequency distribution of ADH3\*1/\*1, ADH3\*1/\*2, and ADH3\*2/\*2 genotypes between TCP and HC or healthy alcoholics. ALDH2 gene was monomorphic in our population, and the frequencies for CYP2E1 intron 6 Dra I polymorphism were comparable in all the four groups.

**Conclusion:** This study shows that carriers of ADH3\*1/\*1 individuals consuming alcohol are at higher risk for alcoholic pancreatitis than those with other genotypes such as ADH3\*1/\*2 and ADH3\*2/\*2.

**Keywords:** chronic pancreatitis, genetic polymorphism, alcohol dehydrogenase

## INTRODUCTION

Alcohol tolerance is highly dependent on the presence or absence of alcohol metabolizing enzymes and the subjects having history of alcohol consumption are more susceptible for chronic pancreatitis onset (Czech and Hartleb, 2003; Cichoz-Lach et al., 2008). Alcohol consumption is suggested to be a major cause of chronic pancreatitis in 70–80% cases in developed countries (Singh and Simsek, 1990; Dzienizewski and Gabryelewicz, 2003; Cichoz-Lach et al., 2008). Genetic factors regulating alcohol metabolism could play a role in developing alcoholic pancreatitis (Vonlaufen et al., 2007). Although these factors are critical in the etiology of chronic pancreatitis, their role in alcoholic pancreatitis is not clearly elucidated.

Alcohol is mainly oxidized by oxidative pathway. The major enzyme of oxidative pathway of alcohol metabolism is alcohol dehydrogenase (ADH)- a dimeric Zn-containing protein with a subunit molecular weight of 40 kDa, and aldehyde dehydrogenase (ALDH; Quertemont, 2004; Edenberg, 2007). ADH is encoded by at least seven gene loci located on the long arm of chromosome 4 (chromosome 4q22) in which ADH2 (ADH1B) and ADH3 (ADH1C) are highly polymorphic (Edenberg, 2000, 2007). ADH3 gene has two alleles ADH3\*1, ADH3\*2 encoding subunits  $\gamma^1$  (Arginine at position 272 and isoleucine at position 350), and  $\gamma^2$  (glutamine at position 272 and a valine at position 350), respectively (Edenberg, 2007). The subunit  $\gamma^1$  shows higher ethanol activity than the subunit  $\gamma^2$  (Day et al., 1991; Crabb et al., 1993; Edenberg, 2007). In majority of cases the occurrence of these two SNPs have been observed, indicating very high linkage disequilibrium (Edenberg, 2007). The turnover rate of ADH depends on the association of the subunits and the homodimeric enzyme  $\gamma^1\gamma^1$  have substantially higher rate (70%) than  $\gamma^2\gamma^2$  enzyme (Edenberg, 2007). Additionally, ADH3, which encodes a subunit with threonine at position 352 (*ADH1C\*Thr352*) has been described in the Native Americans, however, the protein have not been properly investigated (Osier et al., 2002; Edenberg, 2007).

Alcohol is first metabolized to acetaldehyde by ADH and then to acetate by ALDH. Four different isozymes of ALDH; ALDH1, ALDH2, ALDH3, and ALDH4 are found predominantly in the liver and can be differentiated with respect to kinetic properties, localization within the cell, tissue distribution, and electrophoretic mobility (Greenfield and Pietruszko, 1977; Harada et al., 1980; Duley et al., 1985). ALDH2 isozyme is located in mitochondria and is the principal enzyme for acetaldehyde metabolism. ALDH2 is located on the chromosome 12 (chromosome 12q24.2) and is polymorphic with two alleles: ALDH2\*1 and ALDH2\*2 with ethnic differences in the polymorphisms and different kinetic properties (Day et al., 1991; Edenberg, 2007).

Moreover, the involvement of Cytochrome P450s (CYPs) has been also implicated in the alcohol metabolism and associated disease progression (Burim et al., 2004; Liu et al., 2007). CYP2E1 is a key microsomal enzyme that metabolizes alcohol in the non-alcohol dehydrogenase pathway (Ingelman-Sundberg et al., 1993; Song, 1996; Liu et al., 2007). Since, CYP2E1 is ethanol inducible and highly polymorphic, it is inevitable to investigate

polymorphism in this gene, which may explain the variation in disease causation by alcohol. One important polymorphism of CYP2E1 in the intron 6 region, Dra I polymorphism, has been found to be associated with chronic pancreatitis (Ingelman-Sundberg et al., 1993; Song, 1996).

Polymorphisms in alcohol metabolizing enzyme genes have been found to be associated with alcoholic liver disease (Khan et al., 2009, 2010); however, not much information regarding the role of alcohol metabolizing genes polymorphisms in pancreatitis in the north Indian population is available. Evidence from epidemiological, laboratory, and clinical studies suggest that only about 5–10% alcoholics suffer from clinically recognized CP (Bisceglie and Segal, 1984; Harber et al., 1995; Perri et al., 2003). This variation in disease causation suggests that there could be differences in underlying genetic susceptibility. It can be speculated that polymorphism in these alcohol metabolizing enzyme genes could predispose subjects consuming alcohol to alcoholic pancreatitis. Therefore, the aim of the present study is to investigate the association of ADH3, ALDH2, and CYP2E1 polymorphisms in alcoholic pancreatitis in North Indian population.

## SUBJECTS AND METHODS

### Subjects

All consecutive patients with pancreatitis attending the Pancreatobiliary Clinic at Sanjay Gandhi Post Graduate Institute of Medical Sciences, Lucknow were included in the study as per inclusion criteria. Diagnosis was based on clinical and radiological criteria. Inclusion criteria for tropical pancreatitis were radiological evidence of pancreatic intraductal calculi, no history of alcohol consumption and no family history of Pancreatitis. However, inclusion criteria for alcoholic pancreatitis were history of >80 g of alcohol ingestion for at least 3 years, radiological evidence of pancreatitis, increased serum amylase, and absence of any other cause for pain (Table 1). Alcohol consumption history was taken in a questionnaire. Almost all the patients in both the group (ACP and TCP) had pain at presentation. The pain was epigastric in nature usually radiating

**TABLE 1 | The inclusion and exclusion criteria for the recruitment of subjects for the study.**

	Inclusion criteria	Exclusion criteria
ACP	<ol style="list-style-type: none"> <li>1. History of &gt;80 g of alcohol intake for at least 3 years</li> <li>2. Radiological evidence of pancreatitis (acute or chronic)</li> <li>3. Increased serum amylase</li> <li>4. Absence of any other cause for pain</li> </ol>	<ol style="list-style-type: none"> <li>1. Presence of pancreatic cancer</li> <li>2. Occasional alcohol intake</li> </ol>
TCP	<ol style="list-style-type: none"> <li>1. Radiological evidence of pancreatic intraductal calcification</li> <li>2. Non-alcoholic</li> <li>3. No family history of Pancreatitis</li> </ol>	<ol style="list-style-type: none"> <li>1. Presence of any other etiological factor like alcoholism, gall stones hypercalcaemia etc.</li> <li>2. Presence of pancreatic cancer</li> </ol>

to back. Frequency of diabetes in TCP group was 28% (21/75) and in ACP group 18.5% (13/72). The main clinical feature in ACP group was presence of pseudopancreatic cyst. It was present in 41.7% (30/72) of the alcoholic pancreatitis patients. History of alcohol intake was taken in all the patients. None of the patients in TCP group had history of alcohol consumption. In brief, 72 patients with alcoholic pancreatitis, 75 patients with tropical calcific pancreatitis (TCP), 100 healthy controls (HC), and 40 alcoholic controls (AC) free from any gastrointestinal disorder were recruited in the study (Table 2). Cases and controls were from the same ethnic group and were residents of the northern parts of India. The ethics committee of the institution approved the study. A written pre-informed consent was obtained prior to blood collection from the individuals.

## Method

### DNA Isolation

Blood sample was collected from each subject in EDTA coated vial. Genomic DNA was extracted by phenol chloroform method previously described (Miller et al., 1998). The purity and integrity of genomic DNA were checked on agarose gel (0.8%) and by calculating the ratio of absorbance at 260/280 nm (1.8–2.0). The DNA content was quantified by absorbance at 260 nm (1.00 optical density was considered equivalent to 50 mg/mL genomic DNA) before PCR amplification. DNA samples were stored at  $-80^{\circ}\text{C}$  till further use.

## GENOTYPING

### Genotyping for ADH3 Polymorphism

DNA was amplified with specific oligonucleotide primers. The forward and reverse primers were 5' GCTTTAAGAGTAAAT ATTCTGTCCC 3' and 5' AATCTACCTCTTTCCGAAGC 3', respectively. PCR amplification was performed in 25  $\mu\text{l}$  reaction mixture using the standardized PCR condition. The PCR condition included initial denaturation at  $95^{\circ}\text{C}$  for 5 min followed by 35 cycles of denaturation, annealing, and extension at  $94^{\circ}\text{C}$  for 30 s,  $55^{\circ}\text{C}$  for 30 s, and  $72^{\circ}\text{C}$  for 30 s, respectively. Final extension was done at  $72^{\circ}\text{C}$  for 10 min. The 145 bp ADH3 amplicons were digested with SspI restriction enzyme at  $37^{\circ}\text{C}$  for overnight. Genotyping was done by analyzing bands on 12% PAGE. ADH3<sup>1-1</sup> genotype: two bands (67 and 63 bp), ADH3<sup>1-2</sup> genotype: three bands ( $\sim$ 130, 67, and 63 bp), ADH3<sup>2-2</sup> genotype: one band (130 bp). A 15 bp cut was also present and not visualized on gel (Groppi et al., 1990).

**TABLE 2 | Age and Gender distribution.**

Characteristics	ACP	TCP	HC	AC
N	72	75	100	40
Sex (male/female)	All male	40/35	60/40	All male
Age (mean $\pm$ SD) in years	38.5 $\pm$ 8.3	32.8 $\pm$ 11.2	32.05 $\pm$ 4.6	35.9 $\pm$ 5.3

ACP, alcoholic chronic pancreatitis; TCP, tropical calcific pancreatitis; HC, healthy control; AC, alcoholic control.

### Genotyping for ALDH2 Polymorphism

ALDH2 genotyping was done by PCR-RFLP. The forward and reverse primers were 5' CAAATTACAGGGTCAAGGGCT 3' and 5' CCACACTCACAGTTTTCTCTT 3', respectively. PCR was done using standardized PCR condition. The PCR condition included initial denaturation at  $95^{\circ}\text{C}$  for 3 min followed by 35 cycles of denaturation, annealing, and extension at  $94^{\circ}\text{C}$  for 15 s,  $59.7^{\circ}\text{C}$  for 45 s, and  $72^{\circ}\text{C}$  for 30 s, respectively. Final extension was done at  $72^{\circ}\text{C}$  for 10 min. The 134 bp ALDH2 amplicons were digested with MboII restriction enzyme at  $37^{\circ}\text{C}$  for overnight. Genotyping was done by analyzing bands on 12% PAGE. ALDH2<sup>1-1</sup> genotype: one band (125 bp), ALDH2<sup>1-2</sup> genotype: two bands (125 and 134 bp), ALDH2<sup>2-2</sup> genotype: one band (134 bp; Muto et al., 2000).

### Genotyping for CYP2E1 Intron 6 (Dra I) Polymorphism

Polymorphism in the intron 6 of CYP2E1 was determined by PCR-RFLP method. Enzyme Dra I was used to detect the D and C polymorphism in intron 6. The specific primers were 5' TCGTCAGTTCCTGAAAGCAGG 3' and 5' GAGCTCTGATGGAAGTATCGCA 3'. The PCR condition were initial denaturation at  $95^{\circ}\text{C}$  for 4 min followed by 40 cycles of denaturation, annealing, and extension at  $94^{\circ}\text{C}$  for 1 min,  $60.4^{\circ}\text{C}$  for 45 s, and  $72^{\circ}\text{C}$  for 3 min, respectively. Final extension was done at  $72^{\circ}\text{C}$  for 10 min. The 995 bp amplified product was then digested with Dra I restriction enzyme at  $37^{\circ}\text{C}$  for 15–16 h. Genotyping was done on 3% agarose gel stained with ethidium bromide. DD genotype: three bands (572, 302, and 121 bp), CD genotype four bands (874, 572, 302, and 121 bp), CC genotype: two bands (874 and 121 bp; Kato et al., 1994; Wu et al., 1998).

## STATISTICAL ANALYSIS

Genotypic frequencies were calculated and odds ratios (OR) were determined by  $2 \times 2$  contingency table with 95% confidence intervals. Statistical significance for OR was calculated by chi-square test using SPSS Version 12 software. Fisher exact test was used, when expected cell frequency was less than five. Age and sex adjusted OR were calculated to assess the disease risk.

## RESULTS

The mean age and sex distribution are shown in the table (Table 2). Mean age of ACP patients was higher in comparison to TCP patients. Sex distribution in the studied groups was dissimilar as ACP and healthy alcoholics groups consist of males only.

### ADH3 Polymorphism

Frequency of ADH3\*1/ADH3\*1, ADH3\*1/ADH3\*2, and ADH3\*2/ADH3\*2 genotypes were 59.7 (43/72), 30.6 (22/72), and 9.7% (7/72), respectively in ACP group (Table 3; Figures 1A,B).

The prevalence of ADH3\*1/ADH3\*1 genotype was higher in ACP group in comparison to TCP, AC, and HC groups. The genotype frequency of ADH3\*1/ADH3\*1 was significantly higher in alcoholic pancreatitis in comparison to HC ( $p = 0.01$ ; OR = 2.6; 95% CI = 1.2–5.7) and AC ( $p = 0.02$ ; OR = 2.4; 95% CI = 1.1–5.4). The genotype frequency of ADH3\*1/ADH3\*1 was also significantly high in ACP group in comparison to TCP ( $p = 0.01$ , OR = 2.3; 95% CI = 1.2–4.5) group (Table 4).

The prevalence of ADH3\*1/ADH3\*1 genotype was 38.7% (29/75) in TCP group. Frequency of ADH3\*1/ADH3\*2 and ADH3\*2/ADH3\*2 genotype were 53.3 (40/75) and 8% (6/75), respectively in TCP group. ADH3\*1/ADH3\*1 genotype frequency in TCP group was comparable with HC and AC groups. Frequencies of both heterozygous ADH3\*1/ADH3\*2 genotype and homozygous ADH3\*2/ADH3\*2 genotype were statistically similar in TCP, HC, and AC groups (Table 3).

### ALDH2 and CYP2E1 Polymorphisms

ALDH2 is an alcohol metabolizing enzyme and polymorphism in this enzyme may affect individual susceptibility to develop the disease (Quertemont, 2004). We performed PCR-RFLP for ALDH2 in 90 patients (ACP: 50, TCP: 40) and 40 each alcoholic (AC) and healthy controls (HC) and found that all were monomorphic having ALDH2\*1/ALDH2\*1 homozygote

genotype. Additionally, we studied the CYP2E1 intron 6 Dra I polymorphism. In ACP group frequency of DD, CD, and CC genotypes were approximately 85% (61/72), 14% (10/72) and 1.4% (1/72), respectively. Frequency of DD and CD genotypes were 84 (63/75) and 16% (12/75), respectively in TCP group. Frequency distribution of CYP2E1 genotype is statistically similar in all the studied groups (Table 5).

### DISCUSSION


Alcohol misuse is now an established cause of chronic pancreatitis (Verlaan et al., 2004), and therefore the understanding of alcohol metabolism, its regulation, and polymorphisms of the alcohol metabolizing enzymes could be critical for the etiopathogenesis of ACP. There are various determinants of polymorphism, such as, age, sex, location, ethnicity, and race. ADH3, which is involved in the alcohol metabolism, has two alleles ADH3\*1 and ADH3\*2 (Edenberg, 2007). Since, ADH3\*1/ADH3\*1 genotype encodes for the fast metabolizing enzyme, the subjects carrying this genotype can metabolize ethanol at much faster rate resulting in rapid formation of acetaldehyde (Edenberg, 2007). To retain dynamic homeostasis, the conversion of acetaldehyde to acetate with same pace is essential. Malfunctioning of these proteins due to any mutation may lead the accumulation of acetaldehyde that may account for the disease causation (Majumdar et al., 1986; Nordback et al., 1991; Vonlaufen et al., 2007).

We observed strong association of ADH3\*1/ADH3\*1 genotype with alcoholic pancreatitis (Tables 3, 4; ACP vs. HC:  $p = 0.01$ ; OR = 2.6; 95% CI = 1.2–5.7 and ACP vs. AC:  $p = 0.02$ ; OR = 2.4; 95% CI = 1.1–5.4). Studies conducted in past on other populations worldwide have demonstrated varying results—some showing strong association of ADH3 polymorphism with ACP (Day et al., 1991; Dumas et al., 1995), some showing either statistically insignificant or no association of ADH3 polymorphism with ACP (Frenzer et al., 2002; Verlaan

TABLE 3 | Summary of frequency distribution of ADH3 genotypes.

Genotype	ACP (n = 72)	TCP (n = 75)	HC (n = 100)	AC (n = 40)
ADH3*1/*1	43 (59.7%)	29 (38.7%)	42 (42%)	15 (37.5%)
ADH3*1/*2	22 (30.6%)	40 (53.3%)	50 (50%)	22 (55%)
ADH3*2/*2	07 (9.7%)	6 (8%)	08 (8%)	03 (7.5%)

ACP, alcoholic chronic pancreatitis; HC, healthy control; AC, alcoholic control; TCP, tropical chronic pancreatitis.



**TABLE 4 | Summary of analysis of ADH3 frequency data.**

Groups	ACP vs. TCP	ACP vs. HC	ACP vs. AC	TCP vs. HC	TCP vs. AC	HC vs. AC
	p-value OR (95% CI)	p-value OR (95% CI)	p-value OR (95% CI)	p-value OR (95% CI)	p-value OR (95% CI)	p-value OR (95% CI)
ADH3 <sup>1-2</sup> vs. ADH3 <sup>1-1</sup> (ref)	0.06; 0.37 (0.18–0.74)	0.01; 0.43 (0.22–0.88)	0.01; 0.34 (0.15–0.8)	0.6; 1.1 (0.61–2.17)	0.88; 0.94 (0.41–2.11)	0.59; 1.2 (0.56–2.6)
ADH3 <sup>2-2</sup> vs. ADH3 <sup>1-1</sup> (ref)	0.69; 0.78 (0.24–2.5)	0.72; 0.85 (0.28–2.5)	0.78; 0.81 (0.18–3.5)	0.88; 1.0 (0.34–3.4)	0.96; 1.0 (0.22–4.7)	0.94; 1.0 (0.24–4.4)
ADH3 <sup>1-1</sup> vs. carriers of ADH3 <sup>2</sup> (ref)	0.01*; 2.3 (1.2–4.5)	0.01*; 2.6 (1.2–5.7)	0.02*; 2.4 (1.1–5.4)	0.7; 0.89 (0.48–1.6)	0.9; 1.0 (0.47–2.3)	0.67; 0.84 (0.39–1.8)

ACP, alcoholic chronic pancreatitis; TCP, tropical calcific pancreatitis; HC, healthy control; AC, alcoholic control; OR, odds ratio; CI, confidence interval.

**TABLE 5 | CYP2E1 frequency data.**

Genotype	ACP (n = 72)	TCP (n = 75)	HC (n = 90)	AC (n = 40)		
CC	1 (1.4%)	0	0	0		
CD	10 (14%)	12 (16%)	22 (24%)	07 (17.5%)		
DD	61 (85%)	63 (84%)	68 (76%)	33 (82.5%)		
P-VALUE AND OR (95% CI)						
	ACP vs. TCP	ACP vs. HC	ACP vs. AC	TCP vs. HC	TCP vs. AC	HC vs. AC
CD	0.9; 1 (0.27–3.7)	0.08; 0.4 (0.14–1.1)	0.5; 0.67 (0.21–3.1)	0.1; 0.5 (0.25–1.2)	0.3; 0.5 (0.1–1.9)	0.3; 0.63 (0.22–1.7)
DD (Ref)	Reference	Reference	Reference	Reference	Reference	Reference

ACP, alcoholic chronic pancreatitis; TCP, tropical calcific pancreatitis; HC, healthy control; AC, alcoholic control; OR, odds ratio; CI, confidence interval.

et al., 2004). This could be explained as association of ADH3 polymorphism and ACP risk highly depends on the ethnicity therefore could vary from one population to another or one ethnic group to another depending upon the localities they occupy and lifestyle factor.

The ALDH2 gene has two alleles, ALDH2\*1 and ALDH2\*2. Since, ALDH2 enzyme is the main form of aldehyde dehydrogenase responsible for acetaldehyde oxidation and polymorphism at its locus has been shown to be associated with chronic pancreatitis (Day et al., 1991), we also investigated ALDH2. Surprisingly, we were unable to detect any polymorphism in ALDH2 and all the subjects were homozygous for ALDH2\*1/ALDH2\*1 genotype. It clearly indicates that ALDH2 gene is monomorphic in the north Indian population and it might be possible that our population predominantly carried ALDH2\*1 allele. Our results were in agreement with the other finding by Bhaskar et al. in six Indian populations, which also showed that all the subjects had common homozygous genotype (ALDH2\*1/\*1; Bhaskar et al., 2007). Some other studies conducted previously in Japan and Korea have also shown variable frequency of ALDH2 among different populations (Kimura et al., 2000; Lee et al., 2001).

Furthermore, we also investigated polymorphism of CYP2E1, a key microsomal enzyme induced by ethanol, which metabolizes

alcohol in non-alcohol dehydrogenase pathway. We particularly investigated CYP2E1 intron 6D polymorphism, which exhibits two alleles: allele D and allele C. There was no significant difference in the frequency distribution of DD or CD genotype of CYP2E1; however, CC homozygous genotype was less frequent. Genotype frequencies were statistically similar in all the four groups. There was no difference observed in allele frequencies also. The results indicate no association of intron 6 Dra I polymorphism of CYP2E1 with tropical or alcoholic pancreatitis in North Indian population. Our results are in agreement with the previous reports showing no association of CYP2E1 polymorphism with alcoholic cirrhosis (Lee et al., 2001; Vidal et al., 2004). Nevertheless, a study of CYP2E1 intron 6 Dra I polymorphism showed a weak positive association of DD genotype to alcoholic pancreatitis (Verlaan et al., 2004).

In conclusion, the results demonstrate that ADH3 polymorphism may be an important contributor for alcoholic chronic pancreatitis among North Indian population. The polymorphism of ALDH2 and CYP2E1, probably do not have important role in ACP.

## ACKNOWLEDGMENTS

Authors sincerely thank Indian Council of Medical Research, New Delhi for providing research fellowship to DS.

## REFERENCES

- Bhaskar, L. V., Thangaraj, K., Osier, M., Reddy, A. G., Rao, A. P., Singh, L., et al. (2007). Single nucleotide polymorphism of the ALDH2 gene in six Indian populations. *Ann. Hum. Biol.* 34, 607–619. doi: 10.1080/03014460701581419
- Bisceglie, A. M., and Segal, I. (1984). Cirrhosis and chronic pancreatitis in alcoholics. *J. Clin. Gastroenterol.* 6, 199–200.
- Burim, R. V., Canalle, R., Martinelli Ade, L., and Takahashi, C. S. (2004). Polymorphisms in glutathione S-transferases GSTM1, GSTT1 and GSTP1 and cytochromes P4502E1 and CYP1A1 and susceptibility to cirrhosis or pancreatitis in alcoholics. *Mutagenesis* 19, 291–298. doi: 10.1093/mutage/geh034
- Cichoz-Lach, H., Celiński, K., and Słomka, M. (2008). Alcohol-metabolizing enzyme gene polymorphisms and alcohol chronic pancreatitis among Polish individuals. *HPB (Oxford)* 10, 138–143. doi: 10.1080/13651820801938909
- Crabb, D. W., Dipple, K. M., and Thomasson, H. R. (1993). Alcohol sensitivity, alcohol metabolism, risk of alcoholism and the role of alcohol and aldehyde dehydrogenase genotypes. *J. Lab. Clin. Med.* 122, 234–240.
- Czech, E., and Hartleb, M. (2003). Genetic polymorphism of alcohol dehydrogenase- pathophysiological implications. *Adv. Clin. Exp. Med.* 12, 801–809.
- Day, C. P., Bashir, R., James, O. F., Bassendine, M. F., Crabb, D. W., Thomasson, H. R., et al. (1991). Investigation of the role of polymorphism at the alcohol and aldehyde dehydrogenase loci in genetic predisposition to alcohol-related end-organ damage. *Hepatology* 14, 798–801. doi: 10.1002/hep.1840140509
- Duley, J. A., Harris, O., and Holmes, R. S. (1985). Analysis of human alcohol- and aldehyde-metabolizing isozymes by electrophoresis and isoelectric focusing. *Alcohol. Clin. Exp. Res.* 9, 263–271. doi: 10.1111/j.1530-0277.1985.tb05747.x
- Dumas, F., Coutelle, C., Gerolami, A., Groppi, A., Cassaigne, A., and Couzigou, P. (1995). Role of alcohol dehydrogenase polymorphism in alcoholic patients with chronic pancreatitis. *Gastroenterology* 104, A302.
- Dzienizewski, J., and Gabryelewicz, A. (2003). Pancreas; Pancreatitis- Polish accomplishments. *J. Physiol. Pharmacol.* 54, 155–166.
- Edenberg, H. J. (2000). Regulation of the mammalian alcohol dehydrogenase genes. *Prog. Nucleic Acid Res. Mol. Biol.* 64, 295–341. doi: 10.1016/S0079-6603(00)64008-4
- Edenberg, H. J. (2007). Role of alcohol dehydrogenase and aldehyde dehydrogenase variants. *Alcohol Res. Health* 30, 5–13.
- Frenzer, A., Butler, W. J., Norton, I. D., Wilson, J. S., Apte, M. V., Pirola, R. C., et al. (2002). Polymorphism in alcohol-metabolizing enzymes, glutathione S-transferases and apolipoprotein E and susceptibility to alcohol-induced cirrhosis and chronic pancreatitis. *J. Gastroenterol. Hepatol.* 17, 177–182. doi: 10.1046/j.1440-1746.2002.02670.x
- Greenfield, N. J., and Pietruszko, R. (1977). Two aldehyde dehydrogenases from human liver. Isolation via affinity chromatography and characterization of the isozymes. *Biochim. Biophys. Acta* 483, 35–45. doi: 10.1016/0005-2744(77)90005-5
- Groppi, A., Begueret, J., and Iron, A. (1990). Improved methods for genotype determination of human alcohol dehydrogenase (ADH) at ADH2 and ADH3 loci by using polymerase chain reaction directed mutagenesis. *Clin. Chem.* 36, 1765–1768.
- Harada, S., Agarwal, D. P., and Goedde, H. W. (1980). Electrophoretic and biochemical studies of human aldehyde dehydrogenase isozymes in various tissues. *Life Sci.* 26, 1773–1780. doi: 10.1016/0024-3205(80)90577-9
- Harber, P., Wilson, J., Apte, M., Korsten, M., and Pirola, R. (1995). Individual susceptibility to alcohol pancreatitis: still an enigma. *J. Lab. Clin. Med.* 125, 305–312.
- Ingelman-Sundberg, M., Johansson, I., Yin, H., Terelius, Y., Eliasson, E., Clot, P., et al. (1993). Ethanol-inducible cytochrome P4502E1: genetic polymorphism, regulation, and possible role in the etiology of alcohol-induced liver disease. *Alcohol* 10, 447–452. doi: 10.1016/0741-8329(93)90063-T
- Kato, S., Shields, P. G., Caporaso, N. E., Sugimura, H., Trivers, G. E., Tucker, M. A., et al. (1994). Analysis of cytochrome P450 2E1 genetic polymorphisms in relation to human lung cancer. *Cancer Epidemiol. Biomarkers Prev.* 3, 515–518.
- Khan, A. J., Husain, Q., Choudhuri, G., and Parmar, D. (2010). Association of polymorphism in alcohol dehydrogenase and interaction with other genetic risk factors with alcoholic liver cirrhosis. *Drug Alcohol Depend.* 109, 190–197. doi: 10.1016/j.drugalcdep.2010.01.010
- Khan, A. J., Ruwali, M., Choudhuri, G., Mathur, N., Husain, Q., and Parmar, D. (2009). Polymorphism in cytochrome P450 2E1 and interaction with other genetic risk factors and susceptibility to alcoholic liver cirrhosis. *Mutat. Res.* 664, 55–63. doi: 10.1016/j.mrfimm.2009.02.009
- Kimura, S., Okabayashi, Y., Inushima, K., Kochi, T., Yutsudo, Y., and Kasuga, M. (2000). Alcohol and aldehyde dehydrogenase polymorphisms in Japanese patients with alcohol-induced chronic pancreatitis. *Dig. Dis. Sci.* 45, 2013–2017. doi: 10.1023/A:1005650902961
- Lee, H. C., Lee, H. S., Jung, S. H., Yi, S. Y., Jung, H. K., Yoon, J. H., et al. (2001). Association between polymorphisms of ethanol-metabolizing enzymes and susceptibility to alcoholic cirrhosis in a Korean male population. *J. Korean Med. Sci.* 16, 745–750. doi: 10.3346/jkms.2001.16.6.745
- Liu, R., Yin, L. H., and Pu, Y. P. (2007). Association of combined CYP2E1 gene polymorphism with the risk for esophageal squamous cell carcinoma in Hua'an population, China. *Chin. Med. J.* 120, 1797–1802.
- Majumdar, A. P., Vesenska, G. D., Dubick, M. A., Yu, G. S., DeMorrow, J. M., and Geokas, M. C. (1986). Morphological and biochemical changes in the pancreas of rats treated with acetaldehyde. *Am. J. Physiol.* 250, G598–606.
- Miller, S. A., Dykes, D. D., and Polesky, H. F. (1998). A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res.* 16:1215. doi: 10.1093/nar/16.3.1215
- Muto, M., Hitomi, Y., Ohtsu, A., Ebihara, S., Yoshida, S., and Esumi, H. (2000). Association of aldehyde dehydrogenase 2 gene polymorphism with multiple oesophageal dysplasia in head and neck cancer patients. *Gut* 47, 256–261. doi: 10.1136/gut.47.2.256
- Nordback, I. H., MacGowan, S., Potter, J. J., and Cameron, J. L. (1991). The role of acetaldehyde in the pathogenesis of acute alcoholic pancreatitis. *Ann. Surg.* 214, 671–678. doi: 10.1097/0000658-199112000-00006
- Osier, M. V., Pakstis, A. J., Goldman, D., Edenberg, H. J., Kidd, J. R., and Kidd, K. K. (2002). A proline-threonine substitution in codon 351 of ADH1C is common in Native Americans. *Alcohol. Clin. Exp. Res.* 26, 1759–1763. doi: 10.1111/j.1530-0277.2002.tb02481.x
- Perri, F., Piepoli, A., Stanziale, P., Merla, A., Zelante, L., and Andriulli, A. (2003). Mutation analysis of the cystic fibrosis transmembrane conductance regulator (CFTR) gene, the cationic trypsinogen (PRSS1) gene, and the serine protease inhibitor, Kazal type 1 (SPINK1) gene in patients with alcoholic chronic pancreatitis. *Eur. J. Hum. Genet.* 11, 687–692. doi: 10.1038/sj.ejhg.5201035
- Quertemont, E. (2004). Genetic polymorphism in ethanol metabolism: acetaldehyde contribution to alcohol abuse and alcoholism. *Mol. Psychiatry* 9, 570–581. doi: 10.1038/sj.mp.4001497
- Singh, M., and Simsek, H. (1990). Ethanol and the pancreas. Current status. *Gastroenterology* 98, 1051–1062. doi: 10.1016/0016-5085(90)90033-W
- Song, B. J. (1996). Ethanol-inducible cytochrome P450 (CYP2E1): biochemistry, molecular biology and clinical relevance: 1996 update. *Alcohol. Clin. Exp. Res.* 20, 138A–146A. doi: 10.1111/j.1530-0277.1996.tb01764.x
- Verlaan, M., Te Morsche, R. H., Roelofs, H. M., Laheij, R. J., Jansen, J. B., Peters, W. H., et al. (2004). Genetic polymorphisms in alcohol-metabolizing enzymes and chronic pancreatitis. *Alcohol Alcohol.* 39, 20–24. doi: 10.1093/alcal/agh001
- Vidal, F., Lorenzo, A., Auguet, T., Olona, M., Broch, M., Gutiérrez, C., et al. (2004). Genetic polymorphisms of ADH2, ADH3, CYP4502E1 Dra-I and Pst-I, and ALDH2 in Spanish men: lack of association with alcoholism and alcoholic liver disease. *J. Hepatol.* 41, 744–750. doi: 10.1016/j.jhep.2003.06.003
- Vonlaufen, A., Wilson, J. S., Pirola, R. C., and Apte, M. V. (2007). Role of alcohol metabolism in chronic pancreatitis. *Alcohol Res. Health* 30, 48–54.
- Wu, X., Amos, C. L., Kemp, B. L., Shi, H., Jiang, H., Wan, Y., et al. (1998). Cytochrome P450 2E1 DraI polymorphisms in lung cancer in minority populations. *Cancer Epidemiol. Biomarkers Prev.* 7, 13–18.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Singh, Negi, Upadhyay and Choudhuri. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.