



Demonstration of Binding Induced Structural Plasticity in a SH2 Domain

Lorenzo Visconti¹, Angelo Toto¹, James A. Jarvis², Francesca Troilo¹,
Francesca Malagrino¹, Alfonso De Simone^{2*} and Stefano Gianni^{1*}

¹ Istituto Pasteur – Fondazione Cenci Bolognetti, Dipartimento di Scienze Biochimiche “A. Rossi Fanelli” and Istituto di Biologia e Patologia Molecolari del CNR, Sapienza Università di Roma, Rome, Italy, ² Department of Life Sciences, Imperial College London, London, United Kingdom

OPEN ACCESS

Edited by:

Gian Gaetano Tartaglia,
Catalan Institution for Research
and Advanced Studies (ICREA), Spain

Reviewed by:

Stefano Ricagno,
University of Milan, Italy
Ylva Ivarsson,
Uppsala University, Sweden

*Correspondence:

Alfonso De Simone
a.de-simon@imperial.ac.uk
Stefano Gianni
stefano.gianni@uniroma1.it

Specialty section:

This article was submitted to
Structural Biology,
a section of the journal
Frontiers in Molecular Biosciences

Received: 16 March 2020

Accepted: 17 April 2020

Published: 12 May 2020

Citation:

Visconti L, Toto A, Jarvis JA,
Troilo F, Malagrino F, De Simone A
and Gianni S (2020) Demonstration
of Binding Induced Structural
Plasticity in a SH2 Domain.
Front. Mol. Biosci. 7:89.
doi: 10.3389/fmolb.2020.00089

SH2 domains are common protein interaction domains able to recognize short aminoacidic sequences presenting a phosphorylated tyrosine (pY). In spite of their fundamental importance for cell physiology there is a lack of information about the mechanism by which these domains recognize and bind their natural ligands. The N-terminal SH2 (N-SH2) domain of PI3K mediates the interaction with different scaffolding proteins and is known to recognize a specific pY-X-X-M consensus sequence. These interactions are at the cross roads of different molecular pathways and play a key role for cell development and division. By combining mutagenesis, chemical kinetics and NMR, here we provide a complete characterization of the interaction between N-SH2 and a peptide mimicking the scaffolding protein Gab2. Our results highlight that N-SH2 is characterized by a remarkable structural plasticity, with the binding reaction being mediated by a diffused structural region and not solely by the residues located in the binding pocket. Furthermore, the analysis of kinetic data allow us to pinpoint an allosteric network involving residues far from the binding pocket involved in specificity. Results are discussed on the light of previous works on the binding properties of SH2 domains.

Keywords: allosteric network, skinetics, mutagenesis, peptide binding, NMR

INTRODUCTION

The interaction between proteins plays a paramount role in biological functions. Frequently, protein-protein recognition is mediated by specific domains such as SH2, SH3, WW, or PDZ domains (Macias et al., 1996; Pawson and Scott, 1997; Bhattacharyya et al., 2006; Teyra et al., 2012; Kurochkina and Guha, 2013). Whilst these domains are typically highly conserved and often very abundant in the cellular environment, they are nevertheless specific, such as cross-reactivity with non-desired partners is minimized (Zarrinpar et al., 2003). It has been observed that PDZ and SH3 domains control their ability to recognize specific consensus sequences through dynamic and energetic allosteric mechanisms (Lockless and Ranganathan, 1999; Fuentes et al., 2006; Gianni et al., 2011; Kumawat and Chakrabarty, 2017; Malagrino et al., 2019). These mechanisms overcome

the classical paradigm of allostery, which is based on major protein structural and conformational changes, and are characterized by fine rearrangement in the dynamic of the system (Kern and Zuiderweg, 2003; Motlagh et al., 2014). This behavior has been demonstrated for different protein-protein interaction domains. However, despite their critical roles in many important biological pathways, little information is currently available on Src Homology 2 (SH2) domains, for which only little evidence of interdomain allosteric regulations has been reported (Tsutsui et al., 2016; Gangopadhyay et al., 2020).

SH2 domains represent a classical example of modular protein interaction domains. SH2 domains are present in a wide range of proteins, including kinases, adaptors, phosphatases and other signaling molecules (Songyang et al., 1993; Payne et al., 1994), counting for more than 110 different proteins encoded by human genome. From a structural point of view, the SH2 domains are composed of about 100 amino acids presenting a conserved architecture comprising three to seven anti-parallel β strands organized in a β sheet, flanked by two α -helices (Waksman et al., 1992, 1993). SH2 domains from different proteins are known to recognize specific consensus sequences containing a phosphorylated tyrosine (pY) (Payne et al., 1994). In addition to the phospho-tyrosine, each SH2 domain recognizes several additional flanking residues, generally three to five amino acids C-terminal to pY (Schaffhausen, 1995). Many signaling pathways require the action of SH2 domains to control the colocalization of a variety of proteins within the signaling cascade, the binding events mediated by these domains allowing a correct signal transduction (Koch et al., 1991; Neel et al., 2003). For these reasons, mutations that affect the binding properties of the SH2 domains lead to an abnormal regulation of important signaling pathways, causing the misregulation of key physiological processes in the cell and consequently to the development of a number of diseases (Tartaglia et al., 2001; Lappalainen et al., 2008).

PI3K proteins interact with various RTKs involved in an ample range of signaling processes (Katso et al., 2001; Engelman et al., 2006). The activation of PI3K requires the interaction of its two SH2 domains with the binding partners, thus driving the phosphorylation of phosphatidylinositols at their 3' position. Both the N-terminal and the C-terminal SH2 domains (N-SH2 and C-SH2 respectively) recognize the consensus pY-X-X-M sequence of the target (Waksman et al., 1992; Songyang et al., 1993). The interaction between PI3K and their ligands is very often mediated by scaffolding proteins like, for example, Gab1, Gab2, IRS-1, FRS2, that act as platforms recruiting several transcription factors and allowing the correct formation of the molecular machinery.

The Grb2-associated binding protein (Gab2) is composed by an N-terminal folded PH domain and C-terminal unfolded region where the binding sites for other signaling proteins are situated. Because of its key role as scaffolding protein and in the binding with several transcription factors, Gab2 is implicated in several cancers of both solid and hematological origin (Ke et al., 2007; Ding et al., 2015). In particular Gab2 is overexpressed in breast (Bentires-Alj et al., 2006), gastric (Lee et al., 2007) and lung

(Xu et al., 2011) cancers, while its expression in healthy mature cells is relatively suppressed.

Here we provide, through a combination of stopped-flow and NMR experiments, a detailed characterization of the interaction between the N-terminal SH2 domain of PI3K and a 13-residue peptide mimicking Gab2, in its wild-type form and a variant M→A affecting the specific consensus pY-X-X-M sequence. Whilst the structural features of the N-SH2 has been previously described both in the free and in the bound state with other partners (Nolte et al., 1996), the mechanism of interaction of this protein with its natural ligands has never been characterized. Our data show that N-SH2 displays an unexpected structural plasticity, which does not reflect as major structural changes upon binding, but as a finely regulated rearrangement of the entire domain. Binding capabilities of N-SH2 are not solely mediated by its binding pocket but are modulated diffusely within its structure. In fact, by combining site-directed mutagenesis with kinetic and NMR experiments we provide compelling evidence highlighting a remarkable structural malleability of N-SH2 domain. The analysis of kinetic data allowed us to depict an allosteric network involving residues that are not located in the binding site but nevertheless appear to play a role in the recognition of the M residue, which is part of the consensus sequence of N-SH2. These results provide the first description of an allosteric network in the binding reaction of a SH2 domain.

MATERIALS AND METHODS

Protein Expression and Purification

N-SH2 wild-type and all the site-directed variants were expressed and purified as described previously (Visconti et al., 2019). All the mutants were obtained using a QuickChange Lightning Site-Directed Mutagenesis kit (Agilent technologies), accordingly to manufacturer instructions. Peptides with sequence TNSEDNpYVPMNPG mimicking wild-type Gab2 and TNSEDNpYVPMNPG mimicking the M457A variant of Gab2 were purchased from GenScript Biotech Corporation.

Stopped-Flow Experiments

Binding kinetics experiments were performed using a single-mixing SX-18 stopped-flow instrument (Applied Photophysics); all binding experiments were conducted in 50 mM Hepes pH 7.4 300 mM NaCl at 10°C. Excitation was at 280 nm and emission was collected using a 475 nm cut-off glass filter. Pseudo-first order binding experiments were performed mixing a constant concentration of dansylated Gab2_{448–460} and Gab2_{448–460} M457A (1 μ M) vs. N-SH2 wt and its mutants at concentrations ranging from 4 to 12 μ M. For each N-SH2 concentration, usually 5 individual traces were averaged and in all cases the fluorescence time courses obtained was satisfactorily fitted by using a single exponential equation.

NMR Experiments

Standard triple resonance experiments [CBCA(CO)NH, HNCACB, HNCO, HN(CA)CO spectra] and HSQC were recorded at 10°C on Bruker spectrometers operating at ¹H

frequencies of 600 MHz. Solution NMR experiments were carried out in buffer 50 mM Hepes pH 7.4 300 mM NaCl using a 100 μ M sample of labeled N-SH2. For the full assignment of the bound N-SH2 we performed titrations of ^1H - ^{15}N HSQC spectra of a 100 μ M sample of N-SH2 with both Gab2_{448–460} and Gab2_{448–460} M457A, which were recorded using progressive concentration of the ligand until 1:1 molar ratio.


RESULTS

Stopped-Flow Binding Kinetics

One of the aims of this study is to depict the molecular details of the interaction between the N-SH2 domain of PI3K and Gab2. A powerful strategy to achieve this goal relies on perturbing the system by producing conservative site-directed variants while monitoring the effect of mutations on the association and dissociation rate constants between the interacting partners (k_{on} and k_{off} respectively). Thus, we designed and successfully produced 21 site-directed variants of the N-SH2 domain and challenged them vs. a peptide mimicking the region of Gab2 ranging from residue 448–460 (Gab2_{448–460}), containing the pY-X-X-M motif that is specifically recognized by the N-SH2 domain (Waksman et al., 1992; Songyang et al., 1993). The choice of mutations was designed in analogy to phi-value analysis methodology in protein folding studies (Fersht and Sato, 2004). Positions mutated are showed in **Figure 1**. To monitor binding by fluorescence, a dansyl group was added at the N-terminus of the peptide. The binding reaction was monitored by stopped-flow experiments following the change in FRET signal, taking advantage of the two Trp residues on the N-SH2 domain in positions 12 and 14 as fluorescence donor and the dansyl group on Gab2_{448–460} as acceptor. The experiments were performed in pseudo-first order conditions with Gab2_{448–460} at fixed concentration of 1 μ M rapidly mixed vs. large excess of N-SH2 wt and its variants, ranging from 4 to 14 μ M, in buffer Hepes 50 mM, NaCl 300 mM, pH 7.4, at 10°C. For all the experiments observed time courses were found to be consistent with a single exponential equation. The dependence of observed rate constants k_{obs} as a function of the concentration of N-SH2 was fitted with a linear equation, suggesting that the binding reaction can be described by a simple two-state mechanism

$$k_{\text{obs}} = [\text{NSH2}]k_{\text{on}} + k_{\text{off}}$$

with k_{on} being the slope of the line and k_{off} the intercept with the y-axis. Since indirect calculation of k_{off} obtained by extrapolation is associated with high experimental error, we performed displacement experiments to directly measure k_{off} by challenging a pre-incubated complex of dansylated Gab2_{448–460} and N-SH2 domain both at the concentration of 1 μ M vs. a large excess of non-dansylated Gab2_{448–460} (50 μ M). The pseudo-first order plots of k_{obs} vs. the concentration of wild-type N-SH2 and its variants are shown in **Figure 2A**. Kinetic parameters extracted from data together with the calculated changes in activation and equilibrium free energies are reported in **Table 1**. Analysis of kinetic data highlights that T48S,




L51A, and L59A variations cause a detectable decrease of the affinity of the domain for Gab2_{448–460} of \sim 10-fold, affecting both the k_{on} and k_{off} . Furthermore, L99A mutation displays a strong destabilization effect on the complex, with an increase of dissociation equilibrium rate constant, K_{D} , of \sim 100-fold, mainly caused by a high increase of k_{off} . Since L99 residue is part of the binding pocket, this dramatic effect on the binding affinity is expected.

SH2 domains bind ligands with a phosphorylated tyrosine in their sequence. The N-SH2 domain of PI3K is known to recognize a specific consensus sequences that present a methionine in position +3 in respect to the phosphotyrosine (M457 in Gab2). To investigate the details of the role of M457 of Gab2 in the binding reaction we conducted kinetic binding experiments between Gab2_{448–460} M457A and all the site-directed variants of N-SH2 domains, under the same experimental conditions that were used for Gab2_{448–460}. In analogy to what observed with the wild-type peptide, the fluorescence change upon binding followed a single exponential decay, and dependences of k_{obs} vs. the concentration of N-SH2 were fitted with a linear equation (**Figure 2B**). Kinetic data obtained from binding and displacement experiments are reported in **Table 1**. Our data show that M457 is not essential for binding, the K_{D} obtained for the complex with N-SH2 wt being $3.7 \pm 0.4 \mu\text{M}$. The analysis of microscopic association and dissociation rate constants highlights that k_{on} is comparable with the one calculated for the binding with Gab2_{448–460}. However, a clear increase in k_{off} is reported for the binding with N-SH2 wt (Gab2_{448–460} $k_{\text{off}} = 3.9 \pm 0.2 \text{ s}^{-1}$; Gab2_{448–460} M457A $k_{\text{off}} = 80 \pm 4 \text{ s}^{-1}$) suggesting that M457 has a role in the dissociation of Gab2 with N-SH2, rather than in the recognition event. Interestingly, the k_{on} values obtained for the binding of Gab2_{448–460} M457A with all N-SH2 variants appear to be mostly unaffected by mutations occurring in N-SH2, whilst a more pronounced effect on the association rate constant is appreciable for the binding with wild type Gab2_{448–460}. This difference

TABLE 1 | Kinetic and thermodynamic parameters obtained from the analysis of the pseudo-first order binding experiments between site-directed variants of N-SH2 and Gab2_{448–460} wild-type and Gab2_{448–460} M457A.

N-SH2 variant	Gab2 _{448–460}				Gab2 _{448–460} M457A				$\Delta\Delta\Delta G$
	k_{on} ($\mu M^{-1} s^{-1}$)	k_{off} (s^{-1})	K_D (μM)	$\Delta\Delta G_{eq}$ (kcal mol ⁻¹)	k_{on} ($\mu M^{-1} s^{-1}$)	k_{off} (s^{-1})	K_D (μM)	$\Delta\Delta G_{eq}$ (kcal mol ⁻¹)	
WT	19.5 ± 1.0	3.9 ± 0.2	0.2 ± 0.1		21.5 ± 2.4	80 ± 4	3.7 ± 0.4		
L7A	15.7 ± 1.2	4.0 ± 0.2	0.3 ± 0.1	0.15 ± 0.03	19.3 ± 2.3	90 ± 4	4.6 ± 0.6	0.12 ± 0.01	**
I17V	24.4 ± 1.7	4.5 ± 0.2	0.2 ± 0.1	-0.05 ± 0.02	22.0 ± 2.3	90 ± 5	4.3 ± 0.5	0.09 ± 0.01	**
V22A	11.2 ± 0.4	7.4 ± 0.4	0.7 ± 0.1	0.68 ± 0.04	17.7 ± 2.6	170 ± 9	9.7 ± 1.5	0.55 ± 0.05	**
T29S	17.5 ± 0.5	4.3 ± 0.2	0.3 ± 0.1	0.13 ± 0.02	19.7 ± 2.7	80 ± 4	4.0 ± 0.6	0.05 ± 0.01	**
A30G	29.2 ± 2.0	3.7 ± 0.2	0.1 ± 0.1	-0.25 ± 0.02	17.1 ± 2.2	90 ± 4	5.1 ± 0.7	0.19 ± 0.10	**
T33S	22.6 ± 2.4	3.6 ± 0.2	0.2 ± 0.1	-0.12 ± 0.02	20.6 ± 2.9	80 ± 4	3.8 ± 0.6	0.02 ± 0.01	**
L35A	7.0 ± 0.6	6.7 ± 0.3	1.0 ± 0.1	0.88 ± 0.09	*	*	*	*	*
T48S	6.4 ± 0.5	23 ± 1	3.7 ± 0.1	1.64 ± 0.36	*	*	*	*	*
T50S	15.6 ± 0.8	9.3 ± 0.5	0.6 ± 0.1	0.62 ± 0.05	20.2 ± 2.0	140 ± 7	6.8 ± 0.8	0.35 ± 0.10	**
L51A	5.6 ± 1.1	18 ± 1	3.2 ± 0.2	1.56 ± 0.64	21.2 ± 3.3	160 ± 8	7.6 ± 1.2	0.41 ± 0.10	1.16 ± 0.42
L59A	6.7 ± 1.0	21 ± 1	3.2 ± 0.2	1.57 ± 0.52	*	*	*	*	*
I60V	9.1 ± 1.1	6.9 ± 0.3	0.8 ± 0.1	0.76 ± 0.10	15.2 ± 1.5	80 ± 4	5.0 ± 0.6	0.18 ± 0.02	0.58 ± 0.16
I62V	15.5 ± 1.4	4.5 ± 0.2	0.3 ± 0.1	0.21 ± 0.03	18.4 ± 1.5	120 ± 6	6.7 ± 0.6	0.34 ± 0.10	**
P74A	48.7 ± 1.5	27 ± 1	0.6 ± 0.1	0.59 ± 0.04	22.5 ± 1.9	60 ± 3	2.6 ± 0.3	-0.20 ± 0.02	0.79 ± 0.12
L75A	8.1 ± 1.1	6.1 ± 0.3	0.8 ± 0.1	0.76 ± 0.11	18.8 ± 3.2	110 ± 5	5.7 ± 1.0	0.25 ± 0.03	0.51 ± 0.17
T76S	16.9 ± 1.3	5.3 ± 0.3	0.3 ± 0.1	0.26 ± 0.03	19.6 ± 2.6	100 ± 5	5.0 ± 0.7	0.17 ± 0.02	**
V80A	16.0 ± 1.2	4.4 ± 0.2	0.3 ± 0.1	0.19 ± 0.03	22.3 ± 5.4	130 ± 6	5.6 ± 1.4	0.24 ± 0.02	**
V81A	12.0 ± 1.4	4.5 ± 0.2	0.4 ± 0.1	0.37 ± 0.05	20.5 ± 3.8	90 ± 5	4.5 ± 0.9	0.11 ± 0.01	**
L99A	8.2 ± 1.7	160 ± 10	20 ± 0.2	2.59 ± 0.30	*	*	*	*	*
V101A	13.4 ± 0.4	7.0 ± 0.4	0.5 ± 0.1	0.55 ± 0.06	14.2 ± 1.7	170 ± 9	12 ± 1	0.68 ± 0.10	**
L104A	16.5 ± 0.2	3.5 ± 0.2	0.2 ± 0.1	0.05 ± 0.02	22.1 ± 1.4	110 ± 6	5.2 ± 0.4	0.19 ± 0.02	**

*Binding was abolished by this mutation. **The absolute value of calculated $\Delta\Delta\Delta G$ was below 0.4 kcal mol⁻¹ and was excluded from analysis.




may suggest that M457 may be energetically connected with the residues on N-SH2 displaying a different behavior, possibly highlighting a complex role in the binding reaction with the N-SH2 and, therefore, demanding further investigations.

LFER Analysis and Characterization of TS of the Binding Reaction

An essential element to characterize the mechanism of the binding reaction of a domain is the analysis of the transition state (TS) of the reaction. In this context, the study of the correlation between the free energy of the activation barrier to that of the ground states, classically denoted as linear free energy relationship plot (LFER plot) (Leffler, 1953), is a widely used strategy in organic chemistry to analyse reactions involving formation of covalent bonds, as well as in enzymology (Toney and Kirsch, 1989), protein folding (Fersht, 2004), and binding studies (Eaton et al., 1991). LFER analysis correlates the changes in free energy of the transition state to the changes in equilibrium free energies; the slope of the correlation being denoted as α representing the position of the transition state along the reaction

coordinate. The mutational analysis performed in Table 1, allowed us to apply a LFER analysis on the interaction of N-SH2 with Gab2₄₄₈₋₄₆₀ and Gab2₄₄₈₋₄₆₀ M457A (Figure 3). Interestingly, in both cases, the measured data return a linear LFER plot. In analogy to what previously discussed in protein folding (Fersht, 2004), this finding is a hallmark of co-operativity and suggests that not only the residues located in the binding pocket, but all the probed residues taken are involved in the binding of the ligand.

It is of interest to compare the dependence of the plots measured for wild-type Gab2₄₄₈₋₄₆₀ and Gab2₄₄₈₋₄₆₀ M457A. In fact, whilst the apparent α value for the TS of N-SH2 upon binding with Gab2₄₄₈₋₄₆₀ is 0.44 ± 0.04 , in the case of Gab2₄₄₈₋₄₆₀ M457A we measured a value of 0.20 ± 0.09 . This finding would suggest that the M457A mutation may influence the structure of the transition state of N-SH2 in the binding reaction, suggesting that this protein may be characterized by a remarkable structural malleability that is dictated by the ligand. Because these differences appear however just above the limit of the experimental detection, to further validate this intriguing



scenario we resorted to perform NMR experiments with the two different ligands.

NMR

We then probed at an atomic resolution the structural malleability of N-SH2 using nuclear magnetic resonance (NMR). The sharp and well-dispersed backbone amide resonances in the ¹H-¹⁵N HSQC of N-SH2 in the free state (**Figure 4A**, in red) indicate that the protein is well folded under the experimental conditions employed *in plastico*. To obtain residue-specific information from these spectra, we assigned the resonances of the protein backbone using a combination of 3D spectra, including CBCA(CO)NH, HNCACB, HNCO, HN(CA)CO, which were analyzed using a computer-aided procedure as described in Fusco et al. (2012). The assigned ¹H-¹⁵N HSQC was employed to map the conformational changes and interaction surfaces of ¹⁵N labeled N-SH2 upon binding with increasing amounts of unlabeled Gab2₄₄₈₋₄₆₀ or Gab2₄₄₈₋₄₆₀ M457A in a titration experiment.

The resulting Chemical Shift Perturbation (CSP) is an established method to study protein-protein interactions, as it enables following the perturbations on the chemical environment of single residues upon ligand binding. Since changes in chemical shifts are very sensitive to structural changes, comparison of HSQC spectra of free and bound state of a protein may reveal important information about the residues involved in the process. In fact, whilst major changes are expected for residues that are in the binding interface, conformational changes occurring on distal residues, not directly involved in binding, may be diagnostic for allosteric communication in the protein.

The comparison between the HSQC of the N-SH2 in the free and in the bound state with Gab2₄₄₈₋₄₆₀ shows marked chemical shift differences, indicating significant perturbation of

the conformational properties of N-SH2 in the free and in the bound state (**Figure 4A**). Analogous results have been obtained for the binding of Gab2₄₄₈₋₄₆₀ M457A (**Figure 4B**) with major chemical shift changes observed as a result of this interaction. CSP analysis clearly indicates the nature of the cooperative mechanism of binding of N-SH2 domain with Gab2₄₄₈₋₄₆₀ and Gab2₄₄₈₋₄₆₀ M457A. This result provides further evidence that the LFER linearity is due to a global structural perturbation of the protein domain upon binding. Interestingly, the comparison of the peak shifts in the bound states with Gab2₄₄₈₋₄₆₀ and Gab2₄₄₈₋₄₆₀ M457A evidences that the mutation occurring in the ligand affects the plasticity of N-SH2 in the complex (**Figure 4C**). Our results suggest that the interaction of M457 with the binding site of N-SH2 generates a structural response in distal residues of the protein. This scenario implies the presence of an allosteric network in N-SH2 that finely regulates the binding properties and conformations of SH2.

Double Mutant Cycles

Protein allostery may be defined as the regulation of a given protein at a site other than the active (or binding) site. Allosteric regulation can be cryptic and escape characterization, occurring in the absence of major conformational changes and involving thermodynamic interactions of single distal residues. A powerful methodology to quantitatively calculate the interaction involved in such elusive allosteric regulations is to measure the energetic coupling between the ligand and residues that are not directly located in the binding pocket.

Double mutant cycle is a very effective strategy to infer energetic coupling and is based on the synergic employment of mutagenesis and binding experiments (Horovitz and Fersht, 1990). Thus, to investigate the selectivity of N-SH2 domain, we resorted to investigate the energetic coupling between its residues




FIGURE 4 | (A) Changes in the ^1H - ^{15}N -HSQC spectra of free N-SH2 (red peaks) and N-SH2 in complex with Gab2₄₄₈₋₄₆₀ (blue peaks). **(B)** Changes in the ^1H - ^{15}N -HSQC spectra of free N-SH2 (red peaks) and N-SH2 in complex with Gab2₄₄₈₋₄₆₀ M457A (green peaks). **(C)** Comparison of ^1H - ^{15}N -HSQC spectra of the bound states of N-SH2 with the two variants of Gab2 peptide. The analysis reveals that the M457A mutation in the ligand modulates the plasticity of N-SH2 in the complex.

and position M457 of Gab2, which represents a key residue in driving the selectivity of the domain. In analogy to previous work on other domains (Chi et al., 2008; Gianni et al., 2011), coupling free energies were calculated by comparing the changes in free energy obtained for the N-SH2 variants when binding to Gab2_{448–460} to those obtained when binding Gab2_{448–460} M457A, following the formalism:


$$\Delta\Delta\Delta G = \Delta\Delta G_{eq\ Gab2wt}^{mut-wt} - \Delta\Delta G_{eq\ Gab2M457A}^{mut-wt}$$

When $\Delta\Delta\Delta G = 0$ the (de)stabilization of the bound state due to the point mutation in N-SH2 when binding to Gab2_{448–460} equals the one occurring in the binding with Gab2_{448–460} M457A, meaning that M457 is not energetically coupled to the mutated residue in N-SH2. On the contrary, a $\Delta\Delta\Delta G \neq 0$ implies that the two positions interact energetically. The energetic couplings for each mutated residue of N-SH2 with M457 of Gab2 were calculated from kinetic parameters obtained from the analysis of pseudo-first order binding experiments and are reported in **Table 1**. We found that 4 residues (L51, I60, L75, and P74) reported a $\Delta\Delta\Delta G > 0.4$ kcal mol⁻¹ upon binding with Gab2 M457A, whilst for residues L35, T48, L59, L99 we could not detect binding. It is of particular interest to analyse the structural distribution of the residues that are affected by M457A mutation (**Figure 5**). In fact, while T48, L59, I60, and L99 are located in the binding site, L35, L51, P74, L75 are far from the binding pocket. These results, under the light of NMR data and LFER analysis, evidence the presence of an allosteric network in the N-SH2 domain that involve the whole domain, suggesting a key role for these specific residues in modulating the recognition of M457 of Gab2 and the affinity for the substrate.

DISCUSSION

In spite of their importance for cell physiology, there is lack of information pertaining the details of the mechanism of binding of SH2 domains. In this paper, we provided a detailed experimental characterization of the binding of N-SH2 to one of its physiological ligands and detected an unexpected structural plasticity. This effect was monitored both directly, by NMR by comparing the free structure with those of the bound states with two different ligands, and indirectly, by analyzing the effects of mutagenesis on the binding capability of the domain. NMR data reported subtle conformational changes in free and bound states. These minor changes in chemical shifts, supported by LFER and $\Delta\Delta\Delta G$ analysis, are indicative of a fine allosteric regulation of the binding mechanism.

It is of interest to compare our findings with those obtained on the SH2 domain from Itk. In fact, in the latter, it was demonstrated binding to be regulated by peptidyl-prolyl cis-trans isomerization (Mallis et al., 2002; Breheny et al., 2003; Pletneva et al., 2006; Severin et al., 2009). These findings appear to contrast the results reported in our work, which suggests that N-SH2 is characterized by a malleable structure, with binding being mediated by a diffused network involving several residues not directly located in the binding pocket, as mirrored by the



linearity of the LFER plot. Remarkably, in stark contrast to what observed in the case of the SH2 domain from Itk, a quantitative analysis of our NMR data revealed no evidence for peptidyl-prolyl cis-trans isomerization upon binding; therefore highlighting a completely different scenario. However, it is important to observe that P74 appears to retain a role in the recognition of M457 of Gab2, reporting a $\Delta\Delta\Delta G = 0.79 \pm 0.12$ kcal mol⁻¹. Thus, this residue, while maintaining a cis conformation both in the free and bound states, nevertheless participates to the long range energetic network modulating binding. Interestingly, P74 is conserved in both N-SH2 and C-SH2 domains of PI3K, that are both known to recognize the pY-X-X-M consensus sequence (Dornan and Burke, 2018).

Long allosteric networks regulating binding reactions have been observed for different adaptor protein domains, such as PDZ and SH3 domains (Gianni et al., 2011; Malagrino et al., 2019). Taken together with the experiments reported in this work, it appears that this mechanism of regulation may be typical of protein families that share the same topology but nevertheless must bind selectively specific ligands in the complex cellular environment. It has been demonstrated that SH2 domains can bind several peptides presenting different aminoacidic sequences containing a phosphotyrosine, with affinities differing of some orders of magnitude (Ladbury and Arold, 2000). We suggest that the conformational plasticity highlighted in our work, mediated by long range interactions, represents an additional mechanism of regulation of the SH2 moiety, tuning the selectivity of the domain, while maintaining a rather conserved topology and binding pocket. Future work on other SH2 domains will provide additional information about the generality of the allosteric regulation of the binding properties of SH2 domain family.

DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in article/supplementary material.

AUTHOR CONTRIBUTIONS

LV, AD, and SG designed the research. LV, AT, JJ, FM, and FT performed the research. LV and AT wrote the first version of the manuscript. All the authors analyzed data and revised the manuscript.

REFERENCES

- Bentires-Alj, M., Gil, S. G., Chan, R., Wang, Z. C., Wang, Y., Imanaka, N., et al. (2006). A role for the scaffolding adapter GAB2 in breast cancer. *Nat. Med.* 12, 114–121. doi: 10.1038/nm1341
- Bhattacharyya, R. P., Reményi, A., Yeh, B. J., and Lim, W. A. (2006). Domains, motifs, and scaffolds: the role of modular interactions in the evolution and wiring of cell signaling circuits. *Annu. Rev. Biochem.* 75, 655–680. doi: 10.1146/annurev.biochem.75.103004.142710
- Breheny, P. J., Laederach, A., Fulton, D. B., and Andreotti, A. H. (2003). Ligand specificity modulated by prolyl imide bond Cis/Trans isomerization in the Itk SH2 domain: a quantitative NMR study. *J. Am. Chem. Soc.* 125, 15706–15707. doi: 10.1021/ja0375380
- Chi, C. N., Elfström, L., Shi, Y., Snäll, T., Engström, A., and Jemth, P. (2008). Reassessing a sparse energetic network within a single protein domain. *Proc. Natl. Acad. Sci. U.S.A.* 105, 4679–4684. doi: 10.1073/pnas.0711732105
- Ding, C. B., Yu, W. N., Feng, J. H., and Luo, J. M. (2015). Structure and function of Gab2 and its role in cancer (Review). *Mol. Med. Rep.* 12, 4007–4014. doi: 10.3892/mmr.2015.3951
- Dornan, G. L., and Burke, J. E. (2018). Molecular mechanisms of human disease mediated by oncogenic and primary immunodeficiency mutations in class IA phosphoinositide 3-kinases. *Front. Immunol.* 9:575. doi: 10.3389/fimmu.2018.00575
- Eaton, W. A., Henry, E. R., and Hofrichter, J. (1991). Application of linear free energy relations to protein conformational changes: the quaternary structural change of hemoglobin. *Proc. Natl. Acad. Sci. U.S.A.* 88, 4472–4475. doi: 10.1073/pnas.88.10.4472
- Engelman, J. A., Luo, J., and Cantley, L. C. (2006). The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat. Rev. Genet.* 7, 606–619. doi: 10.1038/nrg1879
- Fersht, A. R. (2004). Relationship of Leffler (Bronsted) alpha values and protein folding Phi values to position of transition-state structures on reaction coordinates. *Proc. Natl. Acad. Sci. U.S.A.* 101, 14338–14342. doi: 10.1073/pnas.0406091101
- Fersht, A. R., and Sato, S. (2004). Phi-value analysis and the nature of protein-folding transition states. *Proc. Natl. Acad. Sci. U.S.A.* 101, 7976–7981. doi: 10.1073/pnas.0402684101
- Fuentes, E. J., Gilmore, S. A., Mauldin, R. V., and Lee, A. L. (2006). Evaluation of energetic and dynamic coupling networks in a PDZ domain protein. *J. Mol. Biol.* 364, 337–351. doi: 10.1016/j.jmb.2006.08.076
- Fusco, G., De Simone, A., Hsu, S.-T. D., Bemporad, F., Vendruscolo, M., Chiti, F., et al. (2012). 1H, 13C and 15N resonance assignments of human muscle acylphosphatase. *Biomol. NMR Assign.* 6, 27–29. doi: 10.1007/s12104-011-9318-1
- Gangopadhyay, K., Manna, B., Roy, S., Kumari, S., Debnath, O., Chowdhury, S., et al. (2020). An allosteric hot spot in the tandem-SH2 domain of ZAP-70 regulates T-cell signaling. *Biochem. J.* 477, 1287–1308. doi: 10.1042/BCJ20190879
- Gianni, S., Haq, S. R., Montemiglio, L. C., Jürgens, M. C., Engström, Å., Chi, C. N., et al. (2011). Sequence-specific long range networks in PSD-95/discs large/ZO-1 (PDZ) domains tune their binding selectivity. *J. Biol. Chem.* 286, 27167–27175. doi: 10.1074/jbc.M111.239541
- Horowitz, A., and Fersht, A. R. (1990). Strategy for analysing the co-operativity of intramolecular interactions in peptides and proteins. *J. Mol. Biol.* 214, 613–617. doi: 10.1016/0022-2836(90)90275-Q
- Katso, R., Okkenhaug, K., Ahmadi, K., White, S., Timms, J., and Waterfield, M. D. (2001). Cellular function of phosphoinositide 3-kinases: implications for development, homeostasis, and cancer. *Annu. Rev. Cell Dev. Biol.* 17, 615–675. doi: 10.1146/annurev.cellbio.17.1.615
- Ke, Y., Wu, D., Princen, F., Nguyen, T., Pang, Y., Lesperance, J., et al. (2007). Role of Gab2 in mammary tumorigenesis and metastasis. *Oncogene* 26, 4951–4960. doi: 10.1038/sj.onc.1210315
- Kern, D., and Zuiderweg, E. R. P. (2003). The role of dynamics in allosteric regulation. *Curr. Opin. Struct. Biol.* 13, 748–757. doi: 10.1016/j.sbi.2003.10.008
- Koch, C. A., Anderson, D., Moran, M. F., Ellis, C., and Pawson, T. (1991). SH2 and SH3 domains: elements that control interactions of cytoplasmic signaling proteins. *Science* 252, 668–674. doi: 10.1126/science.1708916
- Kumawat, A., and Chakrabarty, S. (2017). Hidden electrostatic basis of dynamic allostery in a PDZ domain. *Proc. Natl. Acad. Sci. U.S.A.* 114, E5825–E5834. doi: 10.1073/pnas.1705311114
- Kurochkina, N., and Guha, U. (2013). SH3 domains: modules of protein-protein interactions. *Biophys. Rev.* 5, 29–39. doi: 10.1007/s12551-012-0081-z
- Ladbury, J. E., and Arold, S. (2000). Searching for specificity in SH domains. *Chem. Biol.* 7, R3–R8. doi: 10.1016/s1074-5521(00)00067-3
- Lappalainen, I., Thusberg, J., Shen, B., and Vihinen, M. (2008). Genome wide analysis of pathogenic SH2 domain mutations. *Proteins* 72, 779–792. doi: 10.1002/prot.21970
- Lee, S. H., Jeong, E. G., Nam, S. W., Lee, J. Y., Yoo, N. J., and Lee, S. H. (2007). Increased expression of Gab2, a scaffolding adaptor of the tyrosine kinase signalling, in gastric carcinomas. *Pathology* 39, 326–329. doi: 10.1080/00313020701329773
- Leffler, J. E. (1953). Parameters for the description of transition states. *Science* 117, 340–341. doi: 10.1126/science.117.3039.340
- Lockless, S. W., and Ranganathan, R. (1999). Evolutionarily conserved pathways of energetic connectivity in protein families. *Science* 286, 295–299. doi: 10.1126/science.286.5438.295
- Macias, M. J., Hyvönen, M., Baraldi, E., Schultz, J., Sudol, M., Saraste, M., et al. (1996). Structure of the WW domain of a kinase-associated protein complexed with a proline-rich peptide. *Nature* 382, 646–649. doi: 10.1038/382646a0
- Malagrino, F., Troilo, F., Bonetti, D., Toto, A., and Gianni, S. (2019). Mapping the allosteric network within a SH3 domain. *Sci. Rep.* 9:8279. doi: 10.1038/s41598-019-44656-8
- Mallis, R. J., Brazin, K. N., Fulton, D. B., and Andreotti, A. H. (2002). Structural characterization of a proline-driven conformational switch within the Itk SH2 domain. *Nat. Struct. Biol.* 9, 900–905. doi: 10.1038/nsb864
- Motlagh, H. N., Wrabl, J. O., Li, J., and Hilser, V. J. (2014). The ensemble nature of allostery. *Nature* 508, 331–339. doi: 10.1038/nature13001
- Neel, B. G., Gu, H., and Pao, L. (2003). The 'Shp'ing news: SH2 domain-containing tyrosine phosphatases in cell signaling. *Trends Biochem. Sci.* 28, 284–293. doi: 10.1016/S0968-0004(03)00091-4

- Nolte, R. T., Eck, M. J., Schlessinger, J., Shoelson, S. E., and Harrison, S. C. (1996). Crystal structure of the PI 3-kinase p85 amino-terminal SH2 domain and its phosphopeptide complexes. *Nat. Struct. Biol.* 3, 364–374. doi: 10.1038/nsb0496-364
- Pawson, T., and Scott, J. D. (1997). Signaling through scaffold, anchoring, and adaptor proteins. *Science* 278, 2075–2080. doi: 10.1126/science.278.5346.2075
- Payne, G., Stolz, L. A., Pei, D., Band, H., Shoelson, S. E., and Walsh, C. T. (1994). The phosphopeptide-binding specificity of Src family SH2 domains. *Chem. Biol.* 1, 99–105. doi: 10.1016/1074-5521(94)90047-7
- Pletneva, E. V., Sundd, M., Fulton, D. B., and Andreotti, A. H. (2006). Molecular details of Itk activation by prolyl isomerization and phospholigand binding: the NMR structure of the Itk SH2 domain bound to a phosphopeptide. *J. Mol. Biol.* 357, 550–561. doi: 10.1016/j.jmb.2005.12.073
- Schaffhausen, B. (1995). SH2 domain structure and function. *Biochim. Biophys. Acta* 1242, 61–75. doi: 10.1016/0304-419x(95)00004-y
- Severin, A., Joseph, R. E., Boyken, S., Fulton, D. B., and Andreotti, A. H. (2009). Proline isomerization preorganizes the Itk SH2 domain for binding to the Itk SH3 domain. *J. Mol. Biol.* 387, 726–743. doi: 10.1016/j.jmb.2009.02.012
- Songyang, Z., Shoelson, S. E., Chaudhuri, M., Gish, G., Pawson, T., Haser, W. G., et al. (1993). SH2 domains recognize specific phosphopeptide sequences. *Cell* 72, 767–778. doi: 10.1016/0092-8674(93)90404-e
- Tartaglia, M., Mehler, E. L., Goldberg, R., Zampino, G., Brunner, H. G., Kremer, H., et al. (2001). Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2, cause Noonan syndrome. *Nat. Genet.* 29, 465–468. doi: 10.1038/ng772
- Teyra, J., Sidhu, S. S., and Kim, P. M. (2012). Elucidation of the binding preferences of peptide recognition modules: SH3 and PDZ domains. *FEBS Lett.* 586, 2631–2637. doi: 10.1016/j.febslet.2012.05.043
- Toney, M. D., and Kirsch, J. F. (1989). Direct Brønsted analysis of the restoration of activity to a mutant enzyme by exogenous amines. *Science* 243, 1485–1488. doi: 10.1126/science.2538921
- Tsutsui, Y., Deredge, D., Wintrodge, P. L., and Hays, F. A. (2016). Imatinib binding to human c-Src is coupled to inter-domain allostery and suggests a novel kinase inhibition strategy. *Sci. Rep.* 6:30832. doi: 10.1038/srep30832
- Visconti, L., Malagrino, F., Toto, A., and Gianni, S. (2019). The kinetics of folding of the NSH2 domain from p85. *Sci. Rep.* 9:4058. doi: 10.1038/s41598-019-40480-2
- Waksman, G., Kominos, D., Robertson, S. C., Pant, N., Baltimore, D., Birge, R. B., et al. (1992). Crystal structure of the phosphotyrosine recognition domain SH2 of v-src complexed with tyrosine-phosphorylated peptides. *Nature* 358, 646–653. doi: 10.1038/358646a0
- Waksman, G., Shoelson, S. E., Pant, N., Cowburn, D., and Kuriyan, J. (1993). Binding of a high affinity phosphotyrosyl peptide to the Src SH2 domain: crystal structures of the complexed and peptide-free forms. *Cell* 72, 779–790. doi: 10.1016/0092-8674(93)90405-f
- Xu, X. L., Wang, X., Chen, Z. L., Jin, M., Yang, W., Zhao, G. F., et al. (2011). Overexpression of Grb2-associated binder 2 in human lung cancer. *Int. J. Biol. Sci.* 7, 496–504. doi: 10.7150/ijbs.7.496
- Zarrinpar, A., Park, S.-H., and Lim, W. A. (2003). Optimization of specificity in a cellular protein interaction network by negative selection. *Nature* 426, 676–680. doi: 10.1038/nature02178

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Visconti, Toto, Jarvis, Troilo, Malagrino, De Simone and Gianni. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.