



Silencing of the tRNA Modification Enzyme Cdkal1 Effects Functional Insulin Synthesis in NIT-1 Cells: tRNA^{Lys3} Lacking ms²- (ms²t⁶A₃₇) is Unable to Establish Sufficient Anticodon:Codon Interactions to Decode the Wobble Codon AAG

Amithi Narendran¹, Sweta Vangaveti², Srivathsan V. Ranganathan³, Emily Eruysal¹, Miranda Craft¹, Omar Alrifai¹, Fu Yee Chua¹, Kathryn Sarachan¹, Breann Litwa¹, Sheetal Ramachandran⁴ and Paul F. Agris^{1,4*}

OPEN ACCESS

Edited by:

May Khanna,
University of Arizona, United States

Reviewed by:

Lluís Ribas De Pouplana,
Institute for Research in Biomedicine,
Spain
Natraj Krishnan,
Mississippi State University,
United States

*Correspondence:

Paul F. Agris
Paul.Agris@duke.edu

Specialty section:

This article was submitted to
Cellular Biochemistry,
a section of the journal
Frontiers in Molecular Biosciences

Received: 16 July 2020

Accepted: 31 December 2020

Published: 09 February 2021

Citation:

Narendran A, Vangaveti S, Ranganathan SV, Eruysal E, Craft M, Alrifai O, Chua FY, Sarachan K, Litwa B, Ramachandran S and Agris PF (2021) Silencing of the tRNA Modification Enzyme Cdkal1 Effects Functional Insulin Synthesis in NIT-1 Cells: tRNA^{Lys3} Lacking ms²- (ms²t⁶A₃₇) is Unable to Establish Sufficient Anticodon:Codon Interactions to Decode the Wobble Codon AAG. *Front. Mol. Biosci.* 7:584228. doi: 10.3389/fmolb.2020.584228

¹The RNA Institute and Department of Biological Sciences, University of Albany, Albany, NY, United States, ²Program in Bioinformatics and Integrative Biology, University of Massachusetts Medical School, Worcester, MA, United States, ³Knight Cancer Institute, Oregon Health Sciences, School of Medicine, Portland, OR, United States, ⁴Department of Medicine, Duke University School of Medicine, Durham, NC, United States

Human Genome Wide Association Studies found a significant risk of Type 2 Diabetes Mellitus (T2DM) in single nucleotide polymorphisms in the *cdkal1* gene. The *cdkal1* gene is remote from the insulin gene and with the surprising function of a specific tRNA modification. Population studies and case control studies acquired evidences of the connection between Cdkal1 protein and insulin production over the years. To obtain biochemical proofs directly linking potential SNPs to their roles in insulin production and availability is challenging, but the development of Cdkal1 knock out mice and knock out cell lines made it possible to extend our knowledge towards therapeutic field of diabetic research. Supporting the evidences, here we show that knock down of the *cdkal1* gene using small interfering and short hairpin RNA in the NIT-1 cell line, a β -cell line inducible for insulin resulted in reduced levels of *cdkal1* and mature insulin mRNAs, increased the level of precursor insulin mRNA, decreased Cdkal1 and insulin proteins, and diminished modification of tRNA^{Lys3} from t⁶A₃₇ to ms²t⁶A₃₇, the specified function of Cdkal1. tRNA^{Lys3} lacking ms²- is incapable of establishing sufficient hydrogen bonding energy and hydrophobic stabilization to decode the wobble codon AAG.

Keywords: Type 2 diabetes, CDKAL1, tRNA modification, Cdkal1 protein, insulin, Wobble decoding on the ribosome

INTRODUCTION


Type 2 Diabetes Mellitus (T2DM) accounts for 90% of the 34 million diabetes cases in the US (Centers for Disease Control and Prevention, 2020). A fundamental genetic heterogeneity has been found in T2DM (Mannino et al., 2019). Genome Wide Association Studies (GWAS) focused on the genetic heterogeneity have discovered a number of loci, *KLF14*, *KCNQ1*, *DUSP9*, *FTO*, *HNF4A*, *IGFBP2*, *CDKN2A/B*, *TCF7L2*, *KCNJ11*, antioxidant genes, *DNAJC3*, *PGC-1 α* , *ADIPOQ*, *CDKAL1*, *POMC*, *PPAR γ 2*, and *SLC30A8*, that could be potential predictors of the disease (Cho et al., 2019; Witka et al., 2019). One among the loci, in particular, CDK5 regulatory subunit associated protein 1-

like 1, *cdk11*, is intriguing as a locus remote from the insulin gene, and a tRNA modification enzyme. The gene product of *cdk11*, Cdkal1 post-transcriptionally modifies tRNA to facilitate accurate translation of the insulin mRNA and processing of proinsulin to mature insulin protein (McCown, et al., 2020). Thus, Cdkal1 has been found to be a potentially important factor in determining whether a patient is at a high risk for diabetes (Adami and Bottai, 2020; Krentz and Gloyn, 2020).

Cdkal1 protein is a post-transcriptional tRNA modification enzyme, a methylthiotransferase which adds a methylthio (ms²-) moiety to the 2-position of an already modified adenosine-37, N⁶-threonylcarbamoyladenine-37 (t⁶A₃₇) in tRNA^{Lys3}. The resulting ms²t⁶A₃₇, 3'-adjacent to the anticodon (Figure 1), augments translational fidelity for the tRNA to bind the lysine codons AAA/G on the ribosome (Arragain et al., 2010; Agris et al., 2017). Only fully modified tRNA^{Lys3} is capable of accurately and efficiently decoding the AAA and AAG codons (Agris et al., 2017). Of the ~five *cdk11* isoforms, only *cdk11* is expressed in human islets and pancreatic cells in culture (Brambillasca et al., 2012). GWAS have found single nucleotide polymorphisms (SNPs) within intron-5 of the human *cdk11* gene on human chromosome 6 that are considered a high risk for the disease (Diabetes Genetics Initiative of Broad Institute et al., 2007; Steinthorsdottir et al., 2007). The homozygous recessive mutation in the *cdk11* gene has a 1.50 risk of T2DM, comparable to *brca1* and 2. We found that the *cdk11* gene is likewise critical to *Drosophila melanogaster*, the organism having insulin-like peptides. The gene *CG6550* catalyzes the methylthiolation of N⁶-threonylcarbamoyladenine leading to

the formation of ms²t⁶A₃₇ in *Drosophila* tRNAs. The homozygous mutant *CG6550* (Df(2R)BSC44/P{EP}G7578), with the exception of rare escaper males, was lethal. Meta-analysis of genetic variations including homozygous recessive mutations in Cdkal1 shows that the occurrence of single nucleotide polymorphisms in the introns of Cdkal1 effect pre-mRNA processing events which impacts the production, processing, and availability of human insulin.

In β-cells, the protein Cdkal1 uniquely modifies just one of ~45 human tRNA species, tRNA^{Lys3}. tRNA^{Lys3} is one of the three tRNAs for the amino acid lysine. Cdkal1 is a member of the iron-sulfur cluster enzymes that use the substrate S-adenosylmethionine (Forouhar et al., 2013; Landgraf et al., 2016). Recently and understandably, iron-deficiency has been associated with T2DM through Cdkal1 under-performing for lack of iron (Santos et al., 2020). The AAG codon in human pre-proinsulin mRNA codes for both Lys53 and Lys88. Importantly, Lys88 is positioned adjacent to Arg89 and establishes the point at which a crucial protease cleavage separates the insulin A-chain from the C-peptide. Thus, in *cdk11*-associated T2DM it is believed that a non-functioning or missing Cdkal1 protein in islet β-cells would result in tRNA^{Lys3} lacking the modification ms²t⁶A₃₇. The tRNA^{Lys3} in turn would not be able to insert lysine at position 88 in response to the AAG codon. As a result, proinsulin would not be cleaved into insulin. In this study, we knocked down the *cdk11* gene in cell culture to show the effect of Cdkal1 silencing in pre-mRNA processing and production of mature insulin. The result was not only the *cdk11* mRNA and Cdkal1 protein were decreased, but mature insulin was also



reduced. The tRNA modification ms²t⁶A₃₇ was reduced relative to its precursor, t⁶A₃₇ in NIT-1 cells. We are first to demonstrate why tRNA^{Lys3} lacking the ms²-modification of ms²t⁶A₃₇ is unable to decode the lysine wobble codon AAG.

MATERIALS AND METHODS

Cell Line

NIT-1 is a β-cell line established from a transgenic mouse with the SV40 large T-antigen. It is grown in Hams F12K medium (F12K with L-Glutamine 90%; heat-inactivated, dialyzed fetal bovine serum 10%, FBS, Sigma; and 1 X penicillin-streptomycin, Invitrogen). Theophylline (10 mM with 5.5 mM glucose), glucose or KCl were used to stimulate insulin production. To knockdown the gene, we transfected an experimental esiRNA (endonuclease-prepared siRNA, Sigma) and a control GFP esiRNA with Lipofectamine 2000 and a Lentiviral shRNA (Sigma).

Stimulation of Insulin Production

NIT-1 has a glucose stimulated insulin response (GSIR) generated with a high glucose concentration (25 mM) when preceded by overnight incubation in low glucose, serum-free medium (SF-DMEM). Insulin production was assayed by ELISA (mouse proinsulin and insulin antibody, ABClonal).

Transfections

Plated cells were transfected with a control GFP esiRNA or *cdk1* esiRNA (30, 50 or 70 nM using Lipofectamine, LF, RNAiMax reagent, 0.5 or 1 μL, Invitrogen) in antibiotic-free medium (6 h). Cells were then incubated in fresh medium (48 h), serum and glucose starved overnight in SF-DMEM, and induced for insulin (90 min) or left unstimulated (SF-DMEM with 600 KIU/ml aprotinin).

Modified Nucleoside Analysis

Small RNAs (<200 nucleotides) were isolated (Ambion mirVana miRNA Isolation Kit) and the RNA was dialyzed extensively against phosphate buffer (10 mM NaH₂PO₄, pH 6.8) and then against water (18 mΩ). The RNA was hydrolyzed to nucleosides enzymatically rather than chemical digestion. The 2-step process cleaves first the phosphodiester bond with nuclease P1 resulting in nucleoside-5'-monophosphates followed by bacterial alkaline phosphatase (BAP) to cleave the 5'-phosphate from the nucleosides resulting in individual nucleosides and phosphoric acid. The modified nucleoside analysis was conducted by UHPLC-MS/MS (triple quadrupole MS (Waters MS) (Basanta-Sanchez et al., 2016).

RT-qPCR of Insulin and *cdk1* mRNA

We used RT-qPCR to assess the level of expression of mature insulin (mouse insulin I and II) or a pre-insulin or precursor containing intron 2 in normal and in esiRNA transfected NIT-1 cells, induced and not induced for insulin production. Large sequence RNA was isolated from normal and transfected


knockdown NIT-1 cells (Qiagen RNEasy Plus kit) and rRNA was removed (Qiagen RNEasy MinElute Kit). A 2-step RT-PCR was conducted to determine the presence of *cdk1*, precursor insulin, and mature insulin mRNAs (BioRad 2step c-DNA Synthesis kit) (Iype et al., 2005). The expression and amount of *cdk1* mRNA was determined by quantitative real-time PCR (RT-qPCR) analysis. Threshold Cq values were normalized to actin levels. Relative expression was calculated using the 2^{ΔΔCq} method (Livak and Schmittgen, 2001).

Molecular Dynamics Simulations

The crystal structure of the mammalian ribosome was obtained from the Protein Data Bank (PDB ID: 5LZS) (Shao et al., 2016). An intact stable fragment of structure was used for simulations, which included the mRNA, the anticodon stem loop (ASL) of the A-site tRNA, ribosomal RNA and ribosomal proteins within 25 Å of the codon and anticodon minihelix at the A-site. The ASL and the mRNA codon were modified to match the human tRNA^{Lys3} ASL sequence and the lysine codon respectively using MOE (Chemical Computing Group, 2019). Six different constructs of the ASL:codon pair were modeled with codons AAA and AAG, each paired with the ASL containing the unmodified nucleoside A₃₇, N⁶-threonylcarbonyl adenosine (t⁶A₃₇) and hypermodified 2-methylthio N⁶-threonylcarbonyl adenosine (ms²t⁶A₃₇).

In order to simulate the modified tRNA, AMBER (Cornell et al., 1995) type force-field parameters were developed for the atoms of the modified nucleosides—pseudouridine Ψ, mcm⁵s²U, t⁶A and ms²t⁶A. The geometry of the modified nucleosides was optimized using Hatree-Fock level theory and 6-31G* basis-sets in Webmo (Schmidt and Polik, 2020). For obtaining the partial charges on the atoms, the online RESP charge-fitting server REDS was used (Cornell et al., 1993; Dupradeau et al., 2010). AMBER-99 force field parameters and AMBER-99 parameters with the Chen-Garcia correction were used for bonded and Lennard-Jones (LJ) interactions, respectively (Cornell et al., 1995; Chen and García 2013).

Molecular dynamics (MD) simulations were performed using Gromacs-2016.4 and Gromacs-2019.6 packages (Abraham et al., 2015). The MD simulations incorporated a leap-frog algorithm with a 2-fs timestep to integrate the equations of motion. The system was maintained at 300 K, using the velocity rescaling thermostat (Bussi et al., 2007). The pressure was maintained at 1 atm using the Berendsen barostat for equilibration (Berendsen et al., 1998; Parrinello and Rahman 1998). Long-range electrostatic interactions were calculated using particle mesh Ewald (PME) algorithm with a real space cut-off of 1.0 nm (Darden et al., 1998). LJ interactions were truncated at 1.0 nm. The TIP3P model was used to represent the water molecules, and the LINCS algorithm was used to constrain the motion of hydrogen atoms bonded to heavy atoms (Bekker et al., 1997). The system was subjected to energy minimization to prevent any overlap of atoms, followed by 0.5 ns of equilibration and a 25-ns production run. During simulations, the ribosomal RNA, proteins and the mRNA




(except the codon) were held in place using position restraints on the heavy atoms of the RNA and protein backbone with a force constant of 1,000 N/nm in each spatial dimension for the simulation. Coordinates of the ribosomal fragment (rRNA, tRNA, and mRNA) were stored every 1 ps for further analysis. The simulations were visualized using Visual Molecular Dynamics software and analyzed using tools from Gromacs (Humphrey et al., 1996; Abraham et al., 2015).

RESULTS

The NIT-1 cell line (ATCC[®] CRL-2055[™]) is inducible for insulin production with high glucose concentrations, theophylline and KCl (**Figure 2A**). The cells were stimulated to produce insulin with Theophylline (10 mM with 5.5 mM glucose) or KCl (40 mM) in SF media (**Figure 2A**). In order to determine the effect of a reduced Cdkal1 protein production, we knocked down *cdk1* gene expression with two distinct methods. The *cdk1* gene was knocked down by transfection of an esiRNA (endonuclease-prepared siRNA, 30, 50 or 70 nM using Lipofectamine, LF; Sigma) and by a Lentiviral shRNA (Sigma). Cells were also transfected with a control GFP esiRNA in antibiotic-free medium (6 h). Cells were glucose starved overnight in SF-DMEM, and induced for insulin (90 min) or left unstimulated (SF-DMEM with 600 KIU/ml aprotinin). Cells transfected with a control GFP esiRNA or Lipofectamine (LF


RNAiMax reagent) alone responded to stimulation similar to cells that had not been transfected. Supernatants of *cdk1* esiRNA (50 nM) knockdown cells assayed for insulin by ELISA had as low as 20% of the mature insulin production compared to a control esiRNA and LF-transfected cultures (**Figure 2B**). The cells were lysed in RIPA buffer with protease inhibitors and the lysates were processed for Western blots that showed decreased Cdkal1 protein as well as decreased insulin production (**Supplemental Figure S1**).


Cells were also infected with four different Lentiviral shRNA constructs against the *cdk1* gene in antibiotic-free medium and selected with puromycin (3.5 μ g). Clones, LC4, LC5, and LC7 that survived puromycin selection were expanded and DNA isolated and sequenced. Insulin induced cells were analyzed with the GSIR assay, ELISA and Western blot. Compared to normal NIT-1 cells, the secreted insulin levels were 50% or lower with stimulation by Theophylline and KCl (**Figure 2C**). When cell lysates were assayed for insulin, LC4 stimulated by KCl showed that the knockdown of *cdk1* by the shRNA construct inhibited insulin production by >60% (**Figure 2D**). Analysis of stimulated NIT-1 cell lysates by Western blotting showed no significant changes in the levels of Cdkal1 protein in normal NIT-1 cells although knockdown cells had lower Cdkal1 protein levels (**Supplemental Figure S1**). The amount of GAPDH control protein was not affected in the Cdkal1 deficient cells.



The function of *Cdk1* protein's modification of tRNA is accurate and efficient translation of AAG/AAA codons. Insulin mRNA requires the *Cdk1* modification of tRNA^{Lys3} from t⁶A₃₇ to ms²t⁶A₃₇ for insertion of lysine at position 88. A significant decrease was observed in ms²t⁶A relative to t⁶A in RNA (<200 nts) isolated from cells that had been knocked down (Figure 3). We isolated RNA from esiRNA knockdown cells, stimulated and unstimulated for insulin production and control Lipofectamine only (LF) cells. The RNA was fractionated so that we could analyze small RNAs less than 200 nucleotides such as tRNAs without rRNA present (Ambion mirVana miRNA Isolation Kit). The RNA was hydrolyzed to nucleosides and the modified nucleoside analysis conducted by UHPLC-MS/MS (triple quadrupole MS (Waters MS)). The ms²t⁶A modification decreased in tRNAs from esiRNA30 and esiRNA50 knockdown cells. As expected, the decrease was most dramatic in tRNA from esiRNA50 cells grown in SF medium, and less so when the cells were stimulated by KCl. The decrease of ms²t⁶A in tRNA is consistent with and probably the cause of a decrease in secreted insulin and increase in precursor insulin mRNA.

Large sequence RNA was isolated from normal and transfected knockdown NIT-1 cells (Qiagen RNeasy Plus kit) and rRNA was removed (Qiagen RNeasy MinElute Kit). Quantitative real-time PCR was conducted to determine the presence of *cdk1*, precursor insulin, and mature insulin mRNAs. Threshold Cq values were normalized to actin levels. Relative expression was calculated using the 2^{ΔΔCq} method. The expression of *cdk1* mRNA levels increased by ~60% in normal NIT-1 cells





following stimulation with KCl (Figure 4A). However, in stimulated *cdk11*-esiRNA treated knockdown cells, the mRNA levels were decreased by ~30% (Figure 4B). To determine the regulation of insulin transcription, we applied a unique strategy using two sets of primers that amplified total mouse insulin I and II mRNA (fully processed mature insulin mRNA) and a precursor mRNA species containing intron 2 (Figure 4C). The levels of mature mRNA and precursor mRNA insulin levels were quantitated and expressed as the real time threshold cycle (C_T) values, in untreated (SF) and KCl stimulated NIT-1 normal or knockdown cells. Although no differences were seen in mature insulin mRNA levels, significant changes in precursor insulin mRNA levels were detected with stimulation to produce insulin (Figure 4C). When the abundance of precursor insulin mRNA relative to mature insulin mRNA in control cells is normalized to 1.00, in stimulated *cdk11* knockdown cells the ratio is 1.95 relative to mature insulin mRNA. In β -cells when the *cdk11* gene is non-functional or missing and the tRNA^{Lys3} modification ms²t⁶A has decreased, *cdk11* mRNA has decreased 30% also. However, the insulin precursor mRNA is significantly increased. Thus, *cdk11* knockdown cells stimulated to produce insulin are yet secreting less mature insulin. Lysates from these cells exhibited significantly decreased insulin and proinsulin (Figure 2D).

We asked why should the modification ms²-play such an important role in tRNA^{Lys3} in translating the lysine wobble codon AAG in insulin mRNA? When the anticodon U₃₄U₃₅U₃₆ with the adjacent ms²t⁶A₃₇ binds the wobble codon G3A2A1, the ms²t⁶A₃₇ is three nucleosides distant from the U₃₄:G₃ pair. There are three posttranscriptional modifications in the anticodon stem and loop (ASL) of tRNA^{Lys3}, 5-methoxycarbonylmethyl-2-thiouridine at wobble position 34 (mcm⁵s²U₃₄), 2-methylthio-N⁶-threonylcarbamoyladenosine at position 37 (ms²t⁶A₃₇) adjacent to the anticodon and pseudouridine (Ψ ₃₉) at position 39 in the stem. The fully modified ms²t⁶A₃₇ and mcm⁵s²U₃₄ are required to achieve wild-type binding activity of human tRNA^{Lys3} to AAA and the wobble codon AAG (Yarian et al., 2000). NMR structure

determination and molecular dynamics simulations (MDS) of the ASL demonstrated that the ms²t⁶-modification of A₃₇ supports the anticodon nucleoside stack 5' to 3' and reduces solvent accessibility of U₃₆ (Stuart et al., 2000; McCrate et al., 2006).

To explore the role of the tRNA^{Lys3} modifications at A₃₇ for recognition and decoding, we performed molecular simulations of the anticodon stem-loop of the tRNA (ASL) bound to the mRNA AAG at the A site of the eukaryotic ribosome. We compared three simulations with the ASL-mcm⁵s²U₃₄ with A₃₇, t⁶A₃₇, and ms²t⁶A₃₇ each bound to the wobble codon AAG on the ribosome. First, we considered the effect of the modifications on the codon-anticodon interaction. We compared the hydrogen bonding between the codon and anticodon nucleosides (A₁:U₃₆, A₂:U₃₅ and G₃:mcm⁵s²U₃₄), in the three systems (Figure 5). Interestingly, we observe that the hydrogen bonding is stronger for all three positions of the codon-anticodon base pairs for the addition of the t⁶-modification to A₃₇ and is further enhanced by the addition of the ms²- to t⁶A₃₇. Remarkably, we find that this enhancement is most pronounced when the mcm⁵s²U₃₄:G₃ base-pair is considered, which is the farthest from the A₃₇.

Next, we asked how does the modification at A₃₇ lead to significant strengthening of codon anticodon base-pairing? The dominant locations of the threonylcarbamoyl-group in both t⁶A₃₇ & ms²t⁶A₃₇ systems has the hydrophilic moieties of the modification (carboxyl and hydroxyl groups) either pointing away from the ASL cavity and remaining well hydrated or are involved in a cross-loop interaction with the backbone (2' hydroxyl group) of C₃₂ (Figure 6A). The rest of the modification fits inside the ASL cavity through hydrophobic and hydrogen bonding interactions, thereby offering stability to the neighboring codon-anticodon base-pairs. Furthermore, we observed transient interaction between the terminal methyl groups of ms²t⁶A₃₇ and mcm⁵s²U₃₄ (Figure 6B), suggesting that the enhancement in stability due to the modification at A₃₇ extends to the codon-anticodon base pair farthest from A₃₇. Most interestingly, we also found that the t⁶-group interaction




FIGURE 6 | The dominant orientation of the hypermodified ms²t⁶A₃₇ shown in space filling model interacting with the anticodon stem loop cavity. **(A)** The terminal carboxyl group of the threonylcarbamoyl-group faces away from the ASL cavity, while the aliphatic carbons and the terminal methyl group fill up the cavity. Stacking between A1, A₃₇ and A₃₈ is enhanced by the addition of the thiomethyl-group to t⁶A₃₇, ms²t⁶A₃₇. **(B)** The interaction between hypermodified ms²t⁶A₃₇ and mcm⁵s²U₃₄ shown in space filling model. Hydrophobic contacts between the two modified nucleotides assists mcm⁵s²U₃₄ in maintaining the anticodon:codon interaction at the wobble position.

with the ASL cavity is more stable with the addition of the ms²-group (see Movie, t⁶A₃₇ in pink and ms²t⁶A₃₇ in yellow). The ms²-group boosts the stacking interaction between A₃₇ and the A1 codon, as a result of which the threonylcarbamoyl-group is held steady in the ASL cavity.

Overall, our molecular simulations reveal a cascading mechanism for ms²t⁶A₃₇, in which hydrogen bonding energy and the hydrophobic interactions of base-stacking by the methylthio-group stabilizes the threonylcarbamoyl-group in the ASL cavity. This in turn facilitates the hydrophobic interaction of the threonylcarbamoyl-group with the methylcarboxymethyl- (mcm⁵-) group on U₃₄ three nucleosides away, stabilizing the codon-anticodon base-pairing at the wobble position for wobble codon AAG recognition.

DISCUSSION

Homozygous recessive mutations in the human *cdk11* gene such as SNPs in intron-5 are a significant risk for T2DM. Several population studies showed the significant role of SNPs in the development of T2DM in different races, but the overall available data is not sufficient to reveal the biochemical role of particular SNPs to their concerned roles towards the development of diabetes. A *cdk11*-deficient mouse presents properties characteristic of human T2DM (Wei et al., 2011). In this study, we investigated the consequences of diminished function of Cdk11 in pancreatic islets cell NIT-1 cell lines and uncovered a biochemical connection of the tRNA modification enzyme to insulin translation and processing. SNPs in intron-5 of *cdk11* could cause alternative mRNA splicing (Zhou et al., 2014) and a reduction of Cdk11 protein synthesis. Here, diminished function of Cdk11 by knockdown of the *cdk11*

gene in mouse NIT-1 cells resulted in not only the reduction of tRNA^{Lys3} modification, but also a decrease in insulin mRNA and mature insulin. Using MDS we were able to determine that the tRNA^{Lys3} lacked hydrogen bonds and stacking to the wobble codon AAG when missing the ms²-modification of t⁶A₃₇. Generation of *cdk11* knockout mouse (Shao et al., 2016) and availability of knockout cell lines increases the possible ways to study the quite challenging SNPs in genetic level to reveal its functional roles in mature insulin production and availability.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

PA conceived and funded the presented idea. AN designed and conducted experiments, analyzed the results and wrote the first draft of the paper. EE, MC, OA, FC, and BL conducted experiments under the supervision of AN and KS. SV and SR conducted and interpreted the MDS experiments. SR and PA verified the methods, results, and wrote the manuscript. All authors discussed the results and contributed to the final manuscript.

FUNDING

Research was supported by the Department of Defense Grant Number W81XWH-16-1-0428 to PFA.

ACKNOWLEDGMENTS

The authors wish to acknowledge Haein Kim in culturing cells and advising for the manuscript, Dr. Prashanth Rangan (The RNA Institute, SUNY) for aiding in the molecular genetics of *Drosophila*, and The RNA Institute and Jennifer Montimurro in facilitating the research.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmolb.2020.584228/full#supplementary-material>.

Supplementary Figure S1. Insulin B protein and Cdkal1 protein in NIT-1 knockdown cell lysates. Western blots were employed to identify levels of insulin, Cdkal1 and control GAPDH proteins in

Control esiRNA (70 nM) and *cdk11* esiRNA knockdown cells unstimulated (Serum Free, SF) and stimulated (KCl) to produce insulin. **A.** Cdkal1 levels in control LF and GFP esiRNA cells stimulated and not stimulated to produce insulin and in *cdk11* esiRNA (70 nM) knockdown cells. **B.** Insulin B protein levels in control LF and GFP esiRNA-treated cells stimulated and not stimulated to produce insulin and in *cdk11* esiRNA (70 nM) knockdown cells. **C.** Mouse antibodies were used in Western blots against Insulin B and Cdkal1 or GAPDH. Insulin production was induced by low and high glucose, Tolbutamide, Theophyllin and KCl. Protein levels in cell lysates in Western blots were quantitated with Image J software and normalized to GAPDH levels and expressed as fold-change from control unstimulated (SF) cell lysates. LF – Lipofectamine (1ul); Control GFP-esiRNA 70 nM; *cdk11*-esiRNA 70 nM. Protein (250 µg) was loaded on SDS-PAGE gels and immune blotted using mouse antibodies against Insulin B (Clone C-12, Santa Cruz), Cdkal1 (clone E9, Santa Cruz), or GAPDH (Clone MA5-GAIR, Thermo Fisher). GAPDH was used as a loading control.

REFERENCES

- Abraham, M. J., Murtola, T., Schulz, R., Páll, S., Smith, J. C., Hess, B., et al. (2015). GROMACS: high performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* 1–2, 19–25. doi:10.1016/j.softx.2015.06.001
- Adami, R., and Bottai, D. (2020). S-adenosylmethionine tRNA modification: unexpected/unsuspected implications of former/new players. *Int. J. Biol. Sci.* 16 (15), 3018–3027. doi:10.7150/ijbs.49302
- Agris, P. F., Narendran, A., Sarachan, K., Våre, V. Y. P., and Eruysal, E. (2017). The Importance of Being Modified: the role of RNA modifications in translational fidelity. *Enzymes* 41, 1–50. doi:10.1016/bs.enz.2017.03.005
- Arragain, S., Handelman, S. K., Forouhar, F., Wei, F. Y., Tomizawa, K., Hunt, J. F., et al. (2010). Identification of eukaryotic and prokaryotic methylthiotransferase for biosynthesis of 2-methylthio-N⁶-threonylcarbamoyladenosine in tRNA. *J. Biol. Chem.* 285, 28425–28433. doi:10.1074/jbc.M110.106831
- Basanta-Sanchez, M., Temple, S., Ansari, S. A., D'Amico, A., and Agris, P. F. (2016). Attomole quantification and global profile of RNA modifications: Epitranscriptome of human neural stem cells. *Nucl. Acids Res.* 44, e26. doi:10.1093/nar/gkv971
- Bekker, H., Berendsen, H. J. C., and Fraaije, J. G. E. M. (1997). LINC: a linear constraint solver for molecular simulations. *J. Comput. Chem.* 18, 1463–1472. doi:10.1002/(sici)1096-987x(19971130)18:15<1930::aid-jcc8>3.0.co;2-p
- Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A., and Haak, J. R. (1998). Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* 81, 3684–3690.
- Brambilla, S., Altkrueger, A., Colombo, S. F., Friederich, A., Eickelmann, P., Mark, M., et al. (2012). CDK5 regulatory subunit-associated protein 1-like 1 (CDKAL1) is a tail-anchored protein in the endoplasmic reticulum (ER) of insulinoma cells. *J. Biol. Chem.* 287, 41808–41819. doi:10.1074/jbc.M112.376558
- Bussi, G., Donadio, D., and Parrinello, M. (2007). Canonical sampling through velocity rescaling. *J. Chem. Phys.* 126, 14101. doi:10.1063/1.2408420
- Centers for Disease Control and Prevention. (2020). *National Diabetes Statistics Report, 2020*. Atlanta, GA: Centers for Disease Control and Prevention, U.S. Dept of Health and Human Services.
- Chemical Computing Group. (2019). *ULC MOE (The Molecular Operating Environment) Version 2019.0102, 1010 Sherbooke St. West Suite #910, Montreal, QC, Canada, H3A 2R7*.
- Chen, A. A., and García, A. E. (2013). High-resolution reversible folding of hyperstable RNA tetraloops using molecular dynamics simulations. *Proc. Natl. Acad. Sci. USA* 110, 16820–16825. doi:10.1073/pnas.1309392110
- Cho, S., Kim, S. C., and Chung, M. G. (2019). Identification of novel population clusters with different susceptibilities to type 2 diabetes and their impact on the prediction of diabetes. *Sci. Rep.* 9, 3329–3338. doi:10.1038/s41598-019-40058-y
- Cornell, W. D., Cieplak, P., Bayly, C. I., Gould, I. R., Merz, K. M., Ferguson, D. M., et al. (1995). A second generation force field for the simulation of proteins, nucleic acids, and organic molecules. *J. Am. Chem. Soc.* 117, 5179–5197. doi:10.1021/ja00124a002
- Cornell, W. D., Cieplak, P., Bayly, C. I., and Kollman, P. A. (1993). Application of RESP charges to calculate conformational energies, hydrogen bond energies, and free energies of solvation. *J. Am. Chem. Soc.* 115, 9620–9631. doi:10.1021/ja00074a030
- Darden, T., York, D., and Pedersen, L. (1998). Particle mesh Ewald: an N-log(N) method for Ewald sums in large systems. *J. Chem. Phys.* 98, 10089–10092.
- Diabetes Genetics Initiative of Broad Institute of HarvardSaxena, R., Voight, B. F., Lyssenko, V., Burt, N. P., de Bakker, P. I., et al. (2007). Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* 316, 1331–1336. doi:10.1126/science.1142358
- Dupradeau, F. Y., Pigache, A., Zaffran, T., Savineau, C., Lelong, R., Grivel, N., et al. (2010). The R.E.D. tools: advances in RESP and ESP charge derivation and force field library building. *Phys. Chem. Chem. Phys.* 12, 7821–7839. doi:10.1039/c0cp00111b
- Forouhar, F., Arragain, S., Atta, M., Gambarelli, S., Mouesca, J. M., Hussain, M., et al. (2013). Two Fe-S clusters catalyze sulfur insertion by radical-SAM methylthiotransferases. *Nat. Chem. Biol.* 9, 333–338. doi:10.1038/nchembio.1229
- Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: visual molecular dynamics. *J. Mol. Graph.* 14, 33–38. doi:10.1016/0263-7855(96)00018-5
- Iype, T., Francis, J., Garmey, J. C., Schisler, J. C., Neshler, R., Weir, G. C., et al. (2005). Mechanism of insulin gene regulation by the pancreatic transcription factor Pdx-1: application of pre-mRNA analysis and chromatin immunoprecipitation to assess formation of functional transcriptional complexes. *J. Biol. Chem.* 280, 16798–16807. doi:10.1074/jbc.M414381200
- Krentz, N. A. J., and Gloyn, A. L. (2020). Insights into pancreatic islet cell dysfunction from type 2 diabetes mellitus genetics. *Nat. Rev. Endocrinol.* 16, 202–212. doi:10.1038/s41574-020-0325-0
- Landgraf, B. J., McCarthy, E. L., and Booker, S. J. (2016). Radical S-Adenosylmethionine enzymes in human health and disease. *Annu. Rev. Biochem.* 85, 485–514. doi:10.1146/annurev-biochem-060713-035504
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25, 402. doi:10.1006/meth.2001.1262

- Mannino, G. C., Andreozzi, F., and Sesti, G. (2019). Pharmacogenetics of type 2 diabetes mellitus, the route toward tailored medicine. *Diab. Metab. Res. Rev.* 35, e3109. doi:10.1002/dmrr.3109
- McCown, P. J., Ruszkowska, A., Kunkler, C. N., Breger, K., Hulewicz, J. P., Wang, M. C., et al. (2020). Naturally occurring modified ribonucleosides. *Wiley Interdiscip. Rev. RNA* 11 (5), e1595. doi:10.1002/wrna.1595
- McCrate, N. E., Varner, M. E., Kim, K. I., and Nagan, M. C. (2006). Molecular dynamics simulations of human Formula: the role of modified bases in mRNA recognition. *Nucl. Acids Res.* 34 (19), 5361–5368. doi:10.1093/nar/gkl580
- Parrinello, M., and Rahman, A. (1998). Polymorphic transitions in single crystals: a new molecular dynamics method. *J. Appl. Phys.* 52, 7182–7190.
- Santos, M. C. F. D., Anderson, C. P., Neschen, S., Zumbrennen-Bullough, K. B., Romney, S. J., Kahle-Stephan, M., et al. (2020). Irf2 regulates insulin production through iron-mediated Cdkal1-catalyzed tRNA modification. *Nat. Commun.* 11, 296. doi:10.1038/s41467-019-14004-5
- Schmidt, J. R., and Polik, W. F. (2020). *WebMO Enterprise*. Holland, MI, USA: WebMO LLC (Accessed March, 20). <https://www.webmo.net>. version.20.
- Shao, S., Murray, J., Brown, A., Taunton, J., Ramakrishnan, V., and Hegde, R. S. (2016). Decoding mammalian ribosome-mRNA states by translational GTPase complexes. *Cell* 167, 1229–e15. doi:10.1016/j.cell.2016.10.046
- Steinthorsdottir, V., Thorleifsson, G., Reynisdottir, I., Benediktsson, R., Jonsdottir, T., Walters, G. B., et al. (2007). A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. *Nat. Genet.* 39, 770–775. doi:10.1038/ng2043
- Stuart, J. W., Gdaniec, Z., Guenther, R., Marszalek, M., Sochacka, E., Malkiewicz, A., et al. (2000). Functional anticodon architecture of human tRNALys3 includes disruption of intraloop hydrogen bonding by the naturally occurring amino acid modification, t6A. *Biochemistry* 39, 13396–13404. doi:10.1021/bi0013039
- Wei, F. Y., Suzuki, T., Watanabe, S., Kimura, S., Kaitsuka, T., Fujimura, A., et al. (2011). Deficit of tRNA(Lys) modification by Cdkal1 causes the development of type 2 diabetes in mice. *J. Clin. Invest.* 121 (9), 3598–3608. doi:10.1172/JCI58056
- Witka, B. Z., Oktaviani, D. J., Marcellino, M., Barliana, M. I., and Abdulah, R. (2019). Type 2 Diabetes-associated genetic polymorphisms as potential disease predictors. *Diab. Metab. Syndr. Obes.* 12, 2689–2706. doi:10.2147/DMSO.S230061
- Yarian, C., Marszalek, M., Sochacka, E., Malkiewicz, A., Guenther, R., Miskiewicz, A., et al. (2000). Modified nucleoside dependent Watson–Crick and wobble codon binding by tRNALysUUUSpecies†. *Biochemistry* 39, 13390–13395. doi:10.1021/bi001302g
- Zhou, B., Wei, F. Y., Kanai, N., Fujimura, A., Kaitsuka, T., and Tomizawa, K. (2014). Identification of a splicing variant that regulates type 2 diabetes risk factor CDKAL1 level by a coding-independent mechanism in human. *Hum. Mol. Genet.* 23, 4639–4650. doi:10.1093/hmg/ddu184

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Narendran, Vangaveti, Ranganathan, Eruysal, Craft, Alrifai, Chua, Sarachan, Litwa, Ramachandran and Agris. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.