



Recent Developments in Data-Assisted Modeling of Flexible Proteins

Cezary Czaplewski¹, Zhou Gong², Emilia A. Lubecka³, Kai Xue⁴, Chun Tang^{4*} and Adam Liwo^{1*}

¹Faculty of Chemistry, University of Gdańsk, Gdańsk, Poland, ²Innovation Academy for Precision Measurement Science and Technology, Chinese Academy of Sciences, Wuhan, China, ³Faculty of Electronics, Telecommunications and Informatics, Gdańsk University of Technology, Gdańsk, Poland, ⁴PKU-Tsinghua Center for Life Sciences, Beijing National Laboratory for Molecular Sciences, College of Chemistry and Molecular Engineering, Peking University, Beijing, China

OPEN ACCESS

Edited by:

Nikolaos G. Sgourakis,
University of Pennsylvania,
United States

Reviewed by:

Robert Oliver Schneider,
Lille University of Science and
Technology, France
Yong Wang,
Zhejiang University, China

*Correspondence:

Adam Liwo
adam.liwo@ug.edu.pl
Chun Tang
Tang_Chun@pku.edu.cn

Specialty section:

This article was submitted to
Structural Biology,
a section of the journal
Frontiers in Molecular Biosciences

Received: 27 August 2021

Accepted: 06 December 2021

Published: 24 December 2021

Citation:

Czaplewski C, Gong Z, Lubecka EA,
Xue K, Tang C and Liwo A (2021)
Recent Developments in Data-
Assisted Modeling of Flexible Proteins.
Front. Mol. Biosci. 8:765562.
doi: 10.3389/fmolb.2021.765562


Many proteins can fold into well-defined conformations. However, intrinsically-disordered proteins (IDPs) do not possess a defined structure. Moreover, folded multi-domain proteins often digress into alternative conformations. Collectively, the conformational dynamics enables these proteins to fulfill specific functions. Thus, most experimental observables are averaged over the conformations that constitute an ensemble. In this article, we review the recent developments in the concept and methods for the determination of the dynamic structures of flexible peptides and proteins. In particular, we describe ways to extract information from nuclear magnetic resonance small-angle X-ray scattering (SAXS), and chemical cross-linking coupled with mass spectroscopy (XL-MS) measurements. All these techniques can be used to obtain ensemble-averaged restraints or to re-weight the simulated conformational ensembles.

Keywords: proteins, data-assisted modeling, conformational ensembles, nuclear magnetic resonance, small-angle X-ray scattering, chemical cross-linking coupled with mass spectroscopy, molecular dynamics, coarse graining

1 INTRODUCTION

Proteins exist as dynamic structures. Many proteins undergo often very significant motions while performing their functions (Henzler-Wildman and Kern, 2007; Boehr et al., 2009). The respective conformational states are sometimes stable enough to be captured through X-ray structure determination if appropriate conditions of protein-sample preparation are applied (Bertelsen et al., 2009; Kityk et al., 2012). Nevertheless, in most instances, the structures of multistate proteins, as well as those of intrinsically disordered proteins (IDPs) or proteins with intrinsically-disordered regions (IDRs) can be described only in terms of conformational ensembles. Over 40% of human proteins contain stretches of disorder longer than 30 residues (van der Lee et al., 2014).

Thus, ensemble-averaged quantities are usually obtained from measurements while studying conformational dynamics of multistate proteins, IDPs, or flexible peptides. The composition of an ensemble can be determined only by combining the results of measurements with advanced molecular modeling (Bonomi et al., 2017; Bonomi and Vendruscolo, 2019; Orioli et al., 2020). In this minireview, we summarize the methods for conformational-ensemble determination using molecular modeling, using the data from nuclear magnetic resonance (NMR), small-angle X-ray scattering (SAXS), and chemical cross-linking coupled with mass spectroscopy (XL-MS). In **Section 2**, we outline the experimental techniques mentioned above and the quantities that they provide,



while in **Section 3** we describe conformational-sampling methods and two major approaches of implementing the experimental quantities in conformational-ensemble determination: simulations with ensemble-averaged restraints and ensemble reweighting. A scheme summarizing the methodologies discussed is shown in **Figure 1**.

2 EXPERIMENTAL METHODS TO STUDY FLEXIBLE PROTEINS

Here we focus on the experimental measurements that can be performed for proteins in solution. We leave out the single-molecule fluorescence resonance energy transfer (FRET), which does not yield ensemble averages and does not have the same issues as those discussed in **Section 3** (Tang and Gong, 2020; Lerner et al., 2021).

2.1 Nuclear Magnetic Resonance

The most complete information about the structure and conformational dynamics of proteins and peptides is provided by NMR (Sekhar and Kay, 2019). NMR remains the method of choice to characterize the conformational dynamics of proteins to atomic resolution in near-physiological conditions. NMR observables, including nuclear Overhauser effect (NOE), chemical shift, dipolar coupling constants, and paramagnetic relaxation enhancement (PRE) are ensemble-averaged over a multitude of conformational states (Salmon et al., 2010; Konrat, 2014; Clore, 2015; Huang et al., 2015; Tang and Gong, 2020). Thus, though the flexible regions in a protein can be easily identified by NMR owing to their favorable relaxation properties, it is difficult to obtain a comprehensive description of the ensemble structure of a multi-domain protein or an IDP as a whole and determine the fractions of the constituting conformational states. To this end, many methods have been

developed to reconstruct the ensembles based on the NMR data (Bertini et al., 2004; Mittag and Forman-Kay, 2007; Delaforge et al., 2015).

Paramagnetic NMR, in particular, paramagnetic relaxation enhancement (PRE), allows the visualization of protein ensemble structures (Otting, 2010; Liu et al., 2016). The PRE is exquisitely sensitive to the sparsely populated conformations, thanks to the large gyromagnetic ratio of an unpaired electron in the paramagnetic probe and an inverse sixth power dependence on the distances to the observed NMR nuclei (Clore, 2015; Liu et al., 2015). On the other hand, covalent attachment of a paramagnetic probe could perturb the structure, which is more likely for an IDP (Sasmal et al., 2017). As a result, paramagnetic cosolute molecules have been developed (Gu et al., 2014; Gong et al., 2017a), which can also be used to assess the dynamic structures of IDPs (Hartlmüller et al., 2019; Spreitzer et al., 2020). Similar to the PREs, the NOEs also provide ensemble-averaged distances between protein nuclei. However, quantitative interpretation of the NOEs is hampered by the complex relaxation pathways. The exact proton-proton distances and the corresponding conformational states of a protein are best extracted on a perdeuterated background (Vögeli et al., 2009; Vögeli et al., 2016).

2.2 Small-Angle Scattering Methods

Compared to NMR, small-angle X-ray and small-angle neutron scattering (SANS) provide less detailed but more global structural information (Konarev et al., 2003; Forster et al., 2008; Schneidman-Duhovny et al., 2011; Trewella et al., 2013). For a multi-state protein, the scattering curve is averaged over a multitude of conformational states. The different states and the associated population can, in theory, be obtained from the deconvolution of the scattering curve. To this end, many algorithms have been developed that include ensemble optimization method (EOM) (Bernadó et al., 2007; Tria et al., 2015), minimal ensemble search (MES) (Pelikan et al., 2009), and Bayesian ensemble SAXS (BE-SAXS) (Antonov et al., 2016). Though the scattering intensity at each scattering angle is normally used as a restraint (Forster et al., 2008), pairwise distance distribution could also be employed for the comparison between different sets of structure ensembles (Gorba et al., 2008; Karczyńska et al., 2018). The different approaches fit different numbers of parameters and use different treatments of the displaced solvent, which inevitably leads to somewhat different solutions.

2.3 Chemical Cross-Linking Coupled With Mass Spectrometry

Cross-linking reactions are initiated either by illumination or chemical reaction followed by enzymatic digestion. The final products are cross-linked peptides, which can be identified by mass spectrometry with high confidence. The cross-linked residues have to be closer in distance than the length of the cross-linker arm. Therefore, each cross-link can be used to derive the restraint imposed on the $C^\alpha \dots, C^\alpha$, $C^\beta \dots, C^\beta$ - or the terminal-atom (e.g., $N^\zeta \dots, N^\zeta$ atom pair of lysine side chains)

distance of the two cross-linked residues. However, the cross-links may artificially pull two protein regions together, in a so-called zipper effect (Belsom and Rappsilber, 2021), which needs to be carefully controlled and ruled out.

The identified cross-links are often found incompatible with the known protein structure, in which the calculated distance exceeds the maximum length of the cross-linker. Such “over-length” cross-links can be explained by alternative protein conformations, e.g., an open-to-closed transition (Ding et al., 2017), or by the transient oligomerization of the protein. The latter can be ascertained with the mixing of “light” and “heavy” proteins with distinct isotope labeling patterns (Gong et al., 2015). Furthermore, cross-linking mass spectrometry (XL-MS) can be used to elucidate dynamic encounters between two proteins (Gong et al., 2017b).

A crosslink restraint is usually imposed on the straight-line distance between the C^α-atoms of the corresponding residues (Leitner et al., 2014; Merkle et al., 2014; Fajardo et al., 2019). Recently, we developed an approach in which restraints are imposed on side-chain ends and implemented it in all-atom (Gong et al., 2020) and coarse-grained (Kogut et al., 2021) molecular dynamics. This approach is more realistic because such distances are close to those between the solvent-accessible surfaces, which are targeted by the cross-linking reagents in the XL-MS experiments.

3 MODELING PROTEIN STRUCTURES WITH EXPERIMENTAL RESTRAINTS

3.1 Conformational Search

Canonical molecular dynamics (MD) (Frenkel and Smit, 2000) and its extensions, namely simulated annealing (SA) (Kirkpatrick et al., 1983), replica-exchange molecular dynamics (REMD) (Hansmann, 1997), and multiplexed replica exchange molecular dynamics (MREMD) (Rhee and Pande, 2003) are usually the methods of choice for sampling the conformational space, owing to their efficiency. All-atom MD is commonly used and a variety of good algorithms and software packages such as e.g., AMBER (Salomon-Ferrer et al., 2013), CHARMM (Brooks et al., 2009), GROMACS (Abraham et al., 2015), LAMMPS (Plimpton, 1995) and DESMOND (Bowers et al., 2006) are available, which also enable the researchers to include experimental information as restraints.

All-atom MD has restricted ability to sample the conformational space extensively (Bottaro and Lindorff-Larsen, 2018). Compared to all-atom approaches, the coarse-grained (CG) approaches, in which several atoms are merged into extended interaction sites, are computationally more efficient and enable us to run simulations at much longer time-scales and for larger systems (Voth, 2008; Kmiecik et al., 2016). The coarse-grained models with which MD for proteins can be run include MARTINI (Marrink and Tieleman, 2013), AWSEM (Davtyan et al., 2012), OPEP (Sterpone et al., 2014), and UNRES (Liwo et al., 2019). CABS (Kolinski, 2004) is another very good CG model of proteins, which was developed to run Monte Carlo dynamics on a high-resolution lattice.

The experimental information can be used as restraints or to filter the conformational ensembles/reweight its conformations to reproduce the experimental observables (Bonomi et al., 2017; Orioli et al., 2020). These two approaches are described in the two subsequent subsections.

3.2 Restrained Simulations of Conformationally Heterogeneous Systems

In restrained simulations, penalty terms are added to the potential energy in MD so that the forces consist of the forces computed from the force field of choice and those due to restraint violation (van Gunsteren et al., 2016). This approach is straightforward if a protein has a well-defined structure and has been implemented in the CYANA (Güntert and Buchner, 2015) and XPLOR-NIH software packages (Schwieters et al., 2018) for structure determination by NMR, as well as is built in the MD packages mentioned in the previous section. For flexible systems, time- and ensemble averaging algorithms to run restrained simulations have been developed.

It should be noted that using restraints from NMR in CG simulations is not straightforward, because the respective quantities depend on all-atom geometry. One method, in which the CG structures are converted into all-atom structures, from which the respective quantities are calculated, was developed (Latek and Koliński, 2011) for use with the CABS model of proteins (Kolinski, 2004). However, this method is not suitable for restrained MD simulations, because it does not provide the forces due to restraints. Recently, we developed ESCASA (Lubecka and Liwo, 2021), an analytical approach to calculating approximate positions of the protons from C^α-trace geometry, thus enabling us to compute the forces due to the penalty function and, consequently, to use the method with coarse-grained MD.

3.2.1 Time-Averaged Restraints

In the time-averaged-restraint method, the quantities obtained from simulations (e.g., interproton distances) are averaged over a time window (Torda et al., 1989; Bonvin et al., 1994). These average quantities are inserted into the penalty terms.

$$\bar{f}(\mathbf{r}(t)) = [\tau(1 - \exp(-t/\tau))]^{-1} \int_0^t \exp(-t'/\tau) f(\mathbf{r}(t-t')) dt' \quad (1)$$

where f is the quantity being averaged, which depends on the coordinates of the atoms of the system contained in vector \mathbf{r} and τ is the length of the time window.

3.2.2 Ensemble Averaged Restraints

The methods that use ensemble-averaged restraints are based on the maximum-entropy and Bayesian principles, according to which a minimally perturbed conformational ensemble compared to that resulting from free simulations is sought and, at the same time, the ensemble-average quantities match their experimental counterparts within the experimental error

(Pitera and Chodera, 2012; White et al., 2015; Amirkulova and While, 2019). If the ensemble-averaged restraints are enforced strictly, the potential-energy function is modified to include the experimental quantities with the weight calculated to maximize the entropy (Pitera and Chodera, 2012).

$$U_{ME}(\mathbf{r}; \alpha_1, \dots, \alpha_M) = U(\mathbf{r}) + \sum_{i=1}^N \alpha_i f_i(\mathbf{r}) \quad (2)$$

where $f_i(\mathbf{r})$ is the value of the i th experimental observable calculated for the conformation described by the vector of coordinates \mathbf{r} , N is the number of observables, U is the potential-energy function used in MD simulations, U_{ME} is the extended energy function and the weights α_i are calculated to minimize $\Gamma(\alpha_1, \dots, \alpha_N)$.

$$\Gamma(\alpha_1, \dots, \alpha_N) = \ln \int \exp[-\beta U_{ME}(\mathbf{r}; \alpha_1, \dots, \alpha_N)] d^{3n} \mathbf{r} - \beta \sum_{i=1}^N \alpha_i f_{i,exp} \quad (3)$$

where $f_{i,exp}$ is the experimental (ensemble-averaged) value of the i th observable, $\beta = 1/RT$, R being the universal gas constant and T absolute temperature, and n is the number of atoms in the system. It should be noted that the integral in **Equation 3** does not have to be evaluated, because minimization of Γ leads to equations which contain the observables averaged over the conformations, which can readily be calculated from MD simulations (Pitera and Chodera, 2012). With this approach, the distribution of conformations is minimally perturbed with respect to that resulting from the force field used. In other words, the experimental constraints enable us to compensate for the inevitable inaccuracy of the force field and to obtain a distribution of conformations in the ensemble, which is closer to the true (Boltzmann) distribution (Cavalli et al., 2013), provided that the experimental data are sufficient in number and quality. In practical implementation, the replica-averaged method is applied (Camilloni et al., 2013; Hummer and Köfinger, 2015), in which several replicas are run with the extended potential energy, U_{ED} , containing harmonic restraints on the experimentally measured quantities that are averaged over all replicas.

$$U_{ED}(\mathbf{r}_i) = U(\mathbf{r}_i) + M \sum_j \frac{(\frac{1}{M} \sum_{k=1}^M f_j(\mathbf{r}_k) - f_{exp,j})^2}{2\sigma_j^2} \quad (4)$$

where the index i runs over replicas, M is the number of replicas, \mathbf{r}_k is the vector of the coordinates of the conformation of the k th replica, and σ_j is the error in the j th observable. It has been demonstrated that this method becomes the maximum-entropy method as the number of replicas increases (Pitera and Chodera, 2012; Cavalli et al., 2013; Roux and Weare, 2013; Hummer and Köfinger, 2015). This approach has been implemented in determining the conformational ensembles from NMR (Camilloni et al., 2013) and SAXS data (Hermann and Hub, 2019). A similar approach termed dynamic ensemble refinement (DER) (Lindorff-Larsen et al., 2005) was developed earlier for the determination of protein dynamical ensembles from NMR data.

3.3 Reweighting the Conformational Ensembles

In the ensemble-reweighting methods, a pool of conformations is generated first in unrestrained simulations and, subsequently, the weights of the conformations are determined to reach the agreement of the conformation-averaged observables with the corresponding experimental quantities (Cavalli et al., 2013; Orioli et al., 2020). An advantage of this approach is that the ensemble can be generated once and used as the results of new experiments are available. However, the state-of-the-art force fields do not produce the true Boltzmann distribution of the conformational states. Consequently, the distribution of conformations obtained in unrestrained simulations could be far from the true distribution; specifically, some regions of conformational space that are, in reality, visited by the system might happen to be under-represented or omitted from the simulated ensemble. It has been demonstrated (Cerioti et al., 2012) that the more the input distribution diverges from the true distribution the greater the error in reweighting. When the experimental information is included in the simulations as maximum-entropy constraints or replica-averaged restraints, the ensemble is driven towards reproducing the experimental data, i.e., closer to the true (unknown) Boltzmann distribution. An example that the quality of the force field becomes less important with increasing the number of data is the work by Joo et al. (Joo et al., 2015), in which a force field that contained only the van der Waals repulsion, stereochemistry, improper-torsion, and chirality terms, in combination with NOE and dihedral-angle restraints, was used with success to determine protein structures from NMR data.

Because the number of conformations in the ensemble (and, thereby, the number of weights) is usually much greater than the number of observables, the fitting problem is underdetermined. It is solved by using either the maximum-parsimony or the maximum-entropy principle (Bonomi et al., 2017).

In the maximum-parsimony approaches, a minimum set of conformations is determined that can reproduce the experimental observables. This method was originated by Nikiforovich and coworkers (Nikiforovich et al., 1987) and, subsequently evolved into a variety of algorithms, including EOM (Bernadó et al., 2007), ASTEROIDS (Nodet et al., 2009), and SES (Berlin et al., 2013), as well as the algorithms developed in our laboratories to determine the conformational ensembles from the SAXS (Kozak et al., 2010) or SAXS, NMR and XL-MS data (Liu et al., 2018). Usually, the ensemble is clustered first and averages are computed over each cluster, the weights of the clusters being determined by least-square fitting the ensemble-averaged observables to the experimental quantities, subject to the condition that all weights are non-zero and the number of clusters with non-zero weights is minimal.

In the maximum-entropy approach, the weights of conformations are determined so that the ensemble-averaged quantities match the experimental counterparts with minimal perturbation of the input ensembles. Usually, the experimental

errors are included in the target function, which results in solving a Bayesian problem, with the prior distribution being equal to that from unrestrained MD simulations.

$$\theta \sum_{j=1}^M w_j \ln w_j + \sum_{i=1}^N \frac{(\sum_{j=1}^M w_j f_{ij} - f_{i,exp})^2}{2\sigma_i^2} = \min \quad (5)$$

where the first term is the negative of the Shannon entropy, θ being the weight of this term, and the weights are required to be normalized to unity and non-negative. Many approaches that use this principle, including ENSEMBLE (Marsh and Forman-Kay, 2012; Krzeminski et al., 2013), EROS (Różycki et al., 2011), COPER (Leung et al., 2016), and others (Groth et al., 1999) were developed.

Recently, Pesce and Lindorff-Larsen (Pesce and Lindorff-Larsen, 2021) designed an iterative maximum-entropy reweighting method for the determination of conformational ensembles from SAXS data, in which background intensity and the scaling factor of the computed average SAXS profile are fitted to match the experimental profile. Subsequently, the weights are determined by minimizing the target function of **Equation 5**. The two steps are iterated until convergence is achieved. The determination of background intensity and scaling factor is a major step forward with respect to the previous approaches, in which only the weights were determined, because these parameters depend on many features of the system studied (e.g., the solvation shell) and on experimental conditions. Also, very recently, an ensemble-reweighting method by using side-chain NMR-relaxation, termed Average Block Selection Using Relaxation Data with Entropy Restraint (ABSURDer), an extension of the ABSURD method of Blackledge and others (Salvi et al., 2016), has been developed by the Lindorff-Larsen group (Kümmerer et al., 2021). This approach takes into account system dynamics, thus enabling us to find the ensemble of trajectories, not just static conformations, consistent with experiment.

4 CONCLUSION AND OUTLOOK

Investigation of the dynamic structures of proteins and other biomolecules in solution is a rapidly growing field, in which the experimental and theoretical methods are complementary to each other (Bonomi et al., 2017; Bonomi and Vendruscolo, 2019; Orioli et al., 2020). Since the experiment provides only average observables (NMR), distance distribution (SAXS, SANS, and WAXS), or just indicates which residues may be close to each other in part of the dynamic structure (XL-MS), dynamic structure determination from the experiment alone is an underdetermined problem. Thus, the development of efficient and reliable conformational-search methods and better force fields is a necessity.

At present, the respective algorithms are based mostly on ensemble reweighting (Bonomi et al., 2017; Bonomi and Vendruscolo, 2019; Orioli et al., 2020), the maximum-entropy

variant of which seems to be better, because it does not leave out any part of the ensemble completely, an important feature given the under-determinability of the reweighting problem (Bonomi et al., 2017). Because the conformational ensemble is generated in unrestrained simulations, this approach depends on the quality of a force field used, which is usually still far from being perfect. Therefore, the development of methods based on replica-averaged restraints, which stem from the maximum-entropy principle (Cavalli et al., 2013; Hummer and Köfner, 2015) seems to be a better approach. Combining this approach with time-averaged restraints (Torda et al., 1989; Bonvin et al., 1994) or posterior ensemble fitting to enrich the averaging is recommended. An efficient conformational search is required regardless of choosing a particular method to include the experimental data, which can be carried out with coarse-grained models (Voth, 2008; Kmiecik et al., 2016). Deep-learning algorithms are also likely to advance the field, especially given their recent tremendous success in predicting the stable structures of proteins at crystallographic accuracy (Baek et al., 2021; Jumper et al., 2021). These methods may be used to generate the initial models for studying the dynamics of multistate proteins.

Another challenge is capturing the full dynamics of the system under study. Time-resolved techniques are an obvious answer here but averages, such as kinetic rate constants, can also be used – an approach has recently been proposed (Brotzakis et al., 2021). This will be particularly important when studying the dynamics of multistate proteins with more than two stable states.

AUTHOR CONTRIBUTIONS

AL and CT designed the manuscript and wrote part of the text. EL, CC, KX, and ZG participated in writing.

FUNDING

This work was supported by grants No. 2018YFA0507700 (National Key R&D Program of China), UMO-2017/25/B/ST4/01026, and UMO-2017/26/M/ST4/00044 from the National Science Center of Poland (Narodowe Centrum Nauki).

ACKNOWLEDGMENTS

Computational resources were provided by (a) the Interdisciplinary Center of Mathematical and Computer Modeling (ICM) the University of Warsaw under grant No. GA71-23, (b) the Centre of Informatics - Tricity Academic Supercomputer and Network (CI TASK) in Gdańsk, (c) the Academic Computer Centre Cyfronet AGH in Krakow under grants unres19 and unres2021 and (d) 796-processor Beowulf cluster at the Faculty of Chemistry, University of Gdańsk.

REFERENCES

- Abraham, M. J., Murtola, T., Schulz, R., Páll, S., Smith, J. C., Hess, B., et al. (2015). GROMACS: High Performance Molecular Simulations through Multi-Level Parallelism from Laptops to Supercomputers. *SoftwareX* 1-2, 19–25. doi:10.1016/j.softx.2015.06.001
- Amirkulova, D. B., and White, A. D. (2019). Recent Advances in Maximum Entropy Biasing Techniques for Molecular Dynamics. *Mol. Simul.* 45, 1285–1294. doi:10.1080/08927022.2019.1608988
- Antonov, L. D., Olsson, S., Boomsma, W., and Hamelryck, T. (2016). Bayesian Inference of Protein Ensembles from SAXS Data. *Phys. Chem. Chem. Phys.* 18, 5832–5838. doi:10.1039/c5cp04886a
- Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G. R., et al. (2021). Accurate Prediction of Protein Structures and Interactions Using a Three-Track Neural Network. *Science* 373, 871–876. doi:10.1126/science.abj8754
- Belsom, A., and Rappilber, J. (2021). Anatomy of a Crosslinker. *Curr. Opin. Chem. Biol.* 60, 39–46. doi:10.1016/j.cbpa.2020.07.008
- Berlin, K., Castañeda, C. A., Schneidman-Duhovny, D., Sali, A., Nava-Tudela, A., and Fushman, D. (2013). Recovering a Representative Conformational Ensemble from Underdetermined Macromolecular Structural Data. *J. Am. Chem. Soc.* 135, 16595–16609. doi:10.1021/ja4083717
- Bernadó, P., Mylonas, E., Petoukhov, M. V., Blackledge, M., and Svergun, D. I. (2007). Structural Characterization of Flexible Proteins Using Small-Angle X-ray Scattering. *J. Am. Chem. Soc.* 129, 5656–5664. doi:10.1021/ja069124n
- Bertelsen, E. B., Chang, L., Gestwicki, J. E., and Zwietering, E. R. P. (2009). Solution Conformation of Wild-type *E. coli* Hsp70 (DnaK) Chaperone Complexed with ADP and Substrate. *Proc. Natl. Acad. Sci.* 106, 8471–8476. doi:10.1073/pnas.0903503106
- Bertini, I., Del Bianco, C., Gelis, I., Katsaros, N., Luchinat, C., Parigi, G., et al. (2004). From the Cover: Experimentally Exploring the Conformational Space Sampled by Domain Reorientation in Calmodulin. *Proc. Natl. Acad. Sci.* 101, 6841–6846. doi:10.1073/pnas.0308641101
- Boehr, D. D., Nussinov, R., and Wright, P. E. (2009). The Role of Dynamic Conformational Ensembles in Biomolecular Recognition. *Nat. Chem. Biol.* 5, 789–796. doi:10.1038/nchembio.232
- Bonomi, M., and Vendruscolo, M. (2019). Determination of Protein Structural Ensembles Using Cryo-Electron Microscopy. *Curr. Opin. Struct. Biol.* 56, 37–45. doi:10.1016/j.sbi.2018.10.006
- Bonomi, M., Heller, G. T., Camilloni, C., and Vendruscolo, M. (2017). Principles of Protein Structural Ensemble Determination. *Curr. Opin. Struct. Biol.* 42, 106–116. doi:10.1016/j.sbi.2016.12.004
- Bonvin, A. M., Boelens, R., and Kaptein, R. (1994). Time- and Ensemble-Averaged Direct NOE Restraints. *J. Biomol. NMR* 4, 143–149. doi:10.1007/BF00178343
- Bottaro, S., and Lindorff-Larsen, K. (2018). Biophysical Experiments and Biomolecular Simulations: A Perfect Match? *Science* 361, 355–360. doi:10.1126/science.aat4010
- Bowers, K. J., Chow, E., Xu, H., Dror, R. O., Eastwood, M. P., Gregersen, B. A., et al. (2006). “Scalable Algorithms for Molecular Dynamics Simulations on Commodity Clusters,” in Proceedings of the ACM/IEEE Conference on Supercomputing (SC06), Tampa, FL, November 11–17, 2006 (Tampa, FL), 43. doi:10.1109/sc.2006.54
- Brooks, B. R., Brooks, C. L., III, MacKerell, A. D., Jr, Nilsson, L., Petrella, R. J., Roux, B., et al. (2009). CHARMM: The Biomolecular Simulation Program. *J. Comput. Chem.* 30, 1545–1614. doi:10.1002/jcc.21287
- Brotzakis, Z. F., Vendruscolo, M., and Bolhuis, P. G. (2021). A Method of Incorporating Rate Constants as Kinetic Constraints in Molecular Dynamics Simulations. *Proc. Natl. Acad. Sci. USA* 118, e2012423118. doi:10.1073/pnas.2012423118
- Camilloni, C., Cavalli, A., and Vendruscolo, M. (2013). Replica-averaged Metadynamics. *J. Chem. Theor. Comput.* 9, 5610–5617. doi:10.1021/ct4006272
- Cavalli, A., Camilloni, C., and Vendruscolo, M. (2013). Molecular Dynamics Simulations with Replica-Averaged Structural Restraints Generate Structural Ensembles According to the Maximum Entropy Principle. *J. Chem. Phys.* 138, 094112. doi:10.1063/1.4793625
- Cerretti, M., Brain, G. A. R., Riordan, O., and Manolopoulos, D. E. (2012). The Inefficiency of Re-weighted Sampling and the Curse of System Size in High-Order Path Integration. *Proc. R. Soc. A* 468, 2–17. doi:10.1098/rspa.2011.0413
- Clare, G. M. (2015). Practical Aspects of Paramagnetic Relaxation Enhancement in Biological Macromolecules. *Meth. Enzymol.* 564, 485–497. doi:10.1016/b.s.mie.2015.06.032
- Davtyan, A., Schafer, N. P., Zheng, W., Clementi, C., Wolynes, P. G., and Papoian, G. A. (2012). AWSEM-MD: Protein Structure Prediction Using Coarse-Grained Physical Potentials and Bioinformatically Based Local Structure Biasing. *J. Phys. Chem. B* 116, 8494–8503. doi:10.1021/jp212541y
- Delaforge, E., Milles, S., Bouvignies, G., Bouvier, D., Boivin, S., Salvi, N., et al. (2015). Large-Scale Conformational Dynamics Control H5N1 Influenza Polymerase PB2 Binding to Importin α . *J. Am. Chem. Soc.* 137, 15122–15134. doi:10.1021/jacs.5b07765
- Ding, Y.-H., Gong, Z., Dong, X., Liu, K., Liu, Z., Liu, C., et al. (2017). Modeling Protein Excited-State Structures from “Over-length” Chemical Cross-Links. *J. Biol. Chem.* 292, 1187–1196. doi:10.1074/jbc.m116.761841
- Fajardo, J. E., Shrestha, R., Gil, N., Belsom, A., Crivelli, S. N., Czaplewski, C., et al. (2019). Assessment of Chemical-crosslink-assisted Protein Structure Modeling in CASP13. *Proteins* 87, 1283–1297. doi:10.1002/prot.25816
- Förster, F., Webb, B., Krukenberg, K. A., Tsuruta, H., Agard, D. A., and Sali, A. (2008). Integration of Small-Angle X-ray Scattering Data into Structural Modeling of Proteins and Their Assemblies. *J. Mol. Biol.* 382, 1089–1106. doi:10.1016/j.jmb.2008.07.074
- Frenkel, D., and Smit, B. (2000). *Understanding Molecular Simulation: From Algorithms to Applications*. New York: Academic Press.
- Gong, Z., Ding, Y.-H., Dong, X., Liu, N., Zhang, E. E., Dong, M.-Q., et al. (2015). Visualizing the Ensemble Structures of Protein Complexes Using Chemical Cross-Linking Coupled with Mass Spectrometry. *Biophys. Rep.* 1, 127–138. doi:10.1007/s41048-015-0015-y
- Gong, Z., Gu, X.-H., Guo, D.-C., Wang, J., and Tang, C. (2017a). Protein Structural Ensembles Visualized by Solvent Paramagnetic Relaxation Enhancement. *Angew. Chem. Int. Ed.* 56, 1002–1006. doi:10.1002/anie.201609830
- Gong, Z., Liu, Z., Dong, X., Ding, Y.-H., Dong, M.-Q., and Tang, C. (2017b). Protocol for Analyzing Protein Ensemble Structures from Chemical Cross-Links Using DynaXL. *Biophys. Rep.* 3, 100–108. doi:10.1007/s41048-017-0044-9
- Gong, Z., Ye, S.-X., and Tang, C. (2020). Tightening the Crosslinking Distance Restraints for Better Resolution of Protein Structure and Dynamics. *Structure* 28, 1160–1167. doi:10.1016/j.str.2020.07.010
- Corba, C., Miyashita, O., and Tama, F. (2008). Normal-mode Flexible Fitting of High-Resolution Structure of Biological Molecules toward One-Dimensional Low-Resolution Data. *Biophys. J.* 94, 1589–1599. doi:10.1529/biophysj.107.122218
- Groth, M., Malicka, J., Czaplewski, C., Oldziej, S., Łankiewicz, L., Wiczak, W., et al. (1999). Maximum Entropy Approach to the Determination of Solution Conformation of Flexible Polypeptides by Global Conformational Analysis and NMR Spectroscopy – Application to DNS1-C-[D-A2bu2,Trp4,Leu5]enkephalin and DNS1-C-[D-A2bu2,Trp4,D-Leu5]enkephalin. *J. Biomol. NMR* 15, 315–330. doi:10.1023/a:1008349424452
- Gu, X.-H., Gong, Z., Guo, D.-C., Zhang, W.-P., and Tang, C. (2014). A Decadentate Gd(III)-coordinating Paramagnetic Cosolvent for Protein Relaxation Enhancement Measurement. *J. Biomol. NMR* 58, 149–154. doi:10.1007/s10858-014-9817-3
- Güntert, P., and Buchner, L. (2015). Combined Automated NOE Assignment and Structure Calculation with CYANA. *J. Biomol. NMR* 62, 453–471. doi:10.1007/s10858-015-9924-9
- Hansmann, U. H. E. (1997). Parallel Tempering Algorithm for Conformational Studies of Biological Molecules. *Chem. Phys. Lett.* 281, 140–150. doi:10.1016/s0009-2614(97)01198-6
- Hartmüller, C., Spreitzer, E., Göbl, C., Falsone, F., and Madl, T. (2019). NMR Characterization of Solvent Accessibility and Transient Structure in Intrinsically Disordered Proteins. *J. Biomol. NMR* 73, 305–317. doi:10.1007/s10858-019-00248-2
- Henzler-Wildman, K., and Kern, D. (2007). Dynamic Personalities of Proteins. *Nature* 450, 964–972. doi:10.1038/nature06522
- Hermann, M. R., and Hub, J. S. (2019). SAXS-restrained Ensemble Simulations of Intrinsically Disordered Proteins with Commitment to the Principle of Maximum Entropy. *J. Chem. Theor. Comput.* 15, 5103–5115. doi:10.1021/acs.jctc.9b00338

- Huang, Y. J., Mao, B., Xu, F., and Montelione, G. T. (2015). Guiding Automated NMR Structure Determination Using a Global Optimization Metric, the NMR DP Score. *J. Biomol. NMR* 62, 439–451. doi:10.1007/s10858-015-9955-2
- Hummer, G., and Köfinger, J. (2015). Bayesian Ensemble Refinement by Replica Simulations and Reweighting. *J. Chem. Phys.* 143, 243150. doi:10.1063/1.4937786
- Joo, K., Joung, I., Lee, J., Lee, J., Lee, W., Brooks, B., et al. (2015). Protein Structure Determination by Conformational Space Annealing Using NMR Geometric Restraints. *Proteins* 83, 2251–2262. doi:10.1002/prot.24941
- Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., et al. (2021). Highly Accurate Protein Structure Prediction with AlphaFold. *Nature* 596, 593–589. doi:10.1038/s41586-021-03819-2
- Karczyńska, A. S., Mozolewska, M. A., Krupa, P., Giełdoń, A., Liwo, A., and Czaplewski, C. (2018). Prediction of Protein Structure with the Coarse-Grained UNRES Force Field Assisted by Small X-ray Scattering Data and Knowledge-Based Information. *Proteins* 86 (S1), 228–239. doi:10.1002/prot.25421
- Kirkpatrick, S., Gelatt, C. D. J., and Vecchi, M. P. (1983). Optimization by Simulated Annealing. *Science* 220, 671–680. doi:10.1126/science.220.4598.671
- Kityk, R., Kopp, J., Sinning, I., and Mayer, M. P. (2012). Structure and Dynamics of the ATP-Bound Open Conformation of Hsp70 Chaperones. *Mol. Cell* 48, 863–874. doi:10.1016/j.molcel.2012.09.023
- Kmiecik, S., Gront, D., Kolinski, M., Wieteska, L., Dawid, A. E., and Kolinski, A. (2016). Coarse-grained Protein Models and Their Applications. *Chem. Rev.* 116, 7898–7936. doi:10.1021/acs.chemrev.6b00163
- Kogut, M., Gong, Z., Tang, C., and Liwo, A. (2021). Pseudopotentials for Coarse-grained Cross-link-assisted Modeling of Protein Structures. *J. Comput. Chem.* 42, 2054–2067. doi:10.1002/jcc.26736
- Kolinski, A. (2004). Protein Modeling and Structure Prediction with a Reduced Representation. *Acta Biochim. Pol.* 51, 349–371. doi:10.18388/abp.2004_3575
- Konarev, P. V., Volkov, V. V., Sokolova, A. V., Koch, M. H. J., and Svergun, D. I. (2003). PRIMUS: a Windows PC-Based System for Small-Angle Scattering Data Analysis. *J. Appl. Cryst.* 36, 1277–1282. doi:10.1107/s0021889803012779
- Konrat, R. (2014). NMR Contributions to Structural Dynamics Studies of Intrinsically Disordered Proteins. *J. Magn. Reson.* 241, 74–85. doi:10.1016/j.jmr.2013.11.011
- Kozak, M., Lewandowska, A., Oldziej, S., Rodziewicz-Motowidło, S., and Liwo, A. (2010). Combination of SAXS and NMR Techniques as a Tool for the Determination of Peptide Structure in Solution. *J. Phys. Chem. Lett.* 1, 3128–3131. doi:10.1021/jz101178t
- Krzeminski, M., Marsh, J. A., Neale, C., Choy, W.-Y., and Forman-Kay, J. D. (2013). Characterization of Disordered Proteins with Ensemble. *Bioinformatics* 29, 398–399. doi:10.1093/bioinformatics/bts701
- Kümmerer, F., Orioli, S., Harding-Larsen, D., Hoffmann, F., Gavrilov, Y., Teilum, K., et al. (2021). Fitting Side-Chain NMR Relaxation Data Using Molecular Simulations. *J. Chem. Theor. Comput.* 17, 5262–5275. doi:10.1021/acs.jctc.0c01338
- Latek, D., and Kolinski, A. (2011). CABS-NMR-De Novo Tool for Rapid Global Fold Determination from Chemical Shifts, Residual Dipolar Couplings and Sparse Methyl-Methyl Noes. *J. Comput. Chem.* 32, 536–544. doi:10.1002/jcc.21640
- Leitner, A., Joachimiak, L. A., Unverdorben, P., Walzthoeni, T., Frydman, J., Förster, F., et al. (2014). Chemical Cross-Linking/mass Spectrometry Targeting Acidic Residues in Proteins and Protein Complexes. *Proc. Natl. Acad. Sci. U.S.A.* 111, 9455–9460. doi:10.1073/pnas.1320298111
- Lerner, E., Barth, A., Hendrix, J., Ambrose, B., Birkedal, V., Blanchard, S. C., et al. (2021). FRET-based Dynamic Structural Biology: Challenges, Perspectives and an Appeal for Open-Science Practices. *eLife* 10, e60416. doi:10.7554/eLife.60416
- Leung, H. T. A., Bignucolo, O., Aregger, R., Dames, S. A., Mazur, A., Bernèche, S., et al. (2016). A Rigorous and Efficient Method to Reweight Very Large Conformational Ensembles Using Average Experimental Data and to Determine Their Relative Information Content. *J. Chem. Theor. Comput.* 12, 383–394. doi:10.1021/acs.jctc.5b00759
- Lindorff-Larsen, K., Best, R. B., DePristo, M. A., Dobson, C. M., and Vendruscolo, M. (2005). Simultaneous Determination of Protein Structure and Dynamics. *Nature* 433, 128–132. doi:10.1038/nature03199
- Liu, Z., Gong, Z., Jiang, W. X., Yang, J., Zhu, W. K., Guo, D. C., et al. (2015). Lys63-linked Ubiquitin Chain Adopts Multiple Conformational States for Specific Target Recognition. *eLife* 4, e05767. doi:10.7554/eLife.05767
- Liu, Z., Gong, Z., Dong, X., and Tang, C. (2016). Transient Protein-Protein Interactions Visualized by Solution NMR. *Biochim. Biophys. Acta Proteins Proteomics* 1864, 115–122. doi:10.1016/j.bbapap.2015.04.009
- Liu, Z., Gong, Z., Cao, Y., Ding, Y.-H., Dong, M.-Q., Lu, Y.-B., et al. (2018). Characterizing Protein Dynamics with Integrative Use of Bulk and Single-Molecule Techniques. *Biochemistry* 57, 305–313. doi:10.1021/acs.biochem.7b00817
- Liwo, A., Sieradzian, A. K., Lipska, A. G., Czaplewski, C., Joung, I., Żmudzka, W., et al. (2019). A General Method for the Derivation of the Functional Forms of the Effective Energy Terms in Coarse-Grained Energy Functions of Polymers. III. Determination of Scale-Consistent Backbone-Local and Correlation Potentials in the UNRES Force Field and Force-Field Calibration and Validation. *J. Chem. Phys.* 150, 155104. doi:10.1063/1.5093015
- Lubecka, E. A., and Liwo, A. (2021). ESCASA : Analytical Estimation of Atomic Coordinates from Coarse-grained Geometry for Nuclear-magnetic-resonance-assisted Protein Structure Modeling. I. Backbone and H β Protons. *J. Comput. Chem.* 42, 1579–1589. doi:10.1002/jcc.26695
- Marrink, S. J., and Tieleman, D. P. (2013). Perspective on the Martini Model. *Chem. Soc. Rev.* 42, 6801–6822. doi:10.1039/c3cs60093a
- Marsh, J. A., and Forman-Kay, J. D. (2012). Ensemble Modeling of Protein Disordered States: Experimental Restraint Contributions and Validation. *Proteins* 80, 556–572. doi:10.1002/prot.23220
- Merkley, E. D., Rysavy, S., Kahraman, A., Hafen, R. P., Daggett, V., and Adkins, J. N. (2014). Distance Restraints from Crosslinking Mass Spectrometry: Mining a Molecular Dynamics Simulation Database to Evaluate Lysine-Lysine Distances. *Protein Sci.* 23, 747–759. doi:10.1002/pro.2458
- Mittag, T., and Forman-Kay, J. D. (2007). Atomic-level Characterization of Disordered Protein Ensembles. *Curr. Opin. Struct. Biol.* 17, 3–14. doi:10.1016/j.sbi.2007.01.009
- Nikiforovich, G. V., Vesterman, B., Betins, J., and Podins, L. (1987). The Space Structure of a Conformationally Labile Oligopeptide in Solution: Angiotensin. *J. Biomol. Struct. Dyn.* 4, 1119–1135. doi:10.1080/07391102.1987.10507702
- Nodet, G., Salmon, L., Ozenne, V., Meier, S., Jensen, M. R., and Blackledge, M. (2009). Quantitative Description of Backbone Conformational Sampling of Unfolded Proteins at Amino Acid Resolution from NMR Residual Dipolar Couplings. *J. Am. Chem. Soc.* 131, 17908–17918. doi:10.1021/ja9069024
- Orioli, S., Larsen, A. H., Bottaro, S., and Lindorff-Larsen, K. (2020). How to Learn from Inconsistencies: Integrating Molecular Simulations with Experimental Data. *Prog. Mol. Biol. Transl. Sci.* 170, 123–176. doi:10.1016/bs.pmbts.2019.12.006
- Otting, G. (2010). Protein NMR Using Paramagnetic Ions. *Annu. Rev. Biophys.* 39, 387–405. doi:10.1146/annurev.biophys.093008.131321
- Pelikan, M., Hura, G., and Hammel, M. (2009). Structure and Flexibility within Proteins as Identified through Small Angle X-ray Scattering. *Gen. Physiol. Biophys.* 28, 174–189. doi:10.4149/gpb_2009_02_174
- Pesce, F., and Lindorff-Larsen, K. (2021). Refining Conformational Ensembles of Flexible Proteins against Small-Angle X-ray Scattering Data. *Biophys. J.* 120, 5124–5135. doi:10.1016/j.bpj.2021.10.003
- Pitera, J. W., and Chodera, J. D. (2012). On the Use of Experimental Observations to Bias Simulated Ensembles. *J. Chem. Theor. Comput.* 8, 3445–3451. doi:10.1021/ct300112v
- Plimpton, S. (1995). Fast Parallel Algorithms for Short-Range Molecular Dynamics. *J. Comput. Phys.* 117, 1–19. doi:10.1006/jcph.1995.1039
- Rhee, Y. M., and Pande, V. S. (2003). Multiplexed-replica Exchange Molecular Dynamics Method for Protein Folding Simulation. *Biophys. J.* 84, 775–786. doi:10.1016/s0006-3495(03)74897-8
- Roux, B., and Weare, J. (2013). On the Statistical Equivalence of Restrained-Ensemble Simulations with the Maximum Entropy Method. *J. Chem. Phys.* 138, 084107. doi:10.1063/1.4792208
- Rózycki, B., Kim, Y. C., and Hummer, G. (2011). SAXS Ensemble Refinement of ESCRT-III CHMP3 Conformational Transitions. *Structure* 19, 109–116. doi:10.1016/j.str.2010.10.006
- Salmon, L., Nodet, G., Ozenne, V., Yin, G., Jensen, M. R., Zweckstetter, M., et al. (2010). NMR Characterization of Long-Range Order in Intrinsically Disordered Proteins. *J. Am. Chem. Soc.* 132, 8407–8418. doi:10.1021/ja101645g
- Salomon-Ferrer, R., Case, D. A., and Walker, R. C. (2013). An Overview of the Amber Biomolecular Simulation Package. *WIREs Comput. Mol. Sci.* 3, 198–210. doi:10.1002/wcms.1121

- Salvi, N., Abyzov, A., and Blackledge, M. (2016). Multi-Timescale Dynamics in Intrinsically Disordered Proteins from NMR Relaxation and Molecular Simulation. *J. Phys. Chem. Lett.* 7, 2483–2489. doi:10.1021/acs.jpcllett.6b00885
- Sasmal, S., Lincoff, J., and Head-Gordon, T. (2017). Effect of a Paramagnetic Spin Label on the Intrinsically Disordered Peptide Ensemble of Amyloid- β . *Biophys. J.* 113, 1002–1011. doi:10.1016/j.bpj.2017.06.067
- Schneidman-Duhovny, D., Hammel, M., and Sali, A. (2011). Macromolecular Docking Restrained by a Small Angle X-ray Scattering Profile. *J. Struct. Biol.* 173, 461–471. doi:10.1016/j.jsb.2010.09.023
- Schwieters, C. D., Bermejo, G. A., and Clore, G. M. (2018). Xplor-NIH for Molecular Structure Determination from NMR and Other Data Sources. *Protein Sci.* 27, 26–40. doi:10.1002/pro.3248
- Sekhar, A., and Kay, L. E. (2019). An NMR View of Protein Dynamics in Health and Disease. *Annu. Rev. Biophys.* 48, 297–319. doi:10.1146/annurev-biophys-052118-115647
- Spreitzer, E., Usluer, S., and Madl, T. (2020). Probing Surfaces in Dynamic Protein Interactions. *J. Mol. Biol.* 432, 2949–2972. doi:10.1016/j.jmb.2020.02.032
- Sterpone, F., Melchionna, S., Tuffery, P., Pasquali, S., Mousseau, N., Cragnolini, T., et al. (2014). The OPEP Protein Model: from Single Molecules, Amyloid Formation, Crowding and Hydrodynamics to DNA/RNA Systems. *Chem. Soc. Rev.* 43, 4871–4893. doi:10.1039/c4cs00048j
- Tang, C., and Gong, Z. (2020). Integrating Non-NMR Distance Restraints to Augment NMR Depiction of Protein Structure and Dynamics. *J. Mol. Biol.* 432, 2913–2929. doi:10.1016/j.jmb.2020.01.023
- Torda, A. E., Scheek, R. M., and van Gunsteren, W. F. (1989). Time-dependent Distance Restraints in Molecular Dynamics Simulations. *Chem. Phys. Lett.* 157, 289–294. doi:10.1016/0009-2614(89)87249-5
- Trewhella, J., Hendrickson, W. A., Kleywegt, G. J., Sali, A., Sato, M., Schwede, T., et al. (2013). Report of the wwPDB Small-Angle Scattering Task Force: Data Requirements for Biomolecular Modeling and the PDB. *Structure* 21, 875–881. doi:10.1016/j.str.2013.04.020
- Tria, G., Mertens, H. D. T., Kachala, M., and Svergun, D. I. (2015). Advanced Ensemble Modelling of Flexible Macromolecules Using X-ray Solution Scattering. *Int. Union Crystallogr. J.* 2, 207–217. doi:10.1107/s205225251500202x
- van der Lee, R., Buljan, M., Lang, B., Weatheritt, R. J., Daughdrill, G. W., Dunker, A. K., et al. (2014). Classification of Intrinsically Disordered Regions and Proteins. *Chem. Rev.* 114, 6589–6631. doi:10.1021/cr400525m
- van Gunsteren, W. F., Allison, J. R., Daura, X., Dolenc, J., Hansen, N., Mark, A. E., et al. (2016). Deriving Structural Information from Experimentally Measured Data on Biomolecules. *Angew. Chem. Int. Ed. Engl.* 55, 15990–16010. doi:10.1002/anie.201601828
- Vögeli, B., Segawa, T. F., Leitz, D., Sobol, A., Choutko, A., Trzesniak, D., et al. (2009). Exact Distances and Internal Dynamics of Perdeuterated Ubiquitin from NOE Buildups. *J. Am. Chem. Soc.* 131, 17215–17225. doi:10.1021/ja905366h
- Vögeli, B., Olsson, S., Güntert, P., and Riek, R. (2016). The Exact NOE as an Alternative in Ensemble Structure Determination. *Biophys. J.* 110, 113–126. doi:10.1016/j.bpj.2015.11.031
- Voth, G. (2008). *Coarse-Graining of Condensed Phase and Biomolecular Systems*. 1st edn. Boca Raton: CRC Press, Taylor & Francis Group.
- White, A. D., Dama, J. F., and Voth, G. A. (2015). Designing Free Energy Surfaces that Match Experimental Data with Metadynamics. *J. Chem. Theor. Comput.* 11, 2451–2460. doi:10.1021/acs.jctc.5b00178

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Czaplewski, Gong, Lubecka, Xue, Tang and Liwo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.