



Crosstalk between Cdk5 and GSK3 β : implications for Alzheimer's disease

Olivia Engmann and Karl Peter Giese*

Institute of Psychiatry, King's College London, London, UK

Edited by:

William Wisden, Imperial College, UK

Reviewed by:

Jochen Herms, Ludwig-Maximilians-Universität München, Germany

A. Martinez, University of California, USA

Karen Duff, Columbia University Center for Neuroscience Initiatives, USA

*Correspondence:

Karl Peter Giese, Centre for the Cellular Basis of Behaviour, Institute of Psychiatry, King's College London, James Black Centre, 125 Coldharbour Lane, London, SE5 9NU, UK.
e-mail: peter.giese@iop.kcl.ac.uk

GSK3 β and Cdk5 are the two kinases in the center of research on Alzheimer's disease (AD), involved in the pathological symptoms of AD, A β plaque formation, tau hyperphosphorylation and neurodegeneration. So far, both kinases have mostly been examined in isolation, leading to a schism of the research field into defenders of the GSK3 β -versus the Cdk5 hypotheses of AD. However, in this debate the fact that activities of GSK3 β and Cdk5 can influence each other deserves more attention. Recent evidence from p25 transgenic mice suggests that there is a dynamic crosstalk: during aging or prolonged overactivation of Cdk5, GSK3 β activity may alter in favor of AD pathogenesis. In this review we summarize the connections between GSK3 β and Cdk5 and discuss implications for AD hypotheses.

Keywords: cyclin-dependent kinase 5, glycogen synthase-kinase 3 β , Alzheimer's disease, tau hyperphosphorylation

INTRODUCTION

Alzheimer's disease (AD) is the most prevalent form of dementia, currently affecting more than 24 million people worldwide (Ferri et al., 2005). The predominant symptoms include memory impairments, later followed by a decline of intellectual skills and loss of control over body functions (Mayeux, 2003). Pathologically, AD is characterized by three hallmarks: (1) Neurofibrillary tangles (NFT), consisting of filamentous aggregates of hyperphosphorylated tau (Lovestone and Reynolds, 1997). (2) Senile plaques, which are mostly derived from β -amyloid or "A β " (Selkoe, 1994); A β ₁₋₄₂ is an aberrant cleavage product from amyloid precursor protein (APP) and can aggregate to complexes with toxic properties (Hardy and Higgins, 1992). (3) Brain atrophy, which is caused by loss of both synapses and whole neurons (Gómez-Isla et al., 1997).

As putative mediators of these symptoms, two proline-directed serine/threonine kinases became the focus of molecular AD research: Glycogen synthase kinase 3 β (GSK3 β) and cyclin-dependent kinase 5 (Cdk5), whose activities appear to be increased in the AD brain (Lee et al., 1999; Pei et al., 1997, 1998; Yamaguchi et al., 1996).

GSK3 β is starring in a variety of physiological processes such as regulation of cellular morphology, neuronal outgrowth and motility and synaptic plasticity (Peineau et al., 2008). GSK3 β integrates a variety of intracellular and extracellular pathways, including wnt signalling, the insulin pathway, G-protein coupled receptors and others – the effect is a change in GSK3 β activity by phosphorylation, protein binding and/or cleavage (Jope and Johnson, 2004).

Cdk5 is named after its structural similarity to members of the serine/threonine cyclin-dependent kinase family. Cdk5 reaches a peak kinase activity in neurons due to restricted expression of its activators p35 and p39 (Hellmich et al., 1992). Cleavage of p35 to p25 by calpain leads to overactivation of Cdk5 (Patrick et al., 1999). Cdk5 is a main player in processes of neural development,

synaptic signalling, learning and memory (Angelo et al., 2006). Under physiological conditions, p25 occurs in only minor quantities. Formation of p25 may play a role in learning processes (Angelo et al., 2003; Fischer et al., 2005). It has been suggested that p25 levels and therefore Cdk5 activity are increased in AD brain (Patrick et al., 1999), but studies by other groups did not confirm this finding or even reported a down regulation of p25 in AD (Takashima et al., 2001; Yoo and Lubec, 2001).

Like GSK3 β , Cdk5 integrates various signalling pathways, induced by NMDA receptor activity, Rac and growth factors (Dhavan and Tsai, 2001; Nikolic, 2002). Several risk factors of AD such as inflammation and oxidative stress have been shown to lead to increased Cdk5 activity (Muylleert et al., 2008; Strocchi et al., 2003).

Both GSK3 β and Cdk5 can regulate synaptic plasticity via regulation of long-term potentiation (LTP), synaptic vesicle release and modification of the NMDA Receptor (Angelo et al., 2006; Giese, 2009).

Besides sharing both up- and downstream pathways, Cdk5 and GSK3 β are linked more directly in neurons. In this review we will summarize the mechanisms of crosstalk between GSK3 β and Cdk5 and discuss its implications for tau hyperphosphorylation, a main pathological feature of AD.

CO-LOCALIZATION AND MUTUAL REGULATION OF GSK3 β AND Cdk5

The first studies addressing crosstalk between Cdk5 and GSK3 β were performed by Morfini et al. (2004) using cultured neurons: Their investigations to determine a regulatory pathway for kinesin-driven motility in axons revealed that Cdk5 is necessary for kinesin-driven motility within the axon but cannot directly phosphorylate kinesin. Inhibition of Cdk5 by olomoucine enhanced phosphorylation of

kinesin, suggesting that the effect was mediated by negative regulation of other kinases. Therefore activities of the other kinases Erk, CK1 and GSK3 β were tested after olomoucine treatment. Altered phosphorylation of CREB phosphopeptide, a substrate of GSK3 β , suggested that GSK3 β may be the mediator of Cdk5 activity in kinesin-mediated axonal transport.

Activity of GSK3 β is regulated by phosphorylation; phosphorylation of Tyr216 can enhance enzyme activity, and conversely phosphorylation of Ser9 is inhibitory. Inhibition of Cdk5 resulted in dephosphorylation of the inhibitory Ser9 site on GSK3 β and an increase of activating phosphorylation at Tyr216. Cdk5 does not phosphorylate GSK3 β directly on these sites and Morfini et al. (2004) investigated whether the altered inhibitory phosphorylation at Ser9 may be mediated by a phosphatase. They found that GSK3 β and phosphatase 1 (PP1) co-immunoprecipitate and that Cdk5 can indirectly inhibit GSK3 β activity via inhibition of PP1 (Figure 1). GSK3 β is activated upon PP1-dependent dephosphorylation and is therefore inhibited by Cdk5. In support of this idea, immunostaining revealed that Cdk5, PP1 and GSK3 β co-localize at growth cones. Additionally, incubation of cortical neurons with the non-specific phosphatase inhibitor okadaic acid increased phosphorylation of GSK3 β -Ser9. Later evidence indicated that PP1 is a substrate of Cdk5 (Li et al., 2007). Additionally, regulation of PP1 by Cdk5 may occur via phosphorylation of PP1 inhibitor 1 (Nguyen et al., 2007). Furthermore, Morfini et al. (2004) showed for the first time that Cdk5 can influence tau phosphorylation

indirectly via regulation of GSK3 β . However, these studies were performed on cell culture and the results had to be confirmed by *in vivo* experiments.

TAU HYPERPHOSPHORYLATION

Tau is a microtubule binding protein, which occurs in the healthy brain and functions in axonal transport, assembly and stabilization of microtubules (Buée et al., 2000). Compared to healthy controls, the AD brain contains four to eight times more tau, which is hyperphosphorylated (Khatoun et al., 1994). More than 40 phosphorylation sites have been detected on tau (Hanger et al., 1998) and several kinases and phosphatases have been suggested as deregulators of tau in AD (Buée et al., 2000; Iqbal and Grundke-Iqbal, 2008) – amongst these, Cdk5 and GSK3 β are thought to be the major tau kinases *in vivo* (Flaherty et al., 2000). The fact that the number of NFTs, but not senile plaques correlates highly with the severity of dementia in AD (Dickson et al., 1988) has stimulated a great amount of research into the misregulation of tau. GSK3 β and Cdk5 were proposed as tau kinases after isolating them in a complex with neurofibrillary tangles from AD brain (Yamaguchi et al., 1996). This association between GSK3 β , Cdk5 and tau has been confirmed in post-mortem tissue (Pei et al., 1998, 1999), as well as in cell and mouse models (Flaherty et al., 2000; Li et al., 2006; Noble et al., 2003; Pei et al., 1997; Yamaguchi et al., 1996). However, it is currently debated which kinase is most strongly associated with tangle formation.




FIGURE 1 | In young mutant mice with low p25 expression, GSK3 β is inhibited by phosphorylation at Ser 9. Cdk5-p25 may influence the balance between p-GSK3 and an activatory dephosphorylation via regulation of phosphatases and the PI3K/Akt pathway.

Conditional GSK3 β transgenic mice demonstrate tau hyperphosphorylation and neurodegeneration (Lucas et al., 2001), indicating that GSK3 β activity is sufficient to hyperphosphorylate tau. Despite the accumulating evidence for GSK3 β as major tau kinase, Cdk5 overactivation may also play a role in tau phosphorylation. Tau is a substrate of Cdk5 and is regulated by phosphorylation under physiological conditions, e.g. during G-protein mediated growth cone collapse (Nakayama et al., 1999). Furthermore, numerous studies have shown that Cdk5 can phosphorylate tau at sites which are hyperphosphorylated in AD (Imahori and Uchida, 1997; Lew and Wang, 1995; Sengupta et al., 1997; Tang and Wang, 1996).

A negative correlation between activities of the tau kinases Cdk5 and GSK3 β , as observed by Morfini et al. (2004), may provide a new perspective on the interplay between tau and its kinases. Studies with p25 transgenic mice, expressing low levels of p25 predominantly in the hippocampus (Angelo et al., 2003) that is affected in the early stages of AD (Braak and Braak, 1991), showed that there is crosstalk between Cdk5 and GSK-3 β *in vivo* (Plattner et al., 2006). This was confirmed with another p25 mouse line (Wen et al., 2008). In young age the p25-induced increase in Cdk5 activity inhibits GSK-3 β activity by enhancing the inhibitory phosphorylation at Ser-9 of GSK-3 β (Plattner et al., 2006; Wen et al., 2008). Importantly, at this age the increased Cdk5 activity does not cause tau hyperphosphorylation. However, in old age tau becomes hyperphosphorylated in these p25 mutants (Plattner et al., 2006; Wen et al., 2008). This tau hyperphosphorylation is a result of an age-dependent loss of the Cdk5 inhibition of GSK-3 β . In fact in the older age GSK-3 β activity is enhanced in the p25 mutants. Thus, increased GSK-3 β activity, and not Cdk5 activity, leads to tau hyperphosphorylation.

Plattner et al. (2006) inspected whether in young age Cdk5 and GSK3 β interact with other signalling molecules *in vivo*. They found that GSK3 β and Cdk5 aggregate in a complex with protein phosphatase 2A (PP2A), but not with PP1 as found by Morfini et al. (2004) *in vitro*. Furthermore, blocking phosphatases with okadaic acid resulted in less GSK3 β activity in young p25 mice, which supports the evidence that the inhibitory Ser9 site is regulated by phosphatase activity. Thus, the inhibitory crosstalk on GSK-3 β activity in young age is linked to altered balance in phosphatase activity. However, it is unclear why the inhibitory crosstalk gets lost with age and why in older age GSK-3 β activity is increased in the p25 mutants.

The correlation between the time courses of increasing GSK3 β activity and tau hyperphosphorylation suggests that GSK3 β is the major tau kinase in these p25 transgenic mice. This idea was confirmed by treatment with lithium, an inhibitor of GSK3 β . Mutant mice treated with lithium do not exhibit hyperphosphorylation of tau epitopes. In fact, other mouse models with high p25 expression with tau hyperphosphorylation and tangle formation possessed elevated GSK3 β activity too (Noble et al., 2003) but for other reasons as the activating Tyr-216 phosphorylation is increased in comparison to a control group, which was not found by Plattner et al. (2006). Moreover, in p35 knockout mice tau is hyperphosphorylated and GSK3 β activity increased while activity of Cdk5 is decreased (Hallows et al., 2003). These mice

may reflect the scenario of young p25 mutants, in which Cdk5 activity and GSK3 β activity correlate inversely. At the same time this study shows that Cdk5 activity is not a prerequisite for tau hyperphosphorylation.

Cdk5 and GSK3 β may be linked by more intricate pathways: It has been shown that phosphorylation of Ser9 after Cdk5 activation can be mediated by ErbB and PI3K/Akt pathway (Wen et al., 2008): Cdk5 can phosphorylate T871 on ErbB2 and S1120 on Erb3 (Li et al., 2003) and Erb phosphorylation and GSK3 phosphorylation correlate in p25 transgenic mice (Figure 1).

The activity of GSK3 β was found up-regulated in AD brain (Giese, 2009), while the role of Cdk5 regulators in AD brain is still controversial. Further studies with post-mortem tissue may elucidate the inverse relationship between the two kinases during disease progression.


CONCLUSIONS

Recent evidences have shown that the activities of Cdk5 and GSK3 β are linked and the crosstalk depends on ageing. The proposed mechanisms for inhibition of GSK-3 β by Cdk5 include phosphatases and ErbB signaling. However, it is unclear why the impact from Cdk5 on GSK3 β activity renders from an inhibitory into an activatory state in ageing p25 transgenic mice. One possibility is that ageing neurons may become more sensitive to p25 exposure, in which case research on mouse models of p25 expression should put more emphasis on studying aged mice. On the other hand it is conceivable that GSK3 β activity changes during steady exposure to p25, be it short periods of high levels of p25 or prolonged exposure to low p25 levels. As Plattner et al. (2006) have demonstrated, Cdk5 may exert its effects on tau phosphorylation mostly via GSK3 β since inhibition of GSK3 β by lithium reversed this effect. Therefore, tau hyperphosphorylation in other p25 models may also result from changes in GSK3 β activity (Figure 2).

Young female mice with constant expression of low p25 or inducible mice with short exposure to high p25 levels (Fischer et al., 2005; Ris et al., 2005) exhibit improved learning and memory (L&M). This observation has led to the hypothesis that p25 expression is not a cause of AD but a reaction to stimuli that impair L&M, ultimately turning from beneficial to detrimental as p25 exposure continues (Angelo et al., 2006). The learning ability during p25 expression in mice from early improvements toward neurodegeneration may be mediated by changing GSK3 β activity. Therefore, studies of these mice at various time points should not only include substrates of Cdk5 but also focus on GSK3 β pathways.

P25 has a half life of less than 30 min in cultured neurons (Patrick et al., 1999). If lifelong exposure to p25 insults leads to changes in GSK3 β activity, information about Cdk5 activity may be stored within the cell, for instance in phosphorylation of substrates with low turnover rate such as tau or within the genome. Identifying how this information is stored and resetting it in ageing p25 mutants may be useful to reverse insults that have already occurred during a life time.

Mutant mice with low p25 expression model ageing effects as the most important risk factors of AD. Understanding why GSK3 β



turns from an inhibited into a hyperactive state during ageing of these mice and testing the mutants with drugs that can revert this process seems a promising approach to prevent or treat tau hyperphosphorylation in AD.

REFERENCES

- Angelo, M., Plattner, F., and Giese, K. P. (2006). Cyclin-dependent kinase 5 in synaptic plasticity, learning and memory. *J. Neurochem.* 99, 353–370.
- Angelo, M., Plattner, F., Irvine, E. E., and Giese, K. P. (2003). Improved reversal learning and altered fear conditioning in transgenic mice with regionally restricted p25 expression. *Eur. J. Neurosci.* 18, 423–431.
- Braak, H., and Braak, E. (1991). Alzheimer's disease affects limbic nuclei of the thalamus. *Acta Neuropathol.* 81, 261–268.
- Buée, L., Bussi re, T., Bu e-Scherrer, V., Delacourte, A., and Hof, P. R. (2000). Tau protein isoforms, phosphorylation and role in neurodegenerative disorders. *Brain Res. Brain Res. Rev.* 33, 95–130.
- Dhavan, R., and Tsai, L. H. (2001). A decade of CDK5. *Nat. Rev. Mol. Cell Biol.* 2, 749–759.
- Dickson, D. W., Farlo, J., Davies, P., Crystal, H., Fuld, P., and Yen, S. H. (1988). Alzheimer's disease. A double-labeling immunohistochemical study of senile plaques. *Am. J. Pathol.* 132, 86–101.
- Ferri, C. P., Prince, M., Brayne, C., Brodaty, H., Fratiglioni, L., Ganguli, M., Hall, K., Hasegawa, K., Hendrie, H., Huang, Y., Jorm, A., Mathers, C., Menezes, P. R., Rimmer, E., Sczufca, M., and Alzheimer's Disease International (2005). Global prevalence of dementia: a Delphi consensus study. *Lancet* 366, 2112–2117.
- Fischer, A., Sananbenesi, F., Pang, P. T., Lu, B., and Tsai, L. H. (2005). Opposing roles of transient and prolonged expression of p25 in synaptic plasticity and hippocampus-dependent memory. *Neuron* 48, 825–838.
- Flaherty, D. B., Soria, J. P., Tomasiewicz, H. G., and Wood, J. G. (2000). Phosphorylation of human tau protein by microtubule-associated kinases: GSK3beta and cdk5 are key participants. *J. Neurosci. Res.* 62, 463–472.
- Giese, K. P. (2009). GSK-3: a key player in neurodegeneration and memory. *IUBMB Life* 61, 516–521.
- G mez-Isla, T., Hollister, R., West, H., Mui, S., Growdon, J. H., Petersen, R. C., Parisi, J. E., and Hyman, B. T. (1997). Neuronal loss correlates with but exceeds neurofibrillary tangles in

ACKNOWLEDGEMENTS

We thank Prof. Chris Miller and Dr. Andrew Thompson for helpful comments on the manuscript. This work has been supported by a Medical Research Council Ph.D. studentship.

- Alzheimer's disease. *Ann. Neurol.* 41, 17–24.
- Hallows, J. L., Chen, K., DePinho, R. A., and Vincent, I. (2003). Decreased cyclin-dependent kinase 5 (cdk5) activity is accompanied by redistribution of cdk5 and cytoskeletal proteins and increased cytoskeletal protein phosphorylation in p35 null mice. *J. Neurosci.* 23, 10633–10644.
- Hanger, D. P., Betts, J. C., Loviny, T. L., Blackstock, W. P., and Anderton, B. H. (1998). New phosphorylation sites identified in hyperphosphorylated tau (paired helical filament-tau) from Alzheimer's disease brain using nano-electrospray mass spectrometry. *J. Neurochem.* 71, 2465–2476.
- Hardy, J. A., and Higgins, G. A. (1992). Alzheimer's disease: the amyloid cascade hypothesis. *Science* 256, 184–185.
- Hellmich, M. R., Pant, H. C., Wada, E., and Battey, J. F. (1992). Neuronal cdc2-like kinase: a cdc2-related protein kinase with predominantly neuronal expression. *Proc. Natl. Acad. Sci. U.S.A.* 89, 10867–10871.
- Imahori, K., and Uchida, T. (1997). Physiology and pathology of tau protein kinases in relation to Alzheimer's disease. *J. Biochem.* 121, 179–188.
- Iqbal, K., and Grundke-Iqbal, I. (2008). Alzheimer neurofibrillary degeneration: significance, etiopathogenesis, therapeutics and prevention. *J. Cell Mol. Med.* 12, 38–55.
- Joep, R. S., and Johnson, G. V. (2004). The glamour and gloom of glycogen synthase kinase-3. *Trends Biochem. Sci.* 29, 95–102.
- Khatoun, S., Grundke-Iqbal, I., and Iqbal, K. (1994). Levels of normal and abnormally phosphorylated tau in different cellular and regional compartments of Alzheimer disease and control brains. *FEBS Lett.* 351, 80–84.
- Lee, K. Y., Clark, A. W., Rosales, J. L., Chapman, K., Fung, T., and Johnston, R. N. (1999). Elevated neuronal Cdc2-like kinase activity in the Alzheimer disease brain. *Neurosci. Res.* 34, 21–29.
- Lew, J., and Wang, J. H. (1995). Neuronal cdc2-like kinase. *Trends Biochem. Sci.* 20, 33–37.
- Li, B. S., Ma, W., Jaffe, H., Zheng, Y., Takahashi, S., Zhang, L., Kulkarni, A. B., and Pant, H. C. (2003). Cyclin-dependent kinase-5 is involved in neuregulin-dependent activation of phosphatidylinositol 3-kinase and Akt activity mediating neuronal survival. *J. Biol. Chem.* 278, 35702–35709.
- Li, T., Chalifour, L. E., and Paudel, H. K. (2007). Phosphorylation of protein phosphatase 1 by cyclin-dependent protein kinase 5 during nerve growth factor-induced PC12 cell differentiation. *J. Biol. Chem.* 282, 6619–6628.
- Li, T., Hawkes, C., Qureshi, H. Y., Kar, S., and Paudel, H. K. (2006). Cyclin-dependent protein kinase 5 primes microtubule-associated protein tau site-specifically for glycogen synthase kinase 3 β . *Biochemistry* 45, 3134–3145.
- Lovestone, S., and Reynolds, C. H. (1997). The phosphorylation of tau: a critical stage in neurodevelopment and neurodegenerative processes. *Neuroscience* 78, 309–324.
- Lucas, J. J., Hernández, F., Gómez-Ramos, P., Morán, M. A., Hen, R., and Avila, J. (2001). Decreased nuclear beta-catenin, tau hyperphosphorylation and neurodegeneration in GSK-3 β conditional transgenic mice. *EMBO J.* 20, 27–39.
- Mayeux, R. (2003). Epidemiology of neurodegeneration. *Annu. Rev. Neurosci.* 26, 81–104.
- Morfini, G., Szebenyi, G., Brown, H., Pant, H. C., Pigino, G., DeBoer, S., Beffert, U., and Brady, S. T. (2004). A novel CDK5-dependent pathway for regulating GSK3 activity and kinesin-driven motility in neurons. *EMBO J.* 23, 2235–2245.
- Muylaert, D., Terwel, D., Kremer, A., Sennvik, K., Borghgraef, P., Devijver, H., Dewachter, I., and Van Leuven, F. (2008). Neurodegeneration and neuroinflammation in cdk5/p25-inducible mice: a model for hippocampal sclerosis and neocortical degeneration. *Am. J. Pathol.* 172, 470–485.
- Nakayama, T., Goshima, Y., Misu, Y., and Kato, T. (1999). Role of cdk5 and tau phosphorylation in heterotrimeric G protein-mediated retinal growth cone collapse. *J. Neurobiol.* 41, 326–339.
- Nguyen, C., Nishi, A., Kansy, J. W., Fernandez, J., Hayashi, K., Gillardon, F., Hemmings, H. C. Jr, Nairn, A. C., and Bibb, J. A. (2007). Regulation of protein phosphatase inhibitor-1 by cyclin-dependent kinase 5. *J. Biol. Chem.* 282, 16511–16520.
- Nikolic, M. (2002). The role of Rho GTPases and associated kinases in regulating neurite outgrowth. *Int. J. Biochem. Cell Biol.* 34, 731–745.
- Noble, W., Olm, V., Takata, K., Casey, E., Mary, O., Meyerson, J., Gaynor, K., LaFrancois, J., Wang, L., Kondo, T., Davies, P., Burns, M., Veeranna, Nixon, R., Dickson, D., Matsuoka, Y., Ahljianian, M., Lau, L. F., and Duff, K. (2003). Cdk5 is a key factor in tau aggregation and tangle formation *in vivo*. *Neuron* 38, 555–565.
- Patrick, G. N., Zukerberg, L., Nikolic, M., de la Monte, S., Dikkes, P., and Tsai, L. H. (1999). Conversion of p35 to p25 deregulates Cdk5 activity and promotes neurodegeneration. *Nature* 402, 615–622.
- Pei, J. J., Braak, E., Braak, H., Grundke-Iqbal, I., Iqbal, K., Winblad, B., and Cowburn, R. F. (1999). Distribution of active glycogen synthase kinase 3 β (GSK-3 β) in brains staged for Alzheimer disease neurofibrillary changes. *J. Neuropathol. Exp. Neurol.* 58, 1010–1019.
- Pei, J. J., Grundke-Iqbal, I., Iqbal, K., Bogdanovic, N., Winblad, B., and Cowburn, R. F. (1998). Accumulation of cyclin-dependent kinase 5 (cdk5) in neurons with early stages of Alzheimer's disease neurofibrillary degeneration. *Brain Res.* 797, 267–277.
- Pei, J. J., Tanaka, T., Tung, Y. C., Braak, E., Iqbal, K., and Grundke-Iqbal, I. (1997). Distribution, levels, and activity of glycogen synthase kinase-3 in the Alzheimer disease brain. *J. Neuropathol. Exp. Neurol.* 56, 70–78.
- Peineau, S., Bradley, C., Taghibiglou, C., Doherty, A., Bortolotto, Z. A., Wang, Y. T., and Collingridge, G. L. (2008). The role of GSK-3 in synaptic plasticity. *Br. J. Pharmacol.* 153(Suppl 1), S428–S437.
- Plattner, F., Angelo, M., and Giese, K. P. (2006). The roles of cyclin-dependent kinase 5 and glycogen synthase kinase 3 in tau hyperphosphorylation. *J. Biol. Chem.* 281, 25457–25465.
- Ris, L., Angelo, M., Plattner, F., Capron, B., Errington, M. L., Bliss, T. V., Goudaux, E., and Giese, K. P. (2005). Sexual dimorphisms in the effect of low-level p25 expression on synaptic plasticity and memory. *Eur. J. Neurosci.* 21, 3023–3033.
- Selkoe, D. J. (1994). Alzheimer's disease: a central role for amyloid. *J. Neuropathol. Exp. Neurol.* 53, 438–447.
- Sengupta, A., Wu, Q., Grundke-Iqbal, I., Iqbal, K., and Singh, T. J. (1997). Potentiation of GSK-3-catalyzed Alzheimer-like phosphorylation of human tau by cdk5. *Mol. Cell Biochem.* 167, 99–105.
- Strocchi, P., Pession, A., and Dozza, B. (2003). Up-regulation of cDK5/p35 by oxidative stress in human neuroblastoma IMR-32 cells. *J. Cell Biochem.* 88, 758–765.
- Takahashi, A., Murayama, M., Yasutake, K., Takahashi, H., Yokoyama, M., and Ishiguro, K. (2001). Involvement of cyclin dependent kinase5 activator p25 on tau phosphorylation in mouse brain. *Neurosci. Lett.* 306, 37–40.
- Tang, D., and Wang, J. H. (1996). Cyclin-dependent kinase 5 (Cdk5) and neuron-specific Cdk5 activators. *Prog. Cell Cycle Res.* 2, 205–216.
- Wen, Y., Planel, E., Herman, M., Figueroa, H. Y., Wang, L., Liu, L., Lau, L. F., Yu, W. H., and Duff, K. E. (2008). Interplay between cyclin-dependent kinase 5 and glycogen synthase kinase 3 β mediated by neuregulin signaling leads to differential effects on tau phosphorylation and amyloid precursor protein processing. *J. Neurosci.* 28, 2624–2632.
- Yamaguchi, H., Ishiguro, K., Uchida, T., Takashima, A., Lemere, C. A., and Imahori, K. (1996). Preferential labeling of Alzheimer neurofibrillary tangles with antisera for tau protein kinase (TPK) I/glycogen synthase kinase-3 β and cyclin-dependent kinase 5, a component of TPK II. *Acta Neuropathol.* 92, 232–241.
- Yoo, B. C., and Lubec, G. (2001). p25 Protein in neurodegeneration. *Nature* 411, 763–764; Discussion 764–765.

Conflict of Interest Statement: We declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 06 April 2009; paper pending published: 11 May 2009; accepted: 11 May 2009; published online: 29 May 2009.
 Citation: Engmann O and Giese KP (2009) Crosstalk between Cdk5 and GSK3 β implications for Alzheimer's disease. *Front. Mol. Neurosci.* (2009) 2:2. doi:10.3389/neuro.02.002.2009
 Copyright © 2009 Engmann and Giese. This is an open-access article subject to an exclusive license agreement between the authors and the Frontiers Research Foundation, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.