



Foxb1 Regulates Negatively the Proliferation of Oligodendrocyte Progenitors

Yuanfeng Zhang^{1,2}, Elti Hoxha¹, Tianyu Zhao³, Xunlei Zhou¹
and Gonzalo Alvarez-Bolado^{1*}

¹Department of Neuroanatomy, University of Heidelberg, Heidelberg, Germany, ²Department of Urology, The 2nd Affiliated Hospital of Chongqing Medical University, Chongqing, China, ³Key Laboratory of Oral Disease and Biomedical Sciences, Stomatological Hospital, Chongqing Medical University, Chongqing, China

Oligodendrocyte precursor cells (OPC), neurons and astrocytes share a neural progenitor cell (NPC) in the early ventricular zone (VZ) of the embryonic neuroepithelium. Both switch to produce either of the three cell types and the generation of the right number of them undergo complex genetic regulation. The components of these regulatory cascades vary between brain regions giving rise to the unique morphological and functional heterogeneity of this organ. *Forkhead b1* (*Foxb1*) is a transcription factor gene expressed by NPCs in specific regions of the embryonic neuroepithelium. We used the mutant mouse line *Foxb1-Cre* to analyze the cell types derived from *Foxb1*-expressing NPCs (the *Foxb1* cell lineage) from two restricted regions, the medulla oblongata (MO; hindbrain) and the thalamus (forebrain), of normal and *Foxb1*-deficient mice. *Foxb1* cell lineage derivatives appear as clusters in restricted regions, including the MO (hindbrain) and the thalamus (forebrain). *Foxb1*-expressing NPCs produce mostly oligodendrocytes (OL), some neurons and few astrocytes. *Foxb1*-deficient NPCs generate mostly OPC and immature OL to the detriment of neurons, astrocytes and mature OL. The axonal G-ratio however is not changed. We reveal *Foxb1* as a novel modulator of neuronal and OL generation in certain restricted CNS regions. *Foxb1* biases NPCs towards neuronal generation and inhibits OPC proliferation while promoting their differentiation.

Keywords: Claudin11, GalC, lineage analysis, medulla oblongata, neuroepithelium, NG2, thalamus

OPEN ACCESS

Edited by:

Yun-Qing Li,
Fourth Military Medical University,
China

Reviewed by:

Andrea Wizenmann,
University of Tübingen, Germany
Herbert Hildebrandt,
Hannover Medical School, Germany

*Correspondence:

Gonzalo Alvarez-Bolado
alvarez@ana.uni-heidelberg.de

Received: 17 May 2017

Accepted: 19 June 2017

Published: 05 July 2017

Citation:

Zhang Y, Hoxha E, Zhao T, Zhou X
and Alvarez-Bolado G (2017) *Foxb1*
Regulates Negatively the Proliferation
of Oligodendrocyte Progenitors.
Front. Neuroanat. 11:53.
doi: 10.3389/fnana.2017.00053

INTRODUCTION

Oligodendrocytes (OL) produce and maintain the myelin sheaths around axons of the central nervous system (CNS). In addition, OL can also provide trophic support to axons and promote their viability (Emery, 2010; Mitew et al., 2014). Defects in myelination, the process by which OL wrap axons in layers of myelin, cause severe pathological conditions in humans. In addition, pathological conditions affecting the maturation of the brain white matter in premature infants are on the rise, and their most common result is mental retardation. Since myelination occurs mostly during CNS development, much research has focused on the development of OL. In particular, the mechanisms regulating the proliferation and differentiation of OL, the cells responsible for the formation of the white matter, are currently under intense scrutiny (Goldman and Kuypers, 2015; Marinelli et al., 2016).

OL generate from bipolar, migratory oligodendrocyte progenitor cells (OPC), probably heterogeneous in morphology and physiology, that arise from specific zones of the neuroepithelium ventricular zone (VZ). OPC are identified by the expression of a series of specific antigens, including NG2 chondroitin sulfate proteoglycan (NG2), platelet-derived growth factor- α receptor (PDGFR α) and oligodendrocyte transcription factor 2 (Olig2). The OPC specifically express a number of transcription factors as they migrate to colonize the CNS and produce OL. Each phase in the development of OL (OPC, premyelinating OL and myelinating OL) is associated with intense expression of specific sets of transcription factors; once at the final destination, OL mature and acquire the expression of specific markers: MBP, PLP, MAG and Claudin11 among others and acquire their typical morphology as well as the ability to myelinate (Noll and Miller, 1993; Rowitch and Kriegstein, 2010; de Castro et al., 2013; Goldman and Kuypers, 2015). A key point in this pathway, the decision by the OPC to stop proliferating and start producing OL, is under strong control (Dugas et al., 2010).

There are different dorsal and ventral VZ specific areas generating OPC in the hindbrain (Davies and Miller, 2001; Vallstedt et al., 2005). Here we report expression of transcription factor gene *Forkhead b1* (*Foxb1*) in one of the ventral OPC-generating regions in the hindbrain. *Foxb1* belongs to a gene family encoding hundreds of transcription factors whose DNA binding domain has a winged helix configuration (Weigel and Jäckle, 1990; Carlsson and Mahlapuu, 2002). *Foxb1* is expressed widespread in the early developing VZ of the neural tube and is later restricted to areas of the spinal cord, hindbrain, thalamus and hypothalamus (Ang et al., 1993; Kaestner et al., 1993, 1996; Alvarez-Bolado et al., 1999; Zhao et al., 2007). Involvement of *Foxb1* in OL development has not been reported.

Here we use a *Foxb1-Cre* knockin-knockout mouse line together with reporter mouse lines Z/AP and ROSA26R in order to approach the following questions: (1) Which specific cell types of the CNS are generated by *Foxb1*-expressing VZ (i.e., what is the *Foxb1* cell lineage in the brain)? (2) What is the role of *Foxb1* in oligodendrocyte development? Our results show that *Foxb1* is a novel player in OL development, whose role involves inhibiting OPC proliferation and promoting oligodendrocyte maturation.

MATERIALS AND METHODS

Mouse Handling

All mouse lines were housed and fed according to the German Animal Welfare Act (Tierschutzgesetz) and the European Communities Council Directive in the Interfaculty Biomedical Facility (IBF), University of Heidelberg. The authorization for collecting mice brain and handling animal samples was approved by the Regierungspraesidium Karlsruhe, Baden-Wuerttemberg. The experiments were carried out in the Neuroanatomy Department, University of Heidelberg and all the procedures were performed according to the Animal Welfare Act. Adult mice were killed by cervical dislocation. The staining protocol for some antibodies required mice to be anesthetized with

isofluorane and then intracardially perfused with fixative. For embryos, pregnant mice were sacrificed by cervical dislocation and E12.5, E15.5 or E18.5 pups were taken out of the uterus and processed for histology.

Mutant Mouse Lines

Foxb1-Cre Mouse Line

In this mouse line, transcriptional activation of the *Foxb1* locus initiates expression of *Cre* recombinase and enhanced green fluorescent protein (*EGFP*); as well as *Foxb1* itself in heterozygotes; Zhao et al., 2007, 2008).

Z/AP Reporter Mouse Line

In *Foxb1^{Cre} x Z/AP* mice, *Foxb1*-expressing cells and their progeny permanently express human placental alkaline phosphatase (hPLAP; Lobe et al., 1999). hPLAP is a GPI-linked cell surface marker. In this reporter line, cells not expressing *Cre* recombinase are labeled by expression of β -galactosidase (negative control).

ROSA26R Mouse Reporter Line

When dealing with antibodies detecting proteins enriched in the cell nucleus (for instance, transcription factors, like Olig2), the use of a lineage reporter also expressed in the cell nucleus makes it easier to ascertain cellular colocalization of both marker and reporter. ROSA26R (Soriano, 1999) is such a reporter mouse line, carrying β -galactosidase as reporter gene; upon *Cre* recombination, antibody detection of β -galactosidase produces a characteristic punctate pattern in the cell nucleus (Soriano, 1999). In *Foxb1^{Cre} x ROSA26R* mice the presence of β -galactosidase in the nucleus of *Cre*-expressing cells can be detected from E8.5 on (Zhao et al., 2007, 2008).

Labeling Alkaline Phosphatase Activity

Phosphatase/NBT staining has been described (Lobe et al., 1999; Gierut et al., 2014). Briefly, brains or embryos were dissected in PBS buffer on ice; fixed in 4% paraformaldehyde solution with 0.02% NP-40 and 0.01% sodium deoxycholate at 4°C for 30 min; washed in PBS three times for 30 min at 4°C; incubated in PBS at 72°C for 30 min to inactivate endogenous alkaline phosphatase; rinsed three times in PBS for 10 min at room temperature; washed in alkaline phosphatase buffer two times 10 min; stained with 100 mg/ml NBT and 50 mg/ml BCIP in AP buffer at 4°C until optimal results appeared; washed in PBS extensively to reduce the background; mounted in Mowiol.

Immunohistochemistry

To detect specific cell markers as well as reporter proteins hPLAP and β -galactosidase, the following antibodies and conditions were used: anti-hPLAP (Sigma A2951) 1:2000 on paraffin sections (PS) or fixed frozen tissue (FFT); anti- β -galactosidase (Abcam ab9361) 1:200 (FFT); anti-Claudin11 (Abcam ab53041) 1:500 (FFT); anti-GalC (Chemicon AB142) 1:100 (PS); anti-myelin basic protein MBP (Sigma M3821) 1:100 (FFT); anti-platelet derived growth factor receptor alpha (PDGFR α ; BD Pharmingen 558774) 1:300 on fresh frozen tissue; anti-NG2 (Chemicon AB5320) 1:100 (PS); anti-Olig2 (Chemicon

AB9610) 1:200 (FFT); anti-GFAP (Chemicon AB5804) 1:300 (FFT); anti- β -tubulin III (Abcam ab18207) 1:600 (FFT); anti-NeuN (Abcam ab177487) 1:500 (FFT).

Labeling of Thalamic Neurons through Transfection with Two-Reporter Construct

We cloned a DNA construct carrying a constitutive GFP reporter and a Cre-recombination-dependent tdTomato reporter. Every cell transfected will express the green GFP reporter, but transfected cells expressing Cre will express the red tdTomato reporter instead. Our construct was based on MSCV FLIPi P2G_Thy1.1 Dbl (p53, PTEN), a gift from Richard Hynes (Addgene plasmid #19746; Stern et al., 2008). We introduced the following modifications for our purposes: (1) removal of the cassette Puromycin 2a-GFP-Thy1.1-miR-WRPE; (2) insertion of the reporter tdTomato in reverse orientation in the place of Thy1.1-miR; (3) addition of the CAG promoter. The resulting DNA construct was transfected into the thalamus of E12.5 wild type mice through *in utero* electroporation. The electroporation technique has been described before (Tabata and Nakajima, 2001; Haddad-Tóvolli et al., 2013).

Proliferation Assay

For proliferation studies, 50 mg/kg of body weight of Bromodeoxy-uridine (BrdU) was intraperitoneally injected as aqueous solution into P10 mice three times, at 4 h intervals (Bu et al., 2004). Mice were sacrificed 2 h after the last injection. The brains were dissected and fixed in 4% PFA overnight at 4°C, washed in PBS for 4–6 h, cryoprotected in 0.1 M PBS containing 30% sucrose, embedded in OCT compound on Dry Ice and stored at –80°C. Previous to immunodetection of BrdU and Olig2, cryosections (20 μ m) were incubated in 1 M HCl for 10 min on ice and then in 2 M HCl for 30 min at 37°C and neutralized with 0.1 M sodium borate buffer at room temperature.

Confocal Microscope

The results were observed and analyzed under a confocal laser-scanning microscope (Zeiss LSM 510) with ZEN 2010 software.

Electron Microscopy and G-Ratio

The G-ratio was calculated on measurements performed on electron microscopy photographs of transverse sections of medulla oblongata (MO). The tissue samples were obtained from *Foxb1*^{Cre/+} and *Foxb1*^{Cre/Cre} mouse brains (age P56, three mice per genotype), then processed according to current electron microscopy protocols. The diameter of 80 randomly chosen axons and the corresponding fibers (axon + myelin) were measured for each individual mouse. The data were analyzed with the Mann-Whitney test.

Statistical Analysis

Cell countings were performed on 20 μ m sections from the brains of three animals per genotype. Sagittal sections were cut in four series (A, B, C and D) from an area spanning the middle (rostr-caudally) of the hindbrain. Eight sections were counted per mouse. The counting bin for marker-labeled cells was 600 μ m \times 600 μ m in MO. The counting bin for Olig2 plus

BrdU was 0.1 mm² in MO. The localization of the bins was chosen randomly. Statistical assessment was performed with Prism 5 software (GraphPad Software, San Diego, CA, USA). The Mann-Whitney test was used and the results are represented in bar graphs with mean \pm SD; **p* < 0.05, ***p* < 0.01, ****p* < 0.005.

RESULTS



Foxb1 Is Expressed in Oligodendrocyte-Generating Regions of the Embryonic Hindbrain

Foxb1 is expressed in two narrow longitudinal bands in the VZ of the brainstem (Alvarez-Bolado et al., 1999) reminiscent of the domains of expression of OPC marker genes *Olig1* and *Olig2* (Zhou et al., 2000). We asked if the lineage of the *Foxb1*-expressing neuroepithelium included OL and if *Foxb1* was involved in oligodendrocyte development. We approached this question by using a lineage-labeling mouse line, the *Foxb1-Cre-EGFP x Z/AP*. *Foxb1-Cre-EGFP* is a knockin-knockout of the Cre recombinase and the expression reporter EGFP into the *Foxb1* locus (Zhao et al., 2007). By crossing these lines with the *Z/AP* reporter line (Lobe et al., 1999), *Foxb1*-expressing neuroepithelial cells as well as their entire progeny (even if they do not express *Foxb1*) will permanently express hPLAP in this way becoming identifiable. First, we examined the brains of mouse embryos heterozygous for this insertion. *Foxb1*-heterozygotes have normal phenotype (Dou et al., 1997; Alvarez-Bolado et al., 2000; Kloetzli et al., 2001) and were used here as proxies for the wild type. Treating early embryonic brains with a procedure to detect alkaline phosphatase activity, we detected as expected the bands in the hindbrain (Figures 1A–C). Later in development, we detected AP⁺ cells in the MO (Figure 1D).

By staining MO sections of these mice for hPLAP, we labeled large groups of cells with multiple processes in brains of every developmental age (Figure 2). As development proceeds, *Foxb1*-lineage cells form clusters in the brainstem as well as axon-like structures (Figures 2A–C). An internal control is built in the reporter mouse *Z/AP* (Lobe et al., 1999), so that cells not undergoing a Cre recombination (i.e., not expressing the *Cre* driver, in this case *Foxb1*) express β -galactosidase as a reporter instead. We labeled expression of both markers on brainstem sections to confirm that the hPLAP-expressing clusters did not overlap with β -galactosidase-expressing regions (Figures 2D–F).


The Foxb1 Lineage Includes Neurons as well as Large Numbers of Oligodendrocytes

In order to identify the cellular components of these clusters, we colocalized hPLAP with neuron-specific markers (either β -tubulin-III or NeuN) on *Foxb1*^{Cre/+} *x Z/AP* brains and found a number of double-labeled cells (Figures 3A–F). We confirmed this on primary culture of *Foxb1*^{Cre/+} *x Z/AP* brain cells (Figures 3G–I). Colocalizing astrocyte-specific protein glial fibrillary acidic protein (GFAP) with hPLAP also showed a small



number of *Foxb1*-lineage astrocytes (not shown). As expected, no hPLAP-labeled cells expressed microglial marker Iba-1, since microglia originates outside of the neural tube and therefore far from the *Foxb1*-expressing neuroepithelium (not shown).

The finding that only small numbers of hPLAP-labeled brain cells were neurons or astrocytes suggested that the



most abundant cellular component of the *Foxb1* lineage could be OL. Expression of myelin basic protein (MBP) colocalized partially with hPLAP (**Figures 4A–C**), in principle confirming this impression. The appearance of the tissue was however complex and did not provide appropriate cellular resolution (**Figures 4A–C**). Expression of Claudin11, a marker of mature OL (Bronstein et al., 2000), colocalized with profiles morphologically similar to myelinated axons (**Figures 4D–F**). On primary cultures of cells from the MO of *Foxb1*^{Cre/+} *x* *Z/AP* mice, Claudin11 and hPLAP colocalized on cells with unequivocal mature oligodendrocyte morphology, including the characteristic lamellae (**Figures 4G–I**). Expression of Galactocerebroside C (GalC), a marker of mature as well as immature OL, also colocalized with hPLAP on sections (**Figures 4J–L**) and on primary cultures (**Figures 4M–O**).

Abundant OPC Belong to the *Foxb1*-Lineage

If most of the observed *Foxb1*-lineage brain cells are OL, it follows that some OPC must also belong to this lineage. To confirm this, we colocalized *Foxb1*-lineage reporters with OPC markers Olig2 and NG2 (**Figure 5**). Since Olig2 protein localizes to the cell nucleus, in order to colocalize it with a reporter we crossed our *Foxb1*^{Cre/+} *x* *Z/AP* mice with the




FIGURE 4 | Abundant mature and immature oligodendrocytes (OL) in the *Foxb1* lineage. Antibody detection of hPLAP, myelin basic protein (MBP), Claudin11 and Galactocerebroside C (GalC) (as indicated) on MO sections or primary cell culture (as indicated) of P56 *Foxb1*^{Cre/+} x *Z/AP* mouse.

(A–C) Colocalization of MBP and hPLAP. (D–F) Claudin 11 colocalizes with hPLAP-expressing axon bundles (arrowheads). (G–I) Claudin 11 colocalizes with hPLAP-expressing cultured brain cells with the morphology of mature OL (arrowheads indicate their typical lamella extensions). (J–L) Colocalization of GalC and hPLAP. Arrows indicate the boundary of the GalC-labeled cytoplasm; arrowheads indicate the corresponding cell nucleus. (M–O) GalC colocalizes with hPLAP-expressing cultured brain cells with the morphology of immature OL. Scale bars 50 μ m.

reporter mouse line ROSA26R (Soriano, 1999), whose cells express β -galactosidase as reporter upon Cre recombination. We found numerous cells expressing both this reporter and Olig2 (Figures 5A–C) in the MO of *Foxb1*^{Cre/+} x *ROSA26R* mice. The β subunit of the receptor for PDGFR β as well as NG2 specifically colocalize in OPC (Richardson et al., 1988; Hart et al., 1989). We found also many cells where NG2 colocalized with hPLAP in histological sections of the MO (Figures 5G–I) as well as in culture (Figures 5J–L; see cell abundance quantitation below, Figure 8).




FIGURE 5 | The *Foxb1* lineage includes oligodendrocyte precursor cells (OPC). Antibody detection of hPLAP, Olig2, NG2 and platelet-derived growth factor-alpha receptor (PDGFR β ; as indicated) on MO sections of P56 *Foxb1*^{Cre/+} x *ROSA26R* mouse (A–C) or P56 *Foxb1*^{Cre/+} x *Z/AP* mouse (A–I) or on primary culture of P56 MO (J–L). (A–C) The *Foxb1*-lineage marker β -galactosidase colocalizes with a subpopulation of Olig2-expressing cells (arrowheads). Large and small arrows indicate cells expressing only Olig2 or only β -galactosidase, respectively. (D–F) Colocalization of OPC markers PDGFR β and NG2 (arrowheads). Large and small arrows indicate cells expressing only NG2 or only PDGFR β , respectively. (G–I) In hPLAP-labeled cell clusters (red), some cells express NG2 (arrowheads). (J–L) hPLAP colocalizes with NG2 on some brain cells (arrowheads) in primary culture. Arrows indicate non-colocalizing cells. Scale bars 50 μ m.

The Thalamus Shows Similar *Foxb1*-Lineage Cell Clusters

OPC originate from certain spatially restricted regions of the neural tube. They then migrate and proliferate to supply the entire CNS with OL. A current question concerns the possibility that OL from different origins have differential properties and if OL from any given region can be used to substitute for OL in any other region of the brain (replacement therapy). In this context, we asked if the basic composition of the *Foxb1* lineage in the hindbrain was similar to that of another regionally restricted subpopulation of *Foxb1*-lineage brain cells situated not in the hindbrain but in the forebrain. The thalamus is a forebrain region with strong *Foxb1* developmental expression and a large part of the thalamus is of *Foxb1* lineage (Zhao et al., 2007, 2008).






FIGURE 6 | *Foxb1*-lineage clusters are also abundant in the thalamus. **(A)** Sagittal view of a P56 *Foxb1*^{Cre/+} *x Z/AP* mouse brain (rostral to the left) showing hPLAP-expressing clusters in thalamus (Th), mammillary body (MBO), superior colliculus (SC), pons (P) and MO. Inset: line profile of the section in **(A)**. **(B)** Transverse section of P56 *Foxb1*^{Cre/+} after electroploration (at E12.5) and culture (see “Materials and Methods” Section for details). *Foxb1*-lineage thalamic neurons and cortico-thalamic axons (arrow) are labeled red. Transfected cells not expressing *Foxb1* are green. **(C–K)** Primary culture of *Foxb1*^{Cre/+} *x Z/AP* thalamus showing colocalization of hPLAP with GFAP **(C–E)**, GalC **(F–H)** and β -tubulin-III **(I–K)**. Scale bars 50 μ m.

Clusters expressing hPLAP were obvious in the thalamus (**Figure 6A**). A characteristic property of the thalamus with important functional consequences is the presence of many neurons sending axonal projections to the cortex. We used *in utero* electroporation with a plasmid expressing upon Cre recombination the red reporter tdTomato (see “Materials and Methods” Section for details), which fills the cell body and axons. The results showed abundant *Foxb1*-lineage thalamo-cortical neurons (**Figure 6B**). We confirmed the presence of OL, neurons and astrocytes colocalizing hPLAP on primary thalamic culture (**Figures 6C–K**; see cell abundance quantitation below, **Figure 9**).


The *Foxb1*-Lineage in *Foxb1* Mutant Mice Produces Abnormally Large Numbers of OPC

Comparison of *Foxb1*^{Cre/+} *x* Z/AP and *Foxb1*^{Cre/Cre} *x* Z/AP mouse brains after whole mount histochemical demonstration of AP activity showed very increased staining in the homozygotes (**Figures 7A–D**). This suggested that *Foxb1* had an influence on the cell abundance and maybe also identity. To investigate this, we performed cell countings on tissue of heterozygotes and homozygotes labeled with specific cell markers as well as hPLAP. Our results show, first, that in homozygotes there are more *Foxb1*-lineage cells (**Figure 8A**). We also found that, in homozygotes, *Foxb1*-lineage neurons, astrocytes and mature OL were reduced (**Figures 8B–E**). However, OPC were increased (**Figures 8F,G**). Interestingly, a first approach to the abundance of different *Foxb1*-lineage cell types in the thalamus yielded similar results (**Figures 9A–D**). This suggests that all *Foxb1*-lineage OL are functionally homogeneous.

Although the exact proportion of *Foxb1*-lineage cells to other brain cells is not central to our study, we thought it was of some interest to obtain at least a rough approximation to this value for neurons, OL and OPC. Therefore we counted non-hPLAP



(i.e., non-*Foxb1*-lineage) cell in the bins used to quantitate the hPLAP-expressing (*Foxb1*-lineage) cells. The results are as follows: (1) mature plus immature OL (i.e., GalC-expressing cells) belonging to the *Foxb1*-lineage are 73.17% of the total GalC-positive cells per bin in heterozygotes, and 90.00% in homozygotes; (2) OPC (i.e., NG2-expressing cells) belonging to the *Foxb1*-lineage are 45.40% of all NG2-positive cells per bin in heterozygotes and 75.89% in homozygotes; and (3) neurons (i.e., cells expressing beta-tubulin III) belonging to the *Foxb1*-lineage are 30% of all beta-tubulin III-expressing cells per bin in heterozygotes and 24.78% in homozygotes. As could be expected, OL and OPC of the *Foxb1* lineage “crowd out” the non-*Foxb1* lineage ones in homozygotes, while the proportion of *Foxb1*-



lineage neurons becomes slightly smaller. These data however cannot be directly compared to our cell countings summarized in **Figure 10**, since those are represented in percent of hPLAP-expressing cells (as opposed to percent of all cells in the bin).

We concluded that, in the *Foxb1*-deficient MO, the OPC increase to the expense of neurons and mature OL (**Figure 10**).


OPC Proliferation Increased in *Foxb1* Homozygotes

In order to confirm this increase in OPC, we wanted to use another OPC marker, Olig2. Since Olig2 is localized in the nucleus, we wanted a reporter of the *Foxb1* lineage also localized in the nucleus. For this reason we used a different reporter, the ROSA26R mouse line (Soriano, 1999). In this line, Cre recombination causes expression of β -galactosidase which can be seen in the nucleus. We colocalized Olig2 and β -galactosidase on MO sections of heterozygotes and homozygotes (**Figures 11A–F**). Cell




counting confirmed a significant increase of Olig2+ cells in the homozygotes (**Figure 11G**). Additionally, we labeled dividing cells with BrdU (see “Materials and Methods” Section for details) and then labeled the tissues with BrdU and Olig2 (**Figure 12**). We found that, in homozygous mutants, more Olig2-expressing cells were dividing than in heterozygotes.

The G-Ratio Is Not Altered in the *Foxb1*-Deficient Hindbrain

Finally, we asked if there would be any obvious defect in myelination in the hindbrain of our mutants. The phenotype of *Foxb1*-deficient mice includes several behavioral and CNS developmental defects (Dou et al., 1997; Labosky et al., 1997; Alvarez-Bolado et al., 2000; Radyushkin et al., 2005; see “Discussion” Section). However, alterations suggesting myelination defects have not been described in these mice. This suggests that the OL in this mutant are able to myelinate correctly. To test this hypothesis, we measured axon and fiber



diameter on electron microscopy images from the hindbrain of homo- and heterozygotes and found that the G-ratio



(Friede and Bischhausen, 1982) does not change in the mutants (Figure 13).

DISCUSSION

Lineage analysis of a mouse mutant line reveals transcription factor *Foxb1* as a novel player in oligodendrocyte development, promoting neuron generation and OPC differentiation over OPC generation and proliferation.

This novel function of *Foxb1* is in line with the reported functions of the Fox superfamily of transcription factors, which regulate genetic cascades controlling proliferation, differentiation, metabolism, aging, survival and apoptosis (Carlsson and Mahlapuu, 2002). A case in point is the Foxo subfamily, consisting of negative regulators of genes crucial for cell cycle progression (Lam et al., 2013). *Foxb1* is reportedly essential for the survival of a specific subpopulation of spinal motoneurons (Dou et al., 1997) and of a subset of hypothalamic cells (Alvarez-Bolado et al., 2000). However, a relation of *Foxb1* with OL or the control of cell proliferation has not been described before.

In the *Foxb1* null mutant mouse, the OL lineage increases to the detriment of the neuronal lineage. All neurons and macroglia share an early progenitor in the VZ of the embryonic neuroepithelium (Battiste et al., 2007; Pinto and Götz, 2007; Kriegstein and Alvarez-Buylla, 2009). Since part of the neuroepithelium belongs to the *Foxb1* lineage (i.e., expresses *Foxb1* or derives from *Foxb1*-expressing neural progenitor cells, NPCs), we have to assume that *Foxb1*-deficient NPCs are biased towards OPC generation. Therefore, a general function (direct or indirect) of *Foxb1* would be to modulate a complex genetic cascade which has been given the name of neurogenesis/gliogenesis switch (Martynoga et al., 2012). In the cortex, the most extensively researched CNS region, neurogenesis requires Fox transcription factor genes like *Foxc1* and *Foxg1*; reviewed in Martynoga et al. (2012). *Foxb1*, not expressed in the cortical VZ, could fulfill a similar role in other neuroepithelial regions.

Another consequence of *Foxb1* deficiency is the predominance of OPC and immature OL over mature OL, indicating an additional function of *Foxb1* inhibiting OPC proliferation and promoting their differentiation. Other members of the Forkhead gene family have roles in OL development. *Foxj3* seems to have the opposite function to *Foxb1*, by stimulating OPC proliferation (Dugas et al., 2010). *Foxa2* is essential for the induction and/or differentiation of OL in zebrafish (Norton et al., 2005) and *Foxg1* inhibits the glial fate choice (including oligodendroglia; Brancaccio et al., 2010).

Our results suggest that *Foxb1* would be one of many components in a cascade of genetic activation regulating a series of choices made by NPCs and the OPCs that derive from them. This function of *Foxb1* would be exerted in the spatially restricted groups of NPCs expressing this gene, the ones giving rise to cell groups in the hindbrain and thalamus (and possibly also in the mammillary body (MBO) of the hypothalamus). Because other Fox genes are involved in similar functions in the cortex, we can assume that, to a degree, different restricted regions of neuroepithelium regulate the generation of the corresponding brain cells by using different members of the same gene families. Each of these members might perform a similar job but with some differences upon which the morphological and functional heterogeneity of CNS regions in the adult would ultimately be based.

Foxb1 is widely expressed in the neuroepithelium of the thalamus, hypothalamus, hindbrain and spinal cord (Alvarez-Bolado et al., 1999; Zhao et al., 2007, 2008). Accordingly, we have observed clusters of *Foxb1* lineage in thalamus, MBO (hypothalamus), pons and MO (we did not investigate the spinal cord) but not, for instance, in the cortex (**Figure 6A**). Therefore, here we uncover specific, spatially restricted subpopulations of OPC and OL characterized by belonging to the *Foxb1* lineage and whose proliferation is modulated by *Foxb1*. In the rodent CNS, OPC originate in discrete loci of the dorsal and ventral neuroepithelium (Woodruff et al., 2001; Rowitch, 2004; Kessaris et al., 2006; Rowitch and Kriegstein, 2010; Huang et al., 2013). Origin heterogeneity has some correlate in cellular properties like proliferation (Young et al., 2013) and marker expression (Butt et al., 1998b; Kleopa et al., 2004). Besides, different morphological classes of OL (Bjartmar et al., 1994; Butt et al., 1998a; Anderson et al., 1999) have been described, perhaps related to place of origin.

To address the question of the functional equivalence of OL of heterogeneous origins, we compared the main characteristics of lineage and phenotype between MO and thalamus. In both regions the *Foxb1* lineage consists mostly of neurons and OL, together with some astrocytes. Interestingly, a robust portion of the thalamo-cortical bundle tract was labeled by our reporter (**Figure 6A**), indicating that a large number of thalamo-cortical projection neurons belong to the *Foxb1* lineage. The *Foxb1* null phenotype is similar in both regions: reduction of neurons, astrocytes and mature OL in favor of immature OL. In principle, this agrees with previous work indicating that OL from OPC originated in different CNS regions are not very different functionally (Kessaris et al., 2006; Psachoulia et al., 2009; Tripathi et al., 2011; Clarke et al., 2012; although white matter and gray matter OL are functionally different from each other; Viganò et al., 2013). But the two subpopulations that we have compared share *Foxb1* lineage. Obviously OL from non-*Foxb1*-expressing regions will express different proliferation modulators. This implies that there must be a series of restricted neuroepithelial “patches” differentially expressing transcription factors which specifically modulate aspects of OL production.

Finally, a myelination phenotype could in principle be expected in the *Foxb1* mouse mutants. However, neurological signs demyelination (like ataxia, unsteady gait, ocular paralysis, weakness and loss of sensation) have not been observed by us or other authors in these mutants. In addition, the G-ratio remained unchanged in the homozygous brainstem. We conclude that *Foxb1* is not required for normal myelination.

CONCLUSION

Forkhead transcription factor *Foxb1* is a novel modulator of neuronal and OL generation acting in certain restricted CNS regions. *Foxb1* biases NPCs towards the neuronal and to a lesser degree the astrocytic lineages and inhibits the proliferation of

OPC, favoring their differentiation. *Foxb1*-lineage OL are a novel subpopulation whose specific properties as well as their ability to replace other subpopulations (for instance, telencephalic OL) as part of a therapeutic approach are open questions.

AUTHOR CONTRIBUTIONS

YZ and EH performed experiments and analyzed results; TZ made the original observations and performed key preliminary experiments; XZ generated the *Foxb1-Cre* mutant line, assisted with the experiments and analyzed results; GA-B and XZ directed the project; YZ, EH and GA-B wrote the article; all authors revised the manuscript and approved the final version.

REFERENCES

- Alvarez-Bolado, G., Cecconi, F., Wehr, R., and Gruss, P. (1999). The fork head transcription factor Fkh5/Mf3 is a developmental marker gene for superior colliculus layers and derivatives of the hindbrain somatic afferent zone. *Dev. Brain Res.* 112, 205–215. doi: 10.1016/s0165-3806(98)00176-x
- Alvarez-Bolado, G., Zhou, X., Voss, A. K., Thomas, T., and Gruss, P. (2000). Winged helix transcription factor Foxb1 is essential for access of mammillothalamic axons to the thalamus. *Development* 127, 1029–1038.
- Anderson, E. S., Bjartmar, C., Westermark, G., and Hildebrand, C. (1999). Molecular heterogeneity of oligodendrocytes in chicken white matter. *Glia* 27, 15–21. doi: 10.1002/(SICI)1098-1136(199907)27:1<15::AID-GLIA2>3.0.CO;2-I
- Ang, S. L., Wierda, A., Wong, D., Stevens, K. A., Cascio, S., Rossant, J., et al. (1993). The formation and maintenance of the definitive endoderm lineage in the mouse: involvement of HNF3/forkhead proteins. *Development* 119, 1301–1315.
- Battiste, J., Helms, A. W., Kim, E. J., Savage, T. K., Lagace, D. C., Mandyam, C. D., et al. (2007). Ascl1 defines sequentially generated lineage-restricted neuronal and oligodendrocyte precursor cells in the spinal cord. *Development* 134, 285–393. doi: 10.1242/dev.02727
- Bjartmar, C., Hildebrand, C., and Loinder, K. (1994). Morphological heterogeneity of rat oligodendrocytes: electron microscopic studies on serial sections. *Glia* 11, 235–244. doi: 10.1002/glia.440110304
- Brancaccio, M., Pivetta, C., Granzotto, M., Filippis, C., and Mallamaci, A. (2010). Emx2 and Foxg1 inhibit gliogenesis and promote neurogenesis. *Stem Cells* 28, 1206–1218. doi: 10.1002/stem.443
- Bronstein, J. M., Tiwari-Woodruff, S., Buznikov, A. G., and Stevens, D. B. (2000). Involvement of OSP/claudin-11 in oligodendrocyte membrane interactions: role in biology and disease. *J. Neurosci. Res.* 59, 706–711. doi: 10.1002/(SICI)1097-4547(20000315)59:6<706::AID-JNR2>3.0.CO;2-D
- Bu, J., Banki, A., Wu, Q., and Nishiyama, A. (2004). Increased NG2⁺ glial cell proliferation and oligodendrocyte generation in the hypomyelinating mutant shiverer. *Glia* 48, 51–63. doi: 10.1002/glia.20055
- Butt, A. M., Ibrahim, M., and Berry, M. (1998a). Axon-myelin sheath relations of oligodendrocyte unit phenotypes in the adult rat anterior medullary velum. *J. Neurocytol.* 27, 259–269.
- Butt, A. M., Ibrahim, M., Gregson, N., and Berry, M. (1998b). Differential expression of the L- and S-isoforms of myelin associated glycoprotein (MAG) in oligodendrocyte unit phenotypes in the adult rat anterior medullary velum. *J. Neurocytol.* 27, 271–280. doi: 10.1023/A:1006996713413
- Carlsson, P., and Mahlapuu, M. (2002). Forkhead transcription factors: key players in development and metabolism. *Dev. Biol.* 250, 1–23. doi: 10.1006/dbio.2002.0780
- Clarke, L. E., Young, K. M., Hamilton, N. B., Li, H., Richardson, W. D., and Attwell, D. (2012). Properties and fate of oligodendrocyte progenitor cells in the corpus callosum, motor cortex and piriform cortex of the mouse. *J. Neurosci.* 32, 8173–8185. doi: 10.1523/JNEUROSCI.0928-12.2012
- Davies, J. E., and Miller, R. H. (2001). Local sonic hedgehog signaling regulates oligodendrocyte precursor appearance in multiple ventricular zone domains in the chick metencephalon. *Dev. Biol.* 233, 513–525. doi: 10.1006/dbio.2001.0224
- de Castro, F., Bribián, A., and Ortega, M. C. (2013). Regulation of oligodendrocyte precursor migration during development, in adulthood and in pathology. *Cell. Mol. Life Sci.* 70, 4355–4368. doi: 10.1007/s00018-013-1365-6
- Dou, C., Ye, X., Stewart, C., Lai, E., and Li, S. C. (1997). TWH regulates the development of subsets of spinal cord neurons. *Neuron* 18, 539–551. doi: 10.1016/s0896-6273(00)80296-2
- Dugas, J. C., Cuellar, T. L., Scholze, A., Ason, B., Ibrahim, A., Emery, B., et al. (2010). Dicer1 and miR-219 Are required for normal oligodendrocyte differentiation and myelination. *Neuron* 65, 597–611. doi: 10.1016/j.neuron.2010.01.027
- Emery, B. (2010). Regulation of oligodendrocyte differentiation and myelination. *Science* 330, 779–782. doi: 10.1126/science.1190927
- Friede, R. L., and Bischhausen, R. (1982). How are sheath dimensions affected by axon caliber and internode length. *Brain Res.* 235, 335–350. doi: 10.1016/0006-8993(82)91012-5
- Gierut, J. J., Jacks, T. E., and Haigis, K. M. (2014). Whole-mount X-Gal staining of mouse tissues. *Cold Spring Harb. Protoc.* 2014, 417–419. doi: 10.1101/pdb.prot073452
- Goldman, S. A., and Kuypers, N. J. (2015). How to make an oligodendrocyte. *Development* 142, 3983–3995. doi: 10.1242/dev.126409
- Haddad-Tóvolli, R., Szabó, N. E., Zhou, X., and Alvarez-Bolado, G. (2013). Genetic manipulation of the mouse developing hypothalamus through in utero electroporation. *J. Vis. Exp.* 77:e50412. doi: 10.3791/50412
- Hart, I. K., Richardson, W. D., Heldin, C. H., Westermark, B., and Raff, M. C. (1989). PDGF receptors on cells of the oligodendrocyte-type-2 astrocyte (O-2A) cell lineage. *Development* 105, 595–603.
- Huang, H., Zhao, X. F., Zheng, K., and Qiu, M. (2013). Regulation of the timing of oligodendrocyte differentiation: mechanisms and perspectives. *Neurosci. Bull.* 29, 155–164. doi: 10.1007/s12264-013-1314-2
- Kaestner, K. H., Lee, K. H., Schlöndorff, J., Hiemisch, H., Monaghan, A. P., and Schutz, G. (1993). Six members of the mouse forkhead gene family are developmentally regulated. *Proc. Natl. Acad. Sci. U S A* 90, 7628–7631. doi: 10.1073/pnas.90.16.7628
- Kaestner, K. H., Schütz, G., and Monaghan, A. P. (1996). Expression of the winged helix genes fkh-4 and fkh-5 defines domains in the central nervous system. *Mech. Dev.* 55, 221–230. doi: 10.1016/0925-4773(96)00507-2
- Kessar, N., Fogarty, M., Iannarelli, P., Grist, M., Wegner, M., and Richardson, W. D. (2006). Competing waves of oligodendrocytes in the forebrain and postnatal elimination of an embryonic lineage. *Nat. Neurosci.* 9, 173–179. doi: 10.1038/nn1620

FUNDING

This research was supported by the National Nature Science Foundation of China (31571510 to TZ), National Key Basic Research Program of China (2015CB942800 to TZ) and Chongqing Science and Technology Commission (cstc2014jcyjA10045 to TZ), and by the Institute of Anatomy and Cell Biology and the Department of Neuroanatomy of the University of Heidelberg.

ACKNOWLEDGMENTS

We thank the members of the Institute of Anatomy and Cell Biology of the University of Heidelberg for stimulating discussion and intellectual support.

- Kleopa, K. A., Orthmann, J. L., Enriquez, A., Paul, D. L., and Scherer, S. S. (2004). Unique distributions of the gap junction proteins connexin29, connexin32, and connexin47 in oligodendrocytes. *Glia* 47, 346–357. doi: 10.1002/glia.20043
- Kloetzli, J. M., Fontaine-Glover, I. A., Brown, E. R., Kuo, M., and Labosky, P. A. (2001). The winged helix gene, Foxb1, controls development of mammary glands and regions of the CNS that regulate the milk-ejection reflex. *Genesis* 29, 60–71. doi: 10.1002/1526-968x(200102)29:2<60::AID-GENE1006>3.0.CO;2-L
- Kriegstein, A., and Alvarez-Buylla, A. (2009). The glial nature of embryonic and adult neural stem cells. *Annu. Rev. Neurosci.* 32, 149–184. doi: 10.1146/annurev.neuro.051508.135600
- Labosky, P. A., Winnier, G. E., Jetton, T. L., Hargett, L., Ryan, A. K., Rosenfeld, M. G., et al. (1997). The winged helix gene, Mf3, is required for normal development of the diencephalon and midbrain, postnatal growth and the milk-ejection reflex. *Development* 124, 1263–1274.
- Lam, E. W., Brosens, J. J., Gomes, A. R., and Koo, C. Y. (2013). Forkhead box proteins: tuning forks for transcriptional harmony. *Nat. Rev. Cancer* 13, 482–495. doi: 10.1038/nrc3539
- Lobe, C. G., Koop, K. E., Kreppner, W., Lomeli, H., Gertsenstein, M., and Nagy, A. (1999). Z/AP, a double reporter for cre-mediated recombination. *Dev. Biol.* 208, 281–292. doi: 10.1006/dbio.1999.9209
- Marinelli, C., Bertalot, T., Zusso, M., Skaper, S. D., and Giusti, P. (2016). Systematic review of pharmacological properties of the oligodendrocyte lineage. *Front. Cell. Neurosci.* 10:27. doi: 10.3389/fncel.2016.00027
- Martynoga, B., Drechsel, D., and Guillemot, F. (2012). Molecular control of neurogenesis: a view from the mammalian cerebral cortex. *Cold Spring Harb. Perspect. Biol.* 4:a008359. doi: 10.1101/cshperspect.a008359
- Mitew, S., Hay, C. M., Peckham, H., Xiao, J., Koenning, M., and Emery, B. (2014). Mechanisms regulating the development of oligodendrocytes and central nervous system myelin. *Neuroscience* 276, 29–47. doi: 10.1016/j.neuroscience.2013.11.029
- Noll, E., and Miller, R. H. (1993). Oligodendrocyte precursors originate at the ventral ventricular zone dorsal to the ventral midline region in the embryonic rat spinal cord. *Development* 118, 563–573.
- Norton, W. H., Mangoli, M., Lele, Z., Pogoda, H. M., Diamond, B., Mercurio, S., et al. (2005). Monorail/Foxa2 regulates floorplate differentiation and specification of oligodendrocytes, serotonergic raphé neurones and cranial motoneurons. *Development* 132, 645–658. doi: 10.1242/dev.01611
- Pinto, L., and Götz, M. (2007). Radial glial cell heterogeneity: the source of diverse progeny in the CNS. *Prog. Neurobiol.* 83, 2–23. doi: 10.1016/j.pneurobio.2007.02.010
- Psachoulia, K., Jamen, F., Young, K. M., and Richardson, W. D. (2009). Cell cycle dynamics of NG2 cells in the postnatal and ageing brain. *Neuron Glia Biol.* 5, 57–67. doi: 10.1017/S1740925X09990354
- Radyushkin, K., Anokhin, K., Meyer, B. I., Jiang, Q., Alvarez-Bolado, G., and Gruss, P. (2005). Genetic ablation of the mammillary bodies in the Foxb1 mutant mouse leads to selective deficit of spatial working memory. *Eur. J. Neurosci.* 21, 219–229. doi: 10.1111/j.1460-9568.2004.03844.x
- Richardson, W. D., Pringle, N., Mosley, M. J., Westermark, B., and Dubois-Dalcq, M. (1988). A role for platelet-derived growth factor in normal gliogenesis in the central nervous system. *Cell* 53, 309–319. doi: 10.1016/0092-8674(88)90392-3
- Rowitch, D. H. (2004). Glial specification in the vertebrate neural tube. *Nat. Rev. Neurosci.* 5, 409–419. doi: 10.1038/nrn1389
- Rowitch, D. H., and Kriegstein, A. R. (2010). Developmental genetics of vertebrate glial-cell specification. *Nature* 468, 214–222. doi: 10.1038/nature09611
- Soriano, P. (1999). Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat. Genet.* 21, 70–71. doi: 10.1038/5007
- Stern, P., Astrof, S., Erkeland, S. J., Schustak, J., Sharp, P. A., and Hynes, R. O. (2008). A system for Cre-regulated RNA interference *in vivo*. *Proc. Natl. Acad. Sci. U S A* 105, 13895–13900. doi: 10.1073/pnas.0806907105
- Tabata, H., and Nakajima, K. (2001). Efficient *in utero* gene transfer system to the developing mouse brain using electroporation: visualization of neuronal migration in the developing cortex. *Neuroscience* 103, 865–872. doi: 10.1016/s0306-4522(01)00016-1
- Tripathi, R. B., Clarke, L. E., Burzomato, V., Kessar, N., Anderson, P. N., Attwell, D., et al. (2011). Dorsally and ventrally derived oligodendrocytes have similar electrical properties but myelinate preferred tracts. *J. Neurosci.* 31, 6809–6819. doi: 10.1523/JNEUROSCI.6474-10.2011
- Vallstedt, A., Klos, J. M., and Ericson, J. (2005). Multiple dorsoventral origins of oligodendrocyte generation in the spinal cord and hindbrain. *Neuron* 45, 55–67. doi: 10.1016/j.neuron.2004.12.026
- Viganò, F., Möbius, W., Götz, M., and Dimou, L. (2013). Transplantation reveals regional differences in oligodendrocyte differentiation in the adult brain. *Nat. Neurosci.* 16, 1370–1372. doi: 10.1038/nn.3503
- Weigel, D., and Jäckle, H. (1990). The fork head domain: a novel DNA binding motif of eukaryotic transcription factors? *Cell* 63, 455–456. doi: 10.1016/0092-8674(90)90439-1
- Woodruff, R. H., Tekki-Kessar, N., Stiles, C. D., Rowitch, D. H., and Richardson, W. D. (2001). Oligodendrocyte development in the spinal cord and telencephalon: common themes and new perspectives. *Int. J. Dev. Neurosci.* 19, 379–385. doi: 10.1016/s0736-5748(00)00083-6
- Young, K. M., Psachoulia, K., Tripathi, R. B., Dunn, S. J., Cossell, L., Attwell, D., et al. (2013). Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling. *Neuron* 77, 873–885. doi: 10.1016/j.neuron.2013.01.006
- Zhao, T., Szabó, N., Ma, J., Luo, L., Zhou, X., and Alvarez-Bolado, G. (2008). Genetic mapping of Foxb1-cell lineage shows migration from caudal diencephalon to telencephalon and lateral hypothalamus. *Eur. J. Neurosci.* 28, 1941–1955. doi: 10.1111/j.1460-9568.2008.06503.x
- Zhao, T., Zhou, X., Szabó, N., Leitges, M., and Alvarez-Bolado, G. (2007). Foxb1-driven Cre expression in somites and the neuroepithelium of diencephalon, brainstem and spinal cord. *Genesis* 45, 781–787. doi: 10.1002/dvg.20356
- Zhou, Q., Wang, S., and Anderson, D. J. (2000). Identification of a novel family of oligodendrocyte lineage-specific basic helix-loop-helix transcription factors. *Neuron* 25, 331–343. doi: 10.1016/s0896-6273(00)80898-3

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Zhang, Hoxha, Zhao, Zhou and Alvarez-Bolado. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.