

OPEN ACCESS

APPROVED BY

Frontiers Editorial Office, Frontiers Media SA, Switzerland

*CORRESPONDENCE

Kathleen E. Whitlock

☑ kathleen.whitlock@uv.cl

†PRESENT ADDRESSES

M. Fernanda Palominos.

Department of Integrative Biology, University of California, Berkeley, Berkeley, CA, United States; Museum of Vertebrate Zoology, University of California, Berkeley, Berkeley, CA, United States

RECEIVED 28 January 2023 ACCEPTED 01 February 2023 PUBLISHED 23 March 2023

CITATION

Whitlock KE and Palominos MF (2023) Corrigendum: The olfactory tract: Basis for future evolution in response to rapidly changing ecological niches. *Front. Neuroanat.* 17:1153062. doi: 10.3389/fnana.2023.1153062

COPYRIGHT

© 2023 Whitlock and Palominos. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: The olfactory tract: Basis for future evolution in response to rapidly changing ecological niches

Kathleen E. Whitlock^{1,2*} and M. Fernanda Palominos^{1,2†}

¹Centro Interdisciplinario de Neurociencia de Valparaíso (CINV), Universidad de Valparaíso, Valparaíso, Chile, ²Instituto de Neurociencia, Universidad de Valparaíso, Valparaíso, Chile

KEYWORDS

gonadotropin releasing hormone (GnRH), immune system, neutrophils, climate change, limbic system, teleost fishes

A corrigendum on

The olfactory tract: Basis for future evolution in response to rapidly changing ecological niches

by Whitlock, K. E., and Palominos, M. F. (2022). Front. Neuroanat. 16:831602. doi: 10.3389/fnana.2022.831602

In the published article there was a missing citation in Figure 1. Figure 1A was adapted from Calvo-Ochoa and Byrd-Jacobs (2019). The corrected Figure 1 caption is below.

Figure 1. The connections from the peripheral olfactory epithelia to the olfactory bulbs are highly conserved in vertebrates. In both in teleost fish (A, zebrafish: modified from - Calvo-Ochoa and Byrd-Jacobs, 2019) and humans (B) the OSNs relay information to the olfactory bulbs (blue) continuing to the dorsal pallium in fishes (A), and the olfactory cortex/lateral pallium (B) in mammals, thus bypassing the thalamus (orange). Both species have projections from the olfactory bulbs (blue) to the amygdala (red, B) and its proposed equivalent in teleosts, the dorsomedial pallium (red, A).

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Whitlock and Palominos 10.3389/fnana.2023.1153062

References

Calvo-Ochoa, E., and Byrd-Jacobs, C. A. (2019). The olfactory system of zebrafish as a model for the study of neurotoxicity and injury: implications

for neuroplasticity and disease. Int. J. Mol. Sci. 20, 1639. doi: 10.3390/ijm s20071639