



Effects of vaccination with altered peptide ligand on chronic pain in experimental autoimmune encephalomyelitis, an animal model of multiple sclerosis

David H. Tian¹, Chamini J. Perera¹, Vasso Apostolopoulos² and Gila Moalem-Taylor^{1*}

¹ School of Medical Sciences, University of New South Wales, Sydney, NSW, Australia

² VA Consulting Services, Melbourne, VIC, Australia

Edited by:

V. Wee Yong, University of Calgary, Canada

Reviewed by:

V. Wee Yong, University of Calgary, Canada

Bradley Kerr, University of Alberta, Canada

*Correspondence:

Gila Moalem-Taylor, School of Medical Sciences, University of New South Wales, Wallace Wurth Building East, Level 3, Room 327, Sydney, NSW 2052, Australia
e-mail: gila@unsw.edu.au

Neuropathic pain is a chronic symptom of multiple sclerosis (MS) and affects nearly half of all MS sufferers. A key instigator of this pain is the pro-inflammatory response in MS. We investigated the behavioral effects of immunization with a mutant peptide of myelin basic protein (MBP), termed altered peptide ligand (APL), known to initiate immune deviation from a pro-inflammatory state to an anti-inflammatory response in experimental autoimmune encephalomyelitis (EAE), an animal model of MS. Male and female Lewis rats were injected with vehicle control or with varying doses of 50 or 100 μ g guinea pig MBP in combination with or without APL. APL-treated animals established significantly lower disease severity compared to encephalitogenic MBP-treated animals. Animals with EAE developed mechanical, but not thermal pain hypersensitivity. Mechanical pain sensitivities were either improved or normalized during periods of clinical disease in male and female APL-treated animals as compared to the encephalitogenic group. No significant changes to thermal latency were observed upon co-immunization with APL. Together these data indicate that APL ameliorates disease states and selectively mediates an analgesic effect on EAE animals.

Keywords: experimental autoimmune encephalomyelitis, altered peptide ligand, multiple sclerosis, mechanical allodynia, thermal hyperalgesia, nociception

INTRODUCTION

Multiple sclerosis (MS) is a chronic, T cell-mediated autoimmune neurological disease of the central nervous system (CNS) (1, 2), characterized by the production of acute multifocal CNS lesions with concurrent perivenular inflammation, demyelination, neuronal degeneration, and gliosis in gray and white matter (3, 4). While the cause of the disease is not known, pro-inflammatory CD4⁺ T cells, CD8⁺ T cells, B cells, macrophages, and natural killer cells have been implicated in disease onset and progression (4, 5). In addition, the dichotomous actions of pro- and anti-inflammatory cytokines are known to play a major role in disease exacerbation and amelioration respectively (6).

Neuropathic pain is a key clinical symptom in MS, with significant interference of quality of life (7–10). It results from damage to the nervous system (11), and presents in various forms such as ongoing extremity pain, paroxysmal neuropathic pain (e.g., trigeminal neuralgia, Lhermitte's phenomenon), hyperalgesia (increased sensitivity to pain), and allodynia (pain produced by innocuous stimuli) (8, 12–14). While the mechanisms underlying MS-related neuropathic pain are not fully understood, lesions of CNS areas associated with pain, activation of T lymphocytes, and pro-inflammatory responses have been shown to contribute to the development and maintenance of neuropathic pain (15–17).

Experimental autoimmune encephalomyelitis (EAE) is a well-established and ubiquitous animal model that exhibits close

clinical and histopathological similarities to various forms of MS (18). Following induction in susceptible animals by injections with a self-antigenic myelin-derived peptide, such as myelin basic protein (MBP), priming of antigen-specific lymphocytes occurs in the periphery. These cells subsequently migrate to the CNS, where they produce inflammatory mediators and cytokines that damage the myelin and axons and activate resident microglia to attract more inflammatory cells, resulting in inflammatory demyelination and neurodegeneration (19). Recent studies in rodents have demonstrated that animals with EAE develop neuropathic pain behaviors, including tactile and cold allodynia, and mechanical and thermal hyperalgesia (20–22). These symptoms have been observed before, during, and after neurological impairment, depending on the strain and the model used (17).

Epitopes derived from autoantigens involved in the autoimmune pathogenesis can be modified to modulate their immunological properties, and are called altered peptide ligands (APLs). APLs are similar to the immunogenic peptide, but with one or more amino acid substitutions in the essential contact positions, with the T cell receptor interfering with the T cell activation. Thus, APLs have the capacity to affect T cell receptor-mediated effector functions (23, 24), such as conferring an anergic effect on specific T cell subsets, or rendering them unresponsive to specific antigens despite presence of functioning antigen presenting cells (25). Different signaling mechanisms can

also be activated to initiate a functional change in the T cell phenotype (24, 26), thereby altering cytokine production and downstream mechanisms (27) to selectively down-regulate pro-inflammatory T_H1 cells (but not T_H2 cells), as well as selectively inducing T lymphocytes that produce T_H2 and T_H0 cytokines (28). The ability for APLs to divert immune responses to a T_H2 profile has been validated in several studies (26, 29–31). Importantly, T_H2 cellular response has been shown to ameliorate EAE mediated by encephalitogenic T_H1 population (28, 29, 31–33), as well as ameliorate pain states (15, 16, 34). As T_H1 cells are widely believed to mediate pro-inflammatory effects, releasing a distinct set of cytokines that exacerbate pain states (35–37), their downregulation could similarly ameliorate clinical symptoms.

Our study builds upon the concept of immune deviation through APLs. While the effectiveness of myelin-derived APL in preventing EAE is well documented, we hypothesize that APL inoculation through immune deviation mediates an analgesic effect in this animal model of MS. Indeed, our results are the first to demonstrate that MBP-derived APL helps to restore EAE-affected mechanical pain thresholds.

MATERIALS AND METHODS

EXPERIMENTAL ANIMALS

Male ($n = 36$) and female ($n = 18$) 6- to 8-week-old Lewis rats (Animal Resource Centre, Perth, WA, Australia) were used. Animals were housed with food and water *ad libitum* under 12-h light cycle, with constant humidity and temperature. Animals that developed mobility impairment were provided with soft jelly foods and easier access to water. Cage beddings changed twice a week, and animals were inspected daily for well-being. All animal experiments were approved by the Animal Care and Ethics Committee of the University of New South Wales, Sydney, NSW, Australia.

PEPTIDES

Encephalitogenic guinea pig MBP (gp-MBP) is known to induce acute T_H1-associated EAE in Lewis rats (38). Doses of 50 or 100 μ g gp-MBP (Sigma-Aldrich, NSW, Australia) were used to induce EAE. Previous studies using site-directed mutagenesis to compare different mutant peptides have shown that the peptide MBP_{87–99}, with double Ala mutations at positions 91,96-[A⁹¹,A⁹⁶]MBP_{87–99}, alters immune responses leading to decrease in EAE severity (39–41). Our study was designed to assess the effects of this mutant peptide on sensory disturbances in EAE animals. In addition, cyclic peptides have been shown to be more stable *in vivo*, and have similar immunological activity to their linear counterparts (42). For increased stability, the [A⁹¹,A⁹⁶]MBP_{87–99} peptide was cyclized from head to tail resulting in cyclo-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99}. The cyclic double mutant cyclo-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99} peptide at a dose of 250 μ g was used as the APL to inhibit EAE. Peptides were synthesized and purchased from Mimotopes Australia with purity greater than 95%.

EAE INDUCTION

To induce EAE, rats were anesthetized (day 0) with 3% Isoflurane in oxygen, and inoculated at the base of the tail with a

single subcutaneous injection of 200 μ L inoculum. The control group was injected with an inoculum containing sterile saline (0.9% NaCl) emulsified with an equal volume of incomplete Freund's adjuvant (IFA; Difco Laboratories, Detroit, MI, USA) supplemented with desiccated 1 mg/mL *Mycobacterium tuberculosis* (strain H37RA, Difco Laboratories). The IFA with *Mycobacterium tuberculosis* is defined as CFA. A second group was injected with either 50 or 100 μ g gp-MBP emulsified in CFA (deemed the MBP group), while a third group was injected with equal dose of gp-MBP, but an additional 250 μ g cyclo-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99} (APL group).

CLINICAL ASSESSMENT

Following disease induction, animals were assessed daily for signs of disease for up to 35 days, graded using the following scale: Grade 0, normal rat; Grade 1, flaccid tail; Grade 2, weak hind limbs with ataxia; Grade 3, hind limb paralysis; Grade 4, paraplegia with forelimb paralysis; Grade 5, moribund. Weight changes were also recorded.

PAIN ASSESSMENT

Behavioral tests were conducted three times a week prior to and up to 4 weeks following EAE induction.

Prior to testing, animals were acclimatized in a clear Perspex 20 cm \times 20 cm box standing 20 cm above the bench for 15 min until they were in a non-agitated state. Thermal hyperalgesia was assessed by exposing the plantar hind paw of the animal to radiant heat through the transparent floor of the Perspex box, using a plantar analgesia meter (Ugo Basile, Varese, Italy). A cut-off of 20 s was applied to prevent tissue damage.

Mechanical allodynia was then assessed subsequent to 30-min break, by an electronic von Frey anesthesiometer (Ugo Basile). The animal, placed upon an elevated wire mesh surrounded by a Perspex box, was exposed to increasing mechanical pressure to the plantar hind paw through a metal filament.

Withdrawal latency and threshold was measured automatically from the initiation of heat or mechanical stimulus to withdrawal of the paw, defined as sudden jerk of paw away from stimulus. This was repeated three times in both left and right hind paw separated by 2–5 min between each stimulus. Mean results for each animal was calculated.

DATA ANALYSIS

In mechanical and thermal tests, raw scores for left and right hind paws were combined and averaged for each time-point. Time periods were classified as pre-disease, disease peak, and recovery, defined as time-points prior to development of clinical signs, presence of clinical signs, and resolution of clinical signs respectively. Disease peak periods are shaded gray in **Figures 1–3**.

For measurements of pain behaviors, data were analyzed with repeated measures two-way ANOVA with Bonferroni's post-test, and for EAE clinical scores data were analyzed with a non-parametric Mann–Whitney test as appropriate, using GraphPad Prism v5.04 (GraphPad Software, Inc., CA, USA). Significance was set at $p < 0.05$. All data are presented as mean \pm SEM.




FIGURE 1 | Pain hypersensitivity (top set) and clinical score (bottom set) in male Lewis rats injected with 100 μ g gp-MBP on day 0. Top series represent response to stimuli, while bottom series represent clinical score. EAE clinical scores of rats immunized with MBP were significantly greater than those of APL-treated or CFA-immunized (control) animals during period of established disease. **(A)** MBP-treated animals demonstrated a decrease in paw withdrawal thresholds to mechanical stimulus, while APL-treated animals had significantly elevated mechanical

pain thresholds, in particular during disease recovery. **(B)** No significant difference was observed in paw withdrawal latency to thermal stimuli between APL-treated and MBP-treated animals. ($n = 6$ per group; **** $p < 0.0001$, MBP compared to control; * $p < 0.05$, *** $p < 0.001$, APL compared to MBP; two-way ANOVA with Bonferroni's post-tests for upper panel, Mann-Whitney test for lower panel). Gray region indicates periods of established disease. Dotted line represents mean baseline values. Data expressed as mean \pm SEM.




FIGURE 2 | Changes in mechanical sensitivity (top set) in male and female Lewis rats inoculated with 50 μ g gp-MBP on day 0 compared to clinical scores (bottom set). In both males and females, EAE clinical scores of rats immunized with MBP were significantly greater than those of APL-treated or CFA-immunized (control) animals during period of established disease. **(A)** In males, APL conferred significant increase in mechanical threshold compared to MBP-treated rats on day 11, 14, 16. **(B)** In females,

MBP animals generally exhibited lower pain thresholds, while APL animals maintained thresholds similar to control. ($n = 6$ per group; * $p < 0.05$, **** $p < 0.0001$, MBP compared to control; * $p < 0.05$, ** $p < 0.01$, APL compared to MBP; two-way ANOVA with Bonferroni's post-tests for upper panel, Mann-Whitney test for lower panel). Gray region indicates periods of established disease. Dotted line represents mean baseline values. Data expressed as mean \pm SEM.

RESULTS


EFFECTS OF CO-IMMUNIZATION WITH APL, CYCLO-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99}, IN MALE LEWIS RATS INJECTED WITH 100 μ g gp-MBP

We first studied the effects of APL vaccination on EAE clinical disease course and changes to mechanical threshold and thermal latency for pain in 18 male Lewis rats immunized with 100 μ g gp-MBP. Three study groups were used, (i) animals injected with CFA only (vehicle control), (ii) animals injected

with CFA + 100 μ g gp-MBP (MBP group) and, (iii) animals co-immunized with CFA + 100 μ g gp-MBP + 250 μ g cyclo-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99} (APL group).

Treatment with APL significantly reduced disease severity in male animals with EAE

Both MBP-treated and APL-treated animal groups demonstrated characteristic clinical deficits beginning on day 8 (**Figures 1A,B**). Vehicle-treated animals displayed minimal clinical deficits. Onset



was characterized by weakness of tail, followed by ascending motor deficits.

Disease severity in MBP-treated animals peaked later on day 17 with a mean maximal score of 1.7 ± 0.5 ($n = 6$), while the APL cohort peaked earlier on day 13 with 82% comparative reduction in average maximal severity (peak mean score 0.9 ± 0.08). APL animals recovered significantly earlier compared to MBP animals (22.5 days for MBP and 18.8 days for APL, Mann–Whitney test, $p < 0.01$).

Overall, MBP animals demonstrated significantly greater EAE clinical scores than vehicle-treated animals when clinical signs were present between day 8 and 23 ($p < 0.0001$, Mann–Whitney test). Compared to MBP, the APL group showed a significant reduction in clinical scores during the same period ($p < 0.005$, Mann–Whitney test).

Treatment with APL significantly increased thresholds to mechanical stimuli following disease resolution

Prior to disease onset on day 8 (Figure 1A) and during disease establishment (day 8–23, gray region), no significant differences in paw withdrawal thresholds to mechanical stimuli were observed between CFA-injected control, MBP-, and APL-treated animals ($p > 0.05$, two-way ANOVA). Following resolution of clinical disease, MBP animals developed reduced mechanical thresholds, while APL animals exhibited significantly elevated thresholds of as much as 6.90 g compared to MBP animals on days 24 and 26 ($p < 0.05$, two-way ANOVA). Although not statistically significant, compared to control animals, MBP animals generally possessed lower thresholds to mechanical pain, particularly after disease peak, whereas APL animals demonstrated elevated thresholds.

Interestingly, a sharp increase in withdrawal threshold was observed in MBP animals on day 16, corresponding closely to the rats' maximal disease severity (score 1.7). While hindlimb paralysis has the potential to confound results, results for

all three groups became relatively stabilized following animal recovery.

No significant changes to thermal latency were observed in animals co-immunized with APL

In addition to evaluating alterations of mechanical pain thresholds, we examined changes to paw withdrawal latency in response to thermal stimulation (Figure 1B). Interestingly, no difference was observed between any of the cohorts prior to, during, and following disease establishment ($p > 0.05$, two-way ANOVA). We conclude that male animals with EAE do not develop thermal pain hypersensitivity and that APL does not confer any significant effects on latency to thermal stimuli.

EFFECTS OF CO-IMMUNIZATION WITH APL, CYCLO-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99}, IN MALE AND FEMALE RATS INJECTED WITH 50 μ g gp-MBP

To reduce confounding effects of severe hind limb paralysis on the ability of animals to withdraw their paw from the stimulus, we halved the MBP dose to 50 μ g in order to minimize physical paralysis. Since previous studies have demonstrated differences between sexes in EAE severity and nociception, we tested both male and female rats for clinical EAE and pain sensitivity. Here we used 18 male and 18 female Lewis rats immunized with CFA only (control), CFA + 50 μ g gp-MBP (MBP group), and CFA + 50 μ g gp-MBP + 250 μ g cyclo-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99} (APL group).

Treatment with APL reduced disease severity in both male and female animals with EAE

Disease onset in male MBP animals occurred around day 10 and in female MBP animals around day 8 (Figures 2A,B). Additionally, the disease course of male animals peaked later (day 16) than female animals (day 14). Recovery for both sexes occurred on day 23.

Significant differences between the disease course of male and female MBP animals were observed. Female animals developed clinical signs that were on average 151% more severe than male animals (result reported as percentage of difference compared to male animals, $p < 0.05$; Mann–Whitney test). For example, three female MBP animals developed scores of 3 (complete paralysis of tail and hind limbs), whilst the male MBP cohorts only managed to develop maximal scores of 1.5 (paralysis of tail only).

Separately, there existed a significant difference in clinical scores between CFA and MBP, and MBP and APL groups in both sexes when clinical signs were present (day 10–22 for males, day 8–22 for females) (Figures 2A,B). Both male and female MBP animals displayed significantly greater EAE clinical scores than CFA animals ($p < 0.0001$ in both, Mann–Whitney test). Both male and female APL animals showed a significant reduction in EAE severity as compared to MBP animals ($p < 0.05$, Mann–Whitney test) during established disease. These results highlight APL's ameliorative effect on disease status in both males and females.

Treatment with APL significantly ameliorated mechanical pain sensitivities in both male and female animals during established disease

In male animals, APL normalized changes in EAE-induced mechanical pain thresholds only during periods when clinical signs were present. No significant difference was observed between the three cohorts prior to and following disease establishment, except a small reduction in threshold in APL-treated animals compared to MBP animals on day 2 ($p < 0.05$, two-way ANOVA). When clinical signs of disease were evident on day 10–22 (Figure 2A, gray region), MBP animals exhibited lower mechanical pain thresholds compared to control, while no significant differences were observed between APL and control ($p > 0.05$, two-way ANOVA). Furthermore, on day 11, 14, and 16, animals co-immunized with APL demonstrated threshold increase of at least 2.6 g against MBP animals ($p < 0.05$, two-way ANOVA) and their paw withdrawal threshold to mechanical stimulus was similar to control animals. This validates the ability for APL to ameliorate changes to mechanical pain thresholds in animals with EAE.

In female animals, APL's ameliorative effects were similarly evident mostly during established disease (Figure 2B, gray region). Prior to disease onset, no significant differences in mechanical threshold were observed between the cohorts ($p > 0.05$, two-way ANOVA). MBP animals developed a sharp decrease in mechanical threshold starting on day 9 following disease induction, followed by a steady increase, potentially associated with the escalating disease severity. The threshold started to reduce again on day 18, following amelioration of paralytic signs. When clinical signs were present between day 8 and 22, MBP animals exhibited a general reduction in threshold, particularly noticeable on day 11 and 21 ($p < 0.05$, two-way ANOVA). During the same period, APL animals displayed similar thresholds to control, which were significantly elevated against MBP on day 18 ($p < 0.05$, two-way ANOVA). Again, this indicates the normalization of mechanical thresholds in APL animals with EAE.

No significant changes to thermal latency were observed in APL-treated animals

In male animals, no significant latency variations to thermal stimuli were observed between any time-points during the course of the experiment (Figure 3A; $p > 0.05$, two-way ANOVA), similar to the results from male animals inoculated with twice the MBP dosage (Figure 1).

In female animals, immunization with only MBP was able to increase thermal latency by 4.8 s on day 14 compared to control (Figure 3B; $p < 0.001$, two-way ANOVA). However, it should be noted that this transient increase occurred concurrent with disease peak, therefore it is possible that animal paralysis has confounded results. At the same time-point, APL animals, which exhibited less severe paralysis, displayed no significant difference compared to control. No other changes were observed during the course of the experiment.

DISCUSSION

Over the last two decades, numerous studies in rodents have demonstrated that several non-encephalitogenic myelin-derived APLs confer protection from the development of EAE, and even reverse established paralytic disease (26, 28, 29). These APLs were shown to induce T cells that are cross-reactive with the native myelin peptide, but modify the immune response and prevent autoimmune encephalomyelitis. Our results here have shown that active immunization with the APL cyclo-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99} in an animal model of EAE, not only mitigated the disease course, but also improved symptoms of mechanical pain hypersensitivity.

We demonstrated that in both male and female Lewis rats, co-immunization with gp-MBP and APL cyclo-(87–99)[A⁹¹,A⁹⁶]MBP_{87–99} has significantly reduced EAE disease severity and shortened the disease course as compared to gp-MBP alone. These results are in line with previous studies (26, 28, 31, 43, 44). We also demonstrated that EAE severity is dependent upon MBP dosage and gender. It is well known that many autoimmune diseases, including MS, are more predisposed toward females than males (45). Our study determined female animals were significantly more affected by equal dosages of MBP, with clinical scores of female rats more than twice that of male animals. A dose of 100 μ g MBP was able to elicit mean peak disease score of 1.7 in male Lewis rats, while 50 μ g MBP was only able to generate considerably lower mean peak score of 0.7. Female rats were more affected by MBP immunization, with only 50 μ g MBP producing a mean peak score of 1.75, comparable to doubling the dosage in male animals. Indeed, it has been shown that immunization with gp-MBP induced disease in all female mice, but only in half of male animals (46). Variations in encephalitogenic peptides and animal strains similarly indicated that female animals were more severely affected (20, 47, 48).

Chronic neuropathic pain arises subsequent to lesion or disease of the somatosensory nervous system (11). Recent studies have shown that T lymphocytes and pro-inflammatory cytokines play a significant role in the development and maintenance of neuropathic pain (34, 49). For example, injection of a T_H1 cell

population producing pro-inflammatory cytokines increased the level of neuropathic pain, whereas injection of a T_H2 cell population producing anti-inflammatory cytokines attenuated pain sensitivity in nerve-injured rats (15). As chronic neuropathic pain affects the majority of MS patients (10, 17), it is believed that modulation of pro-inflammatory T cells and their associated cytokine response will mitigate such symptoms.

Our results show that concurrent inoculation with APL, in addition to disease-causing MBP, normalizes disturbances to mechanical pain threshold, particularly during established disease, although it had no significant effect on thermal latency to pain. Consistent with previous studies (21, 22, 50, 51), we found that animals with EAE display mechanical allodynia during the course of the disease. In addition, we observed normalization or increase in pain threshold to mechanical stimuli during periods of clinical disease in animals co-immunized with APL. While there existed some association between changes to mechanical threshold and disease severity, particularly in MBP animals, threshold trends persisted even following resolution of clinical paralysis. Overall, treatment with APL displayed a tendency to normalize pain thresholds, and hinder the development of mechanical pain hypersensitivity.

In MS, neuro-inflammatory lesions in the CNS produce significant somatosensory deficit, particularly in temperature discrimination, such as paradoxical heat sensations and altered heat/cold thresholds (52–54). In our experiments with male and female MBP animals, no differences in latency to thermal stimuli were observed, except in transience. Similarly, Olechowski et al. (21) reported no change in sensitivity to noxious heat, albeit using a different encephalitogenic peptide and animal model (21). In contrast, thermal hyperalgesia was observed in the tail and forepaws of male and female SJL mice, using a proteolipid protein from the myelin sheath as immunogenic source. These conflicting results underscore the high variability existent between differing animal models and encephalitogenic peptides. Consequently, we were not able to show any differences caused by APL co-immunization. It should be noted, however, that paralysis of the hind paws could have potentially confounded results, a concern shared by others (20, 21). Additional experimental setups that diminish the impact of paralysis on nociceptive testing, such as measuring vocalization response (55) or spontaneous pain (56) in animals, are encouraged.

As pain has only recently been recognized as a key functional disability of MS, a clear understanding of its pathogenesis is still absent. Several theories exist to explain this pain, including damage to somatosensory nerves (57), lesions in the CNS and spinal cord inflammation (17). However, a key factor is the dichotomous role of pro- and anti-inflammatory responses. Indeed, a recent study has shown that animals with EAE did not have altered expression of sensory neuropeptides, but possessed an influx of CD3⁺ T cells and increased astrocyte and microglia/macrophage reactivity in the superficial dorsal horn of the spinal cord, an area associated with pain processing (21). Additionally, a significant increase in the level of tumor necrosis factor (TNF) expression, a key pro-inflammatory cytokine, in the dorsal root ganglia of EAE animals was found at disease peak (58). Gene

therapy with anti-inflammatory interleukin (IL)-10 resulted in prevention of the onset of allodynia in animals with EAE (50). Collectively, these findings suggest that pro-inflammation and gliosis are key mediators in the neuropathic pain behaviors associated with EAE.

The mechanisms underlying the analgesic effect of APL immunization in EAE-induced mechanical pain hypersensitivity are not known, but may include: reduced production of interferon- γ and TNF, pro-inflammatory cytokines that are critical in the pathogenesis of EAE (29); up-regulation of anti-inflammatory cytokines IL-4, IL-10, IL-13 and transforming growth factor- β (32); diverting immune responses from T_H1 to T_H2 (33); and mediating bystander suppression by the generation of regulatory T cells (59), which have been shown to suppress pain hypersensitivity in nerve-injured animals (60). In addition, APL immunization in EAE animals may have affected other pain mediators such as bradykinin, eicosanoids (prostaglandins and leukotrienes), adenosine-5'-triphosphate (ATP), histamine, chemokines (e.g., chemotactic cytokine ligand 2, fractalkine), neurotrophins (e.g., nerve growth factor, brain-derived neurotrophic factor), and reactive oxygen species to reduce mechanical pain hypersensitivity (16). Future studies will have to investigate the impact of APL treatment on immune modulation and inflammatory mediators associated with EAE pain.

While recent studies have mostly focused on individual single T cell clones in animal models (61), clinical trials have underlined the complexity of APLs. Despite APL's effective suppression and reversal of EAE in rodents (29, 30), human trials reported conflicting results. In one phase II trial using [A⁹¹]MBP_{83–99}, a decrease in the size of new MS lesions on MRI scans was observed in human subjects, but the trial was halted due to hypersensitivity reactions in 9% of patients (62). Crucially, there was no increase in disease exacerbation, although this did present in a similar phase II clinical trial (63). Thus, further clinical use of APLs is considered questionable. However, this avenue of research holds great promise, as the immune changes instigated by the APLs could induce 2–4.5 years of T_H2-directed deviation in humans (64). Experimental trends show that clinical benefit and allergy mitigation is related to the correct dose of APL and route of administration, both of which necessitate further investigation (65).

Although our results highlight the restorative effect of APL on mechanical pain thresholds, further work is required to elucidate the mechanisms behind such changes. Challenges also remain in translating results from animal experiments into human therapies. Dosages need to be accurately titrated to maximize disease reduction while minimizing side effects, particularly T_H2-induced hypersensitivity reactions. However, should this avenue of research yield promising results, it will herald a new field of immune deviation as a therapeutic option to neuropathic pain in MS and similar diseases.

ACKNOWLEDGMENTS

This work was supported by grants from the National Health and Medical Research Council of Australia and the NSW Office for Science and Medical Research to Gila Moalem-Taylor. We thank Cristina Kim for technical assistance.

REFERENCES

- McFarland HF, Martin R. Multiple sclerosis: a complicated picture of autoimmunity. *Nat Immunol* (2007) **8**:913–9. doi:10.1038/nl1507
- Goldenberg MM. Multiple sclerosis review. *Pharmacy and Therapeutics* (2012) **37**:175–84.
- Bennett J, Basivireddy J, Kollar A, Biron KE, Reickmann P, Jefferies WA, et al. Blood-brain barrier disruption and enhanced vascular permeability in the multiple sclerosis model EAE. *J Neuroimmunol* (2010) **229**:180–91. doi:10.1016/j.jneuroim.2010.08.011
- Cohen JA, Chun J. Mechanisms of fingolimod's efficacy and adverse effects in multiple sclerosis. *Ann Neurol* (2011) **69**:759–77. doi:10.1002/ana.22426
- Olson JK, Ludovic Croxford J, Miller SD. Innate and adaptive immune requirements for induction of autoimmune demyelinating disease by molecular mimicry. *Mol Immunol* (2004) **40**:1103–8. doi:10.1016/j.molimm.2003.11.010
- Sospedra M, Martin R. Immunology of multiple sclerosis. *Annu Rev Immunol* (2005) **23**:683–747. doi:10.1146/annurev.immunol.23.021704.115707
- Ehde DM, Gibbons LE, Chwastiak L, Bombardier CH, Sullivan MD, Kraft GH. Chronic pain in a large community sample of persons with multiple sclerosis. *Mult Scler* (2003) **9**:605–11. doi:10.1191/1352458503ms939oa
- Svensden KB, Jensen TS, Overvad K, Hansen HJ, Koch-Henriksen N, Bach FW. Pain in patients with multiple sclerosis: a population-based study. *Arch Neurol* (2003) **60**:1089–94. doi:10.1001/archneur.60.8.1089
- Beiske AG, Pedersen ED, Czujko B, Myhr KM. Pain and sensory complaints in multiple sclerosis. *Eur J Neurol* (2004) **11**:479–82. doi:10.1111/j.1468-1331.2004.00815.x
- O'Connor AB, Schwid SR, Herrmann DN, Markman JD, Dworkin RH. Pain associated with multiple sclerosis: systematic review and proposed classification. *Pain* (2008) **137**:96–111. doi:10.1016/j.pain.2007.08.024
- Treede RD, Jensen TS, Campbell JN, Cruccu G, Dostrovsky JO, Griffin JW, et al. Neuropathic pain: redefinition and a grading system for clinical and research purposes. *Neurology* (2008) **70**:1630–5. doi:10.1212/01.wnl.0000282763.29778.59
- Solaro C, Bricchetto G, Amato MP, Cocco E, Colombo B, D'Aleo G, et al. The prevalence of pain in multiple sclerosis: a multicenter cross-sectional study. *Neurology* (2004) **63**:919–21. doi:10.1212/01.WNL.0000137047.85868.D6
- Solaro C, Uccelli MM. Management of pain in multiple sclerosis: a pharmacological approach. *Nat Rev Neurol* (2011) **7**:519–27. doi:10.1038/nrneuro.2011.120
- Truini A, Galeotti F, Cruccu G. Treating pain in multiple sclerosis. *Expert Opin Pharmacother* (2011) **12**:2355–68. doi:10.1517/14656566.2011.607162
- Moalem G, Xu K, Yu L. T lymphocytes play a role in neuropathic pain following peripheral nerve injury in rats. *Neuroscience* (2004) **129**:767–77. doi:10.1016/j.neuroscience.2004.08.035
- Moalem G, Tracey DJ. Immune and inflammatory mechanisms in neuropathic pain. *Brain Res Rev* (2006) **51**:240–64. doi:10.1016/j.brainresrev.2005.11.004
- Tian DH, Perera CJ, Moalem-Taylor G. Neuropathic pain in animal models of nervous system autoimmune diseases. *Mediators Inflamm* (2013) **2013**:298326. doi:10.1155/2013/298326
- Schreiner B, Heppner FL, Becher B. Modeling multiple sclerosis in laboratory animals. *Semin Immunopathol* (2009) **31**:479–95. doi:10.1007/s00281-009-0181-4
- Constantinescu CS, Farooqi N, O'Brien K, Gran B. Experimental autoimmune encephalomyelitis (EAE) as a model for multiple sclerosis (MS). *Br J Pharmacol* (2011) **164**:1079–106. doi:10.1111/j.1476-5381.2011.01302.x
- Aicher SA, Silverman MB, Winkler CW, Bebo BF Jr. Hyperalgesia in an animal model of multiple sclerosis. *Pain* (2004) **110**:560–70. doi:10.1016/j.pain.2004.03.025
- Olechowski CJ, Truong JJ, Kerr BJ. Neuropathic pain behaviours in a chronic-relapsing model of experimental autoimmune encephalomyelitis (EAE). *Pain* (2009) **141**:156–64. doi:10.1016/j.pain.2008.11.002
- Thibault K, Calvino B, Pezet S. Characterisation of sensory abnormalities observed in an animal model of multiple sclerosis: a behavioural and pharmacological study. *Eur J Pain* (2011) **15**:e1–16. doi:10.1016/j.ejpain.2010.07.010
- Kersh GJ, Allen PM. Structural basis for T cell recognition of altered peptide ligands: a single T cell receptor can productively recognize a large continuum of related ligands. *J Exp Med* (1996) **184**:1259–68. doi:10.1084/jem.184.4.1259
- Sloan-Lancaster J, Allen PM. Altered peptide ligand – induced partial T cell activation: molecular mechanisms and role in T cell biology. *Annu Rev Immunol* (1996) **14**:1–27. doi:10.1146/annurev.immunol.14.1.1
- Sloan-Lancaster J, Evavold BD, Allen PM. Th2 cell clonal anergy as a consequence of partial activation. *J Exp Med* (1994) **180**:1195–205. doi:10.1084/jem.180.4.1195
- Gaur A, Boehme SA, Chalmers D, Crowe PD, Pahuja A, Ling N, et al. Amelioration of relapsing experimental autoimmune encephalomyelitis with altered myelin basic protein peptides involves different cellular mechanisms. *J Neuroimmunol* (1997) **74**:149–58. doi:10.1016/S0165-5728(96)00220-2
- Evavold BD, Allen PM. Separation of IL-4 production from Th cell proliferation by an altered T cell receptor ligand. *Science* (1991) **252**:1308. doi:10.1126/science.1833816
- Nicholson LB, Greer JM, Sobel RA, Lees MB, Kuchroo VK. An altered peptide ligand mediates immune deviation and prevents autoimmune encephalomyelitis. *Immunity* (1995) **3**:397–405. doi:10.1016/1074-7613(95)90169-8
- Karin N, Mitchell DJ, Brocke S, Ling N, Steinman L. Reversal of experimental autoimmune encephalomyelitis by a soluble peptide variant of a myelin basic protein epitope: T cell receptor antagonism and reduction of interferon gamma and tumor necrosis factor alpha production. *J Exp Med* (1994) **180**:2227–37. doi:10.1084/jem.180.6.2227
- Brocke S, Gijbels K, Allegretta M, Ferber I, Piercy C, Blankenstein T, et al. Treatment of experimental encephalomyelitis with a peptide analogue of myelin basic protein. *Nature* (1996) **379**:343–6. doi:10.1038/379343a0
- Fischer FR, Santambrogio L, Luo Y, Bermer MA, Hancock WW, Dorf ME. Modulation of experimental autoimmune encephalomyelitis: effect of altered peptide ligand on chemokine and chemokine receptor expression. *J Neuroimmunol* (2000) **110**:195–208. doi:10.1016/S0165-5728(00)00351-9
- Young D, Lowe L, Booth S, Whitters M, Nicholson LB, Kuchroo VK, et al. IL-4, IL-10, IL-13, and TGF-β from an altered peptide ligand-specific Th2 cell clone down-regulate adoptive transfer of experimental autoimmune encephalomyelitis. *J Immunol* (2000) **164**:3563–72.
- Katsara M, Yuriev E, Ramsland PA, Deraos G, Tselios T, Matsoukas J, et al. Mannosylation of mutated MBP83-99 peptides diverts immune responses from Th1 to Th2. *Mol Immunol* (2008) **45**:3661–70. doi:10.1016/j.molimm.2008.04.024
- Austin PJ, Moalem-Taylor G. The neuro-immune balance in neuropathic pain: involvement of inflammatory immune cells, immune-like glial cells and cytokines. *J Neuroimmunol* (2010) **229**:26–50. doi:10.1016/j.jneuroim.2010.08.013
- Mosmann TR, Sad S. The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol Today* (1996) **17**:138–46. doi:10.1016/0167-5699(96)80606-2
- Cata JP, Weng HR, Dougherty PM. Spinal injection of IL-2 or IL-15 alters mechanical and thermal withdrawal thresholds in rats. *Neurosci Lett* (2008) **437**:45–9. doi:10.1016/j.neulet.2008.03.074
- Andrade P, Visser-Vandewalle V, Hoffmann C, Steinbusch HW, Daemen MA, Hoogland G. Role of TNF-alpha during central sensitization in preclinical studies. *Neurosci* (2011) **32**:757–71. doi:10.1007/s10072-011-0599-z
- Pender MP. The pathophysiology of myelin basic protein-induced acute experimental allergic encephalomyelitis in the Lewis rat. *J Neurol Sci* (1988) **86**:277–89. doi:10.1016/0022-510X(88)90105-0
- Tselios T, Apostolopoulos V, Daliani I, Deraos S, Grdadolnik S, Mavromoustakos T, et al. Antagonistic effects of human cyclic MBP87-99 altered peptide ligands in experimental allergic encephalomyelitis and human T-cell proliferation. *J Med Chem* (2002) **45**:275–83. doi:10.1021/jm0102147
- Katsara M, Deraos G, Tselios T, Matsoukas MT, Friligou I, Matsoukas J, et al. Design and synthesis of a cyclic double mutant peptide (cyclo(87-99)[A91,A96]MBP87-99) induces altered responses in mice after conjugation to mannan: implications in the immunotherapy of multiple sclerosis. *J Med Chem* (2009) **52**:214–8. doi:10.1021/jm801250v
- Katsara M, Yuriev E, Ramsland PA, Tselios T, Deraos G, Lourdopoulos A, et al. Altered peptide ligands of myelin basic protein (MBP87-99) conjugated to

- reduced mannan modulate immune responses in mice. *Immunology* (2009) **128**:521–33. doi:10.1111/j.1365-2567.2009.03137.x
42. Katsara M, Tselios T, Deraos S, Deraos G, Matsoukas MT, Lazoura E, et al. Round and round we go: cyclic peptides in disease. *Curr Med Chem* (2006) **13**:2221–32. doi:10.2174/09298670677935113
 43. Santambrogio L, Lees MB, Sobel RA. Altered peptide ligand modulation of experimental allergic encephalomyelitis: immune responses within the CNS. *J Neuroimmunol* (1998) **81**:1–13. doi:10.1016/S0165-5728(97)00138-0
 44. Ruiz PJ, Garren H, Hirschberg DL, Langer-Gould AM, Levite M, Karpuz MV, et al. Microbial epitopes act as altered peptide ligands to prevent experimental autoimmune encephalomyelitis. *J Exp Med* (1999) **189**:1275–83. doi:10.1084/jem.189.8.1275
 45. Koch-Henriksen N, Sorensen PS. The changing demographic pattern of multiple sclerosis epidemiology. *Lancet Neurol* (2010) **9**:520–32. doi:10.1016/S1474-4422(10)70064-8
 46. Fritz RB, Chou CH, McFarlin DE. Relapsing murine experimental allergic encephalomyelitis induced by myelin basic protein. *J Immunol* (1983) **130**:1024–6.
 47. Bebo B, Vandenbark A, Offner H. Male SJL mice do not relapse after induction of EAE with PLP 139–151. *J Neurosci Res* (1996) **45**:680–9. doi:10.1002/(SICI)1097-4547(19960915)45:6<680::AID-JNR4>3.0.CO;2-4
 48. Dalal M, Kim S, Voskuhl RR. Testosterone therapy ameliorates experimental autoimmune encephalomyelitis and induces a T helper 2 bias in the autoantigen-specific T lymphocyte response. *J Immunol* (1997) **159**:3–6.
 49. Sommer C, Galbraith JA, Heckman HM, Myers RR. Pathology of experimental compression neuropathy producing hyperesthesia. *J Neuropathol Exp Neurol* (1993) **52**:223. doi:10.1097/00005072-199305000-00006
 50. Sloane E, Ledeboer A, Seibert W, Coats B, Van Strien M, Maier SF, et al. Anti-inflammatory cytokine gene therapy decreases sensory and motor dysfunction in experimental Multiple Sclerosis: MOG-EAE behavioral and anatomical symptom treatment with cytokine gene therapy. *Brain Behav Immun* (2009) **23**:92–100. doi:10.1016/j.bbi.2008.09.004
 51. Ramos KM, Lewis MT, Morgan KN, Crysdale NY, Kroll JL, Taylor FR, et al. Spinal upregulation of glutamate transporter GLT-1 by ceftriaxone: therapeutic efficacy in a range of experimental nervous system disorders. *Neuroscience* (2010) **169**:1888–900. doi:10.1016/j.neuroscience.2010.06.014
 52. Hansen C, Hopf HC, Treede RD. Paradoxical heat sensation in patients with multiple sclerosis. Evidence for a supraspinal integration of temperature sensation. *Brain* (1996) **119**(Pt 5):1729–36. doi:10.1093/brain/119.5.1729
 53. Bowsher D. Central pain following spinal and supraspinal lesions. *Spinal Cord* (1999) **37**:235–8. doi:10.1038/sj.sc.3100832
 54. Osterberg A, Boivie J. Central pain in multiple sclerosis – sensory abnormalities. *Eur J Pain* (2010) **14**:104–10. doi:10.1016/j.ejpain.2009.03.003
 55. Pender M. Ascending impairment of nociception in rats with experimental allergic encephalomyelitis. *J Neurol Sci* (1986) **75**:317–28. doi:10.1016/0022-510X(86)90079-1
 56. King T, Vera-Portocarrero L, Gutierrez T, Vanderah TW, Dussor G, Lai J, et al. Unmasking the tonic-aversive state in neuropathic pain. *Nat Neurosci* (2009) **12**:1364–6. doi:10.1038/nn.2407
 57. Sandyk R. Serotonergic neuronal sprouting as a potential mechanism of recovery in multiple sclerosis. *Int J Neurosci* (1999) **97**:131–8. doi:10.3109/00207459908994307
 58. Melanson M, Miao P, Eisenstat D, Gong Y, Gu X, Au K, et al. Experimental autoimmune encephalomyelitis-induced upregulation of tumor necrosis factor- α in the dorsal root ganglia. *Mult Scler* (2009) **15**:1135–45. doi:10.1177/1352458509106856
 59. Nicholson LB, Murtaza A, Hafner BP, Sette A, Kuchroo VK. A T cell receptor antagonist peptide induces T cells that mediate bystander suppression and prevent autoimmune encephalomyelitis induced with multiple myelin antigens. *Proc Natl Acad Sci U S A* (1997) **94**:9279–84. doi:10.1073/pnas.94.17.9279
 60. Austin PJ, Kim CF, Perera CJ, Moalem-Taylor G. Regulatory T cells attenuate neuropathic pain following peripheral nerve injury and experimental autoimmune neuritis. *Pain* (2012) **153**:1916–31. doi:10.1016/j.pain.2012.06.005
 61. Heijmans N, Smith PA, Morris-Downes MM, Pryce G, Baker D, Donaldson AV, et al. Encephalitogenic and tolerogenic potential of altered peptide ligands of MOG and PLP in Biozzi ABH mice. *J Neuroimmunol* (2005) **167**:23–33. doi:10.1016/j.jneuroim.2005.06.005
 62. Kappos L, Comi G, Panitch H, Oger J, Antel J, Conlon P, et al. Induction of a non-encephalitogenic type 2 T helper-cell autoimmune response in multiple sclerosis after administration of an altered peptide ligand in a placebo-controlled, randomized phase II trial. The Altered Peptide Ligand in Relapsing MS Study Group. *Nat Med* (2000) **6**:1176–82. doi:10.1038/80525
 63. Bielekova B, Goodwin B, Richert N, Cortese I, Kondo T, Afshar G, et al. Encephalitogenic potential of the myelin basic protein peptide (amino acids 83–99) in multiple sclerosis: results of a phase II clinical trial with an altered peptide ligand. *Nat Med* (2000) **6**:1167–75. doi:10.1038/80516
 64. Kim HJ, Antel J, Duquette P, All-eva DG, Conlon P, Bar-Or A. Persistence of immune responses to altered and native myelin antigens in patients with multiple sclerosis treated with altered peptide ligand. *Clin Immunol* (2002) **104**:105–14. doi:10.1006/clim.2002.5258
 65. Martin R, Sturzebecher C-S, McFarland HF. Immunotherapy of multiple sclerosis: where are we? Where should we go? *Nat Immunol* (2001) **2**:785–8. doi:10.1038/ni0901-785

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 16 July 2013; accepted: 16 October 2013; published online: 29 October 2013.

Citation: Tian DH, Perera CJ, Apostolopoulos V and Moalem-Taylor G (2013) Effects of vaccination with altered peptide ligand on chronic pain in experimental autoimmune encephalomyelitis, an animal model of multiple sclerosis. *Front. Neurol.* **4**:168. doi: 10.3389/fneur.2013.00168

This article was submitted to Multiple Sclerosis and Neuroimmunology, a section of the journal *Frontiers in Neurology*.

Copyright © 2013 Tian, Perera, Apostolopoulos and Moalem-Taylor. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.