



OPEN ACCESS

EDITED AND REVIEWED BY
Svetlana Lorenzano,
Sapienza University of Rome, Italy

*CORRESPONDENCE

Josefine Grundtvig
josefine.liv.gilling.grundtvig@regionh.dk
Hanne Christensen
hanne.krarup.christensen@regionh.dk

SPECIALTY SECTION

This article was submitted to
Neuroepidemiology,
a section of the journal
Frontiers in Neurology

RECEIVED 27 September 2022

ACCEPTED 05 October 2022

PUBLISHED 19 October 2022

CITATION

Grundtvig J, Ovesen C, Steiner T,
Carcel C, Gaist D, Christensen L,
Marstrand J, Meden P, Rosenbaum S,
Iversen HK, Kruuse C, Christensen T,
Ægidius K, Havsteen I and
Christensen H (2022) Corrigendum:
Sex-differences in oral
anticoagulant-related intracerebral
hemorrhage.
Front. Neurol. 13:1055360.
doi: 10.3389/fneur.2022.1055360

COPYRIGHT

© 2022 Grundtvig, Ovesen, Steiner,
Carcel, Gaist, Christensen, Marstrand,
Meden, Rosenbaum, Iversen, Kruuse,
Christensen, Ægidius, Havsteen and
Christensen. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Corrigendum: Sex-differences in oral anticoagulant-related intracerebral hemorrhage

Josefine Grundtvig^{1,2*}, Christian Ovesen¹, Thorsten Steiner^{2,3,4}, Cheryl Carcel⁵, David Gaist⁶, Louisa Christensen^{1,2}, Jacob Marstrand^{1,2}, Per Meden^{1,2}, Sverre Rosenbaum¹, Helle K. Iversen^{2,7}, Christina Kruuse^{2,8}, Thomas Christensen⁷, Karen Ægidius⁶, Inger Havsteen⁹ and Hanne Christensen^{1,2*}

¹Department of Neurology, Bispebjerg Hospital, Copenhagen, Denmark, ²Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark, ³Department of Neurology, Klinikum Frankfurt Höchst, Frankfurt, Germany, ⁴Department of Neurology, Heidelberg University Hospital, Heidelberg, Germany, ⁵The George Institute for Global Health, University of New South Wales, Sydney, NSW, Australia, ⁶Research Unit for Neurology, Odense University Hospital, University of Southern Denmark, Odense, Denmark, ⁷Department of Neurology, Rigshospitalet, Copenhagen, Denmark, ⁸Department of Neurology, Herlev Hospital, Copenhagen, Denmark, ⁹Department of Radiology, Bispebjerg Hospital, Copenhagen, Denmark

KEYWORDS

stroke, sex-differences, ICH, oral anticoagulation, vitamin K-antagonist, stroke in women, intracerebral hemorrhage (ICH), NOAC

A corrigendum on

[Sex-differences in oral anticoagulant-related intracerebral hemorrhage](#)

by Grundtvig, J., Ovesen, C., Steiner, T., Carcel, C., Gaist, D., Christensen, L., Marstrand, J., Meden, P., Rosenbaum, S., Iversen, H. K., Kruuse, C., Christensen, T., Ægidius, K., Havsteen, I., and Christensen, H. (2022). *Front. Neurol.* 13:832903. doi: 10.3389/fneur.2022.832903

In the published article, there was an error in [Table 1](#) as published due to erroneous data entry. The reported NIHSS in the table for the male participants is 14 (10; 15) and it is 14 (8; 15) for the female, $p = 0.88$. The correct number for the male participants is 8 (4; 13) and for the female it is 10 (5; 15), $p = 0.09$. The corrected [Table 1](#) and its caption appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

TABLE 1 Baseline data in 401 men and women with oral anticoagulant (OAC)-related intracerebral hemorrhage (ICH).

	Male (n = 226)	Female (n = 175)	P-value
Mean age (SD)	76.9 (8.3)	81.9 (7.4)	<0.001
Modified Rankin Scale	0 (0; 1)	1 (0; 2)	<0.001
Comorbidity scores			
Median Charlson Comorbidity Index (IQR)	1 (0; 3)	1 (0; 2)	0.62
Median CHA ₂ DS ₂ -VASc (IQR)	2 (1; 3)	2 (1; 3)	0.74
Medicine			
NOAC	60 (26.5 %)	69 (39.4 %)	0.009
VKA	166 (73.5 %)	106 (60.6 %)	
Antiplatelets	40 (18.6 %)	16 (9.9 %)	0.03
Antihypertensives	156 (71.9 %)	123 (75.9 %)	0.44
Lipid lowering agents	94 (43.9 %)	48 (29.6 %)	0.006
Selective serotonin reuptake inhibitors	14 (6.5 %)	20 (12.3 %)	0.08
Smoking status			
Active smoker	22 (13.1 %)	10 (7.8 %)	<0.001
Former smoker	80 (47.6 %)	37 (28.7 %)	
Never smoker	66 (39.3 %)	82 (63.6 %)	
Alcohol/ drug use			
No alcohol use	42 (25.5 %)	54 (44.3 %)	<0.001
Below 14 units per week	89 (53.9 %)	63 (51.6 %)	
Above 14 units per week	34 (20.6 %)	5 (4.1 %)	
Alcohol dependency	20 (8.8 %)	2 (1.1 %)	0.002
Stroke severity scores			
Admission Glasgow Coma Scale	14 (10; 15)	14 (8; 15)	0.03
Admission NIHSS	8 (4; 13)	10 (5; 15)	0.09
Systolic blood pressure limit order, n (%) ^a			
<140 mmHg	53 (23.5 %)	42 (24.0 %)	0.99
140–160 mmHg	27 (11.9 %)	21 (12.0 %)	
160–180 mmHg	4 (1.8 %)	2 (1.1 %)	
No limit ordered	142 (62.8 %)	110 (62.9 %)	
Radiology			
Median hematoma volume, mL (IQR)	22.2 (4.6; 64.6)	19.1 (4.6; 63.8)	0.90

OAC, oral anticoagulant; ICH, intracerebral hemorrhage; SD, standard deviation; IQR, interquartile range; NIHSS, National Institute of Health Stroke Scale; mmHg, millimeter of mercury.

^aNumber of patients with a systolic blood pressure limit order in the patient file.