



OPEN ACCESS

EDITED AND REVIEWED BY
Jo Madeleine Wilmshurst,
University of Cape Town, South Africa

*CORRESPONDENCE
Divyen K. Shah
✉ d.shah@qmul.ac.uk

†Joint senior authors

RECEIVED 01 August 2024
ACCEPTED 13 August 2024
PUBLISHED 30 September 2024

CITATION
Shah DK, Ponnusamy V, Evanson J,
Kapellou O, Ekitzidou G, Gupta N, Clarke P,
Michael-Titus AT and Yip PK (2024)
Corrigendum: Raised plasma neurofilament
light protein levels are associated with
abnormal MRI outcomes in newborns
undergoing therapeutic hypothermia.
Front. Neurol. 15:1474489.
doi: 10.3389/fneur.2024.1474489

COPYRIGHT
© 2024 Shah, Ponnusamy, Evanson, Kapellou,
Ekitzidou, Gupta, Clarke, Michael-Titus and
Yip. This is an open-access article distributed
under the terms of the [Creative Commons
Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other forums is
permitted, provided the original author(s) and
the copyright owner(s) are credited and that
the original publication in this journal is cited,
in accordance with accepted academic
practice. No use, distribution or reproduction
is permitted which does not comply with
these terms.

Corrigendum: Raised plasma neurofilament light protein levels are associated with abnormal MRI outcomes in newborns undergoing therapeutic hypothermia

Divyen K. Shah^{1,2*}, Vennila Ponnusamy^{3,4}, Jane Evanson¹,
Olga Kapellou⁵, Georgia Ekitzidou⁵, Neelam Gupta⁶,
Paul Clarke⁷, Adina T. Michael-Titus^{1†} and Ping K. Yip^{1†}

¹The Royal London Hospital, Barts Health NHS Trust, London, United Kingdom, ²The Centre for Neuroscience and Trauma, Barts and The London School of Medicine and Dentistry, Blizard Institute, Queen Mary University of London, London, United Kingdom, ³Centre for Genomics and Child Health, Barts and The London School of Medicine and Dentistry, Blizard Institute, Queen Mary University of London, London, United Kingdom, ⁴Ashford and St. Peter's Hospitals NHS Foundation Trust, Chertsey, United Kingdom, ⁵Homerton University Hospital NHS Foundation Trust, London, United Kingdom, ⁶University Hospital Southampton, Southampton, United Kingdom, ⁷Norfolk and Norwich University Hospitals NHS Foundation Trust, Norwich, United Kingdom

KEYWORDS

neurofilament proteins, hypoxic-ischemic encephalopathy, therapeutic hypothermia, MRI imaging, neuroprotection, biomarkers

A Corrigendum on

[Raised plasma neurofilament light protein levels are associated with abnormal MRI outcomes in newborns undergoing therapeutic hypothermia](#)

by Shah, D. K., Ponnusamy, V., Evanson, J., Kapellou, O., Ekitzidou, G., Gupta, N., Clarke, P., Michael-Titus, A. T., and Yip, P. K. (2018). *Front. Neurol.* 9:86. doi: 10.3389/fneur.2018.00086

In the published article, there was an error in [Table 2](#) as published. For patients 1 and 3, the cell for the column entitled “Neurologic exam prior to discharge” should read “Severe, died after 18 months” instead of “Severe, died.” The corrected [Table 2](#) and its caption “Patients arranged in descending rank order of neurofilament light levels at S3 in relation to MRI findings and neurological exam at discharge. The latter is classified primarily in terms of abnormality of tone” appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

TABLE 2 Patients arranged in descending rank order of neurofilament light levels at S3 in relation to MRI findings and neurological exam at discharge. The latter is classified primarily in terms of abnormality of tone.

	S1 (pg/mL)	S2 (pg/mL)	S3 (pg/mL)	Age at peak (h)	Electrical seizures (Y/N)	PLIC	BGT	WM	Cortex	Composite MRI Group	Sentinel event (Y/N)	Comment	Neurologic exam prior to discharge
1	2,506	2,255	4,698	100	Y	Loss	Severe	Moderate	Normal	U	N		Severe, died after 18 months
2	1,999	1,615	1,993	18	N	Equivocal	Normal	Severe	Moderate	U	Y	Cord prolapse	Mild
3	0	189	1,626	124	Y	Loss	Severe	Severe	Severe	U	N		Severe, died after 18 months
4	70	423	1,014	114	N	Loss	Severe	Mild	Mild	U	Y	Breech, head entrapment	Moderate, tube feeds
5	1,046	1,512	874	31	Y	Normal	Moderate	Normal	Normal	U	N		Normal
6	516	n/a	849	96	N	Equivocal	Normal	Severe	Severe	U	Y	Placental abruption	Severe, tube feeds
7	413	381	844	118	Y	Normal	Moderate	Severe	Moderate	U	N		Moderate, tube feeds
8	29	225	674	95	N	Loss	Severe	Severe	Severe	U	Y	Shoulder dystocia	Severe, tube feeds
9	0	n/a	563	94	Y	Equivocal	Moderate	Mild	Moderate	U	N		Moderate, tube feeds
10	0	0	511	96	n/a	Loss	Moderate	Mild	Normal	U	N		Severe, tube feeds
11	0	294	506	100	Y	Normal	Mild	Mild	Normal	F	N		Normal
12	681	419	490	18	N	Loss	Severe	Normal	Moderate	U	N		Normal
13	141	321	403	115	N	Normal	Normal	Normal	Normal	F	Y	Placental abruption	Normal
14	99	577	381	67	N	Normal	Normal	Normal	Normal	F	Y	Head entrapment	Normal
15	28	77	336	101	N	Normal	Normal	Mild	Mild	F	Y	Shoulder dystocia	Normal
16	173	159	334	101	N	Loss	Severe	Moderate	Mild	U	N		Normal
17	277	457	324	62	N	Normal	Normal	Normal	Normal	F	N		Mild
18	166	160	242	86	Y	Normal	Mild	Normal	Normal	F	N		Normal
19	0	71	164	108	n/a	Normal	Normal	Mild	Normal	F	N		Normal

(Continued)

TABLE 2 (Continued)

	S1 (pg/mL)	S2 (pg/mL)	S3 (pg/mL)	Age at peak (h)	Electrical seizures (Y/N)	PLIC	BGT	WM	Cortex	Composite MRI Group	Sentinel event (Y/N)	Comment	Neurologic exam prior to discharge
20	148	172	151	48	Y	Loss	Moderate	Severe	Normal	U	N		Moderate, tube feeds
21	0	41	143	115	n/a	Normal	Normal	Normal	Normal	F	N		Normal
22	0	0	136	114	Y	Normal	Normal	Moderate	Normal	F	N		Normal
23	0	0	59	102	N	Normal	Normal	Normal	Normal	F	N		Normal
24	0	0	8	-	n/a	Normal	Normal	Normal	Normal	F	N		Normal
25	0	0	0	-	Y	Normal	Normal	Mild	Mild	F	N		Normal
26	0	0	0	-	N	Normal	Normal	Normal	Normal	F	Y	Uterine rupture	Normal

PLIC, posterior limb of internal capsule; BGT, basal ganglia and thalami; WM, white matter; n/a, sample not available; F, favorable; U, unfavorable; y, yes; n, no.